

THE ISOTROPIC CONSTANT CONJECTURE IS TRUE

J.ASPEGREN

ABSTRACT. In this preprint we will prove the isotropic constant conjecture.

1. INTRODUCTION

All the problems that we investigate in this preprint have their roots in classical mechanics. However, our tools are from the asymptotic convex geometry. We will prove the following theorem.

Theorem 1. *Let the convex body $D \subset \mathbf{R}^n$ be isotropic. Then*

$$\int_D \|x\|^2 dx < nC,$$

where $C > 0$, where C does not depend on n .

Above result implies the hyperplane conjecture [1][3]. Thus, any convex body of volume 1 has a $n - 1$ dimensional hyperplane section that has volume greater than some universal constant [2].

2. DEFINITIONS AND KNOWN RESULTS

A convex body $|D| = 1$ is isotropic if the barycenter is at the origin and

$$\int_D \langle y, x \rangle^2 dx = \alpha^2 \|y\|^2,$$

for some constant $\alpha > 0$ and all $y \in \mathbf{R}^n$. The following integral can be calculated in spherical coordinates.

$$(1) \quad \int_{B_n} \|x\|^2 dx = \frac{n}{n+2} |B_n|,$$

where B_n is the unit ball.

3. THE PROOF OF THE MAIN THEOREM

Let D be an isotropic convex body. Because centralized balls are isotropic then via scaling there exists a ball L with the same isotropic constant α . So the real question is the volume of that ball L . Now, we integrate the following in spherical coordinates and obtain

$$(2) \quad \frac{1}{|L|^{(n+2)/n}} \int_D \|x\|^2 dx = \frac{1}{|L|^{(n+2)/n}} \int_L \|x\|^2 dx = \frac{1}{|L|^{(n+2)/n}} \frac{nR^{n+2}}{n+2} |B_n| \\ = \frac{n}{n+2} |B|^{-2/n},$$

1991 *Mathematics Subject Classification.* 52A20,52A23.

Key words and phrases. Convex Geometry, Bourgain's Slicing Problem, Hyperplane Conjecture, Asymptotic Convex Geometry.

where B_n is the n dimensional unit ball and we used scaling in order to rewrite the expression. Rewriting the above gives

$$(3) \quad \frac{1}{|L|} \int_L ||x||^2 dx = \frac{n}{n+2} |B_n|^{-2/n} |L|^{2/n} < c_n |B_n|^{-2/n},$$

where $c_n > 0$ and

$$(4) \quad \lim_{n \rightarrow \infty} c_n = 1.$$

The bound follows when we consider the bound

$$(5) \quad |L| \leq cn.$$

For the bound (5) see for example [3] and [2]. Now, it follows from (3) that

$$\sqrt{\frac{n}{n+2}} R < \sqrt{c_n} |B_n|^{-1/n} = \sqrt{c_n} r,$$

where r is the radius of the ball with the unit volume. Let $\epsilon > 0$ then there exists $n \geq 1$ s.t

$$R < r + \epsilon$$

Thus,

$$|L|^{1/n} = |B_n|^{1/n} R = r^{-1} * R < \frac{(r + \epsilon)}{r}$$

So it follows that

$$|L| < \left(\frac{r + \epsilon}{r}\right)^n < \left(\frac{r + 1}{r}\right)^n$$

Thus,

$$(6) \quad |L| < C_1$$

and we have theorem 1.

REFERENCES

- [1] J.Bourgain, *On high-dimensional maximal functions associated to convex bodies*, Amer. J. Math., Vol. 108, No. 6, (1986), 1467–1476.
- [2] A.Giannopoulos, *Notes on isotropic convex bodies*, Warsaw, October 2003.
- [3] V.D.Milman and A. Pajor *Isotropic Position and Inertia Ellipsoids and Zonoids of the Unit Ball of a Normed n -Dimensional Space*, GAFA-Seminar 87-88, Springer Lecture Notes in Math., v.1376, 64-104, (1989).

Email address: jaspegren@outlook.com