

THE KAKEYA SET CONJECTURE IS TRUE

J. ASPEGREN

ABSTRACT. In this article we will prove the Kakeya set conjecture. In addition we will prove that in the usual approach to the Kakeya maximal function conjecture we can assume that the tube-sets are maximal. Third, we build a direct connection between line incidence theorems and Kakeya type conjectures.

1. INTRODUCTION

The Kakeya maximal function conjecture and its variations have gained considerable interest especially after an influential paper by Bourgain (?). For example, it would follow from the conjecture that the Kakeya sets and the Nikodym sets have necessarily full dimensions (11; 12; 8). However, the Nikodym set conjecture is implied by the Kakeya set conjecture (8; 12). The case $n = 2$ was proved by Davies see (5) and the finite field case by Dvir (6). A Kakeya is a set that contains a unit line in every direction. For surveys see (16; 13; 3). Almost all the necessary preliminaries for this paper can be found for example in (8), (11) and in (14). Define the δ -tubes in standard way: for all $\delta > 0, \omega \in S^{n-1}$ and $a \in \mathbb{R}^n$, let

$$T_\omega^\delta(a) = \{x \in \mathbb{R}^n : |(x-a) \cdot \omega| \leq \frac{\delta}{2}, |proj_{\omega^\perp}(x-a)| \leq \delta\}.$$

Moreover, let $f \in L^1_{loc}(\mathbb{R}^n)$. Define the Kakeya maximal function $f_\delta^* : S^{n-1} \rightarrow \mathbb{R}$ via

$$f_\delta^*(\omega) = \sup_{a \in \mathbb{R}^n} \frac{1}{|T_\omega^\delta(a)|} \int_{T_\omega^\delta(a)} |f(y)| dy.$$

In this paper any constant can depend on dimension n . In study of the Kakeya maximal function conjecture we are aiming at the following bounds

$$(1) \quad \|f_\delta^*\|_p \leq C_\epsilon \delta^{-n/p+1-\epsilon},$$

for all $\epsilon > 0$. Remarkably, a bound of the form (1) follows from a bound of the form

$$(2) \quad \left\| \sum_{\omega \in \Omega} 1_{T_\omega(a_\omega)} \right\|_{p/(p-1)} \leq C_\epsilon \delta^{-n/p+1-\epsilon},$$

for all $\epsilon > 0$, and for any set of δ -separated δ -tubes. See for example (12) or (8). We will prove that we need to consider only the case where the set Ω is maximal. As usual we define that $A \lesssim B$ iff for all $\epsilon > 0$ and for all $\delta > 0$, it holds that $A \leq C_\epsilon \delta^{-\epsilon} B$. We will prove that

Theorem 1. *Every Kakeya set has full dimension.*

1991 *Mathematics Subject Classification.* 42B37, 28A75.

Key words and phrases. Kakeya conjectures.

2. A REDUCTION TO THE CASE WHERE THE TUBE-SETS ARE MAXIMAL

Let Ω' be any set of δ -separated directions. We will prove that

$$\left\| \sum_{\omega' \in \Omega'} 1_{T_{\omega'}(a_{\omega'})} \right\|_{p/(p-1)} \leq \left\| \sum_{\omega \in \Omega} 1_{T_{\omega}(a_{\omega})} \right\|_{p/(p-1)},$$

where Ω is maximal. We construct the set Ω as follows. Let Ω' be the original direction-set and let $\Omega' \subset \Omega'''$ be maximal. Define

$$\Omega'' := \{\omega'' \in S^{n-1} \mid \omega'' \in \Omega''' / \{\Omega'\}\}.$$

Moreover, let

$$\Omega := \Omega' \cup \Omega''.$$

Clearly, Ω is maximal. We choose the tubes corresponding to directions in Ω' to have origo as their center of masses. Thus, what we do is that we add tubes to the original tube-set so it becomes maximal. Now, we can estimate:

$$\begin{aligned} \left\| \sum_{\omega' \in \Omega'} 1_{T_{\omega'}(a_{\omega'})} \right\|_{p/(p-1)} &\leq \left\| \sum_{\omega' \in \Omega'} 1_{T_{\omega'}(a_{\omega'})} + \sum_{\omega'' \in \Omega''} 1_{T_{\omega''}(0)} \right\|_{p/(p-1)} \\ &= \left\| \sum_{\omega \in \Omega} 1_{T_{\omega}(a_{\omega})} \right\|_{p/(p-1)}. \end{aligned}$$

Thus, we need only to consider the cases where the tube sets are maximal.

3. PREVIOUSLY KNOWN RESULTS

We will use the following bound for the pairwise intersections of δ -tubes:

Lemma 1 (Corbòda). *For any pair of directions $\omega_i, \omega_j \in S^{n-1}$ and any pair of points $a, b \in \mathbb{R}^n$, we have*

$$|T_{\omega_i}^{\delta}(a) \cap T_{\omega_j}^{\delta}(b)| \lesssim \frac{\delta^n}{|\omega_i - \omega_j|}.$$

A proof can be found for example in (8).

For any (spherical) cap $\Omega \subset S^{n-1}$, $|\Omega| \gtrsim \delta^{n-1}$, $\delta > 0$, denote its δ -entropy $N_{\delta}(\Omega)$ as the maximum possible cardinality for an δ -separated subset of Ω .

Lemma 2. *In the notation just defined*

$$1 \leq N_{\delta}(\Omega) \sim \frac{|\Omega|}{\delta^{n-1}}.$$

Again, a proof can essentially be found in (8).

4. A LINE INTERSECTION THEOREM

We consider a set of lines L where no two lines belong to a common plane containing say, A , and no line intersects two or more strict joints. We call any intersection of such lines a strict joint. We will prove a bound for the number of such line intersections.

Theorem 2. *The number of strict joints $|S_k|$ of order k of n lines is less than n/k .*

Proof. We see this by noticing that A , a strict joint B and some non-intersection point of a line C intersecting a strict joint B defines a plane. No strict joint belongs to 2 or more planes by assumption. It's clear from the assumptions that those planes are distinct. To each strict joints there corresponds k unique planes. In total we have n planes. So the number of strict joints is less than n/k . \square

5. A PROOF OF THE KAKEYA SETL FUNCTION CONJECTURE

We consider a maximal δ -separated set of middle line intersecting tubes $E \subset K_\delta$, where K_δ is a δ -neighbourhood of a Keakeya set. We can assume that no two central lines of the tubes belong to a same plane going through origo by rotating the tubes if necessary a small amount. This can only increase the number of intersections of positive volume and does not affect too much to their volumes. So via theorem 2 of the last section, the number of central line intersections is less than $\sim \delta^{1-n}$. Next, we prove a lemma.

Lemma 3. *It holds that*

$$(3) \quad \left| \bigcap_{i=1}^{2^k} T_i \right| \lesssim \delta^{n-1} 2^{-k/(n-1)}.$$

Proof. Let us suppose that $2^k \sim \delta^{-\beta}$, $0 < \beta \leq n-1$, and let's suppose that tube $T_{\omega'}$ intersecting $T_\omega \cap E_{2^k}$ has it's direction outside of a cap of size $\sim \delta^{n-1-\beta}$ on the unit sphere. Then the angle between T_ω and $T_{\omega'}$ is greater than $\sim \delta^{1-\beta/(n-1)}$. Thus by lemma 1 the intersection

$$(4) \quad \left| \bigcap_{i=1}^{2^k} T_i \right| \leq |T_\omega \cap T_{\omega'} \cap E_{2^k}| \leq |T_\omega \cap T_{\omega'}| \lesssim \delta^{n-1+\beta/(n-1)} \sim \delta^{n-1} 2^{-k/(n-1)}.$$

Thus, we can suppose that the directions in the intersection $E_{2^k} \cap T_\omega \cap T_{\omega'}$ belong to a cap of size $\sim \delta^{n-1+\beta}$. If we δ -separate the cap via lemma 2 we get that the cap can contain at most $\sim 2^k$ tube-directions. Thus, for any tube T_ω in the intersection there exists a tube $T_{\omega'}$, such that the angle the angle between T_ω and $T_{\omega'}$ is $\sim \delta^{1-\beta/(n-1)}$ and the inequality (4) is valid. \square

Now, there exists dyadic k such that

$$(5) \quad 2^k |E_{2^k}| \approx 1.$$

Moreover there exists $\delta_1 \leq \delta$ s.t all the intersections are central line intersections. For that k' it holds that

$$2^{k'} |E_{2^{k'}}| \approx \delta_1^{n-1} \delta^{1-n},$$

in order that equation (5) holds we can assume that $k = k'$. So we replace the δ -tubes with δ_1 -tubes and obtain for the same k via scaling that

$$(6) \quad \delta_1^{n-1} \delta^{1-n} \approx 2^k |E'_{2^k}| \lesssim |S'_k| \delta_1^{n-1} 2^{-k/(n-1)} \lesssim 2^{-k/(n-1)} \delta_1^{n-1} \delta^{1-n}.$$

Thus,

$$(7) \quad 2^k \approx 1.$$

Now, above (7) and (5) implies that

$$(8) \quad |E_{2^k}| \approx 1.$$

The above (8) implies easily the Minkowski version of the Keakeya set onjecture and the Hausdorff version follows from (1).

REFERENCES

- [1] J.Aspegren, *The Kakeya Tube Conjecture Implies the Kakeya Conjecture*, available at <http://vixra.org/pdf/1709.0357v2.pdf>
- [2] J. Bourgain, *Besicovitch Type Maximal Operators and Applications to Fourier Analysis*, *Geometric and Functional Analysis 1* (1991), 147-187.
- [3] J. Bourgain, *Harmonic analysis and combinatorics: How much may they contribute to each other?*,IMU/Amer. Math. Soc. (2000), 13–32
- [4] A. Córdoba, *The Kakeya Maximal Function and the Spherical Summation Multipliers*, *American Journal of Mathematics* 99 (1977), 1-22.
- [5] R.O. Davies, *Some Remarks on the Kakeya Problem*, *Proc. Camb. Phil. Soc.* 69 (1971), 417-421.
- [6] Z. Dvir, *On the Size of Kakeya Sets in Finite Fields*, *J. Amer. Math. Soc.* 22 (2009), 1093-1097.
- [7] K.J. Falconer *The Geometry of Fractal Sets*, Cambridge University Press (1985).
- [8] E.Kroc, *The Kakeya problem*, available at <http://ekroc.weebly.com/uploads/2/1/6/3/21633182/mscessay-final.pdf>
- [9] W. Rudin, *Real and Complex Analysis*, McGraw-Hill Education (1986)
- [10] E. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton University Press (1993)
- [11] T. Tao, *Lecture Notes*, available at math.ucla.edu/~tao/254b.1.99s/ (1999)
- [12] T. Tao, *The Bochner-Riesz Conjecture Implies the Restriction Conjecture*, *Duke Math. J.* 96 (1999), 363-375.
- [13] T.Tao, *From rotating needles to stability of waves: emerging connections between combinatorics, analysis, and pde*,*Notices Amer. Math. Soc.*, 48(3),(2001),294–303.
- [14] T. Tao <https://terrytao.wordpress.com/2009/05/15/the-two-ends-reduction-for-the-kakeya-maximal-conjecture/>
- [15] T. Wolff, *An Improved Bound for Kakeya Type Maximal Functions*, *Rev. Mat. Iberoamericana* 11 (1995), 651-674.
- [16] T. Wolff, *Recent work connected with the Kakeya problem* *Prospects in mathematics* (Princeton, NJ, 1996), (1999),129–162.

Email address: jaspegren@outlook.com