

**Two simples proofs of Fermat 's last theorem and Beal conjecture**

31 Octobre 2018

M.Sghiar

msghiar21@gmail.com

9 Allée capitaine Jean Bernard Bossu, 21240, Talant, France

**Abstract** :If after 374 years the famous theorem of Fermat-Wiles was demonstrated in 150 pages by A. Wiles [2], the purpose of this article is to give a simple demonstration and deduce a proof of the Beal conjecture.

**Résumé** : Si après 374 ans le célèbre théorème de Fermat-Wiles a été démontré en 150 pages par A. Wiles [2], le but de cet article est de donner une simple démonstration et d'en déduire une preuve de la conjecture de Beal.

**Keywords** : Fermat, Fermat-Wiles theorem, Fermat's great theorem.

**The Subject Classification Codes** : 11D41 - 11G05 - 11G07 - 26B15 - 26B20 - 28A10 - 28A75 -

**1-Introduction :**

Set out by Pierre de Fermat [2], it was not until more than three centuries ago that Fermat's great theorem was published, validated and established by the British mathematician Andrew Wiles [4] in 1995.

In mathematics, and more precisely in number theory, the last theorem of Fermat [2], or Fermat's great theorem, or since his Fermat-Wiles theorem demonstration [3], is as follows : There are no non-zero integers a, b, and c such that :  $a^n + b^n = c^n$  , as soon as n is an integer strictly greater than 2 ".

The Beal conjecture is the following conjecture in number theory : If  $a^x + b^y = c^z$  where a, b, c, x, y and z are positive integers with x, y, z > 2, then a, b, and c have a common prime factor. Equivalently, There are no solutions to the above equation in positive integers a, b, c, x, y, z with a, b and c being pairwise coprime and all of x, y, z being greater than 2.

If the famous Fermat-Wiles theorem has been demonstrated in **150 pages** by A. Wiles [4], the purpose of this article is to give a simple proof and deduce a proof of the Beal conjecture.

## 2. The proof of Fermat 's last theorem

**Théorème :**

There are no non-zero integers a, b, and c such that:  $a^n + b^n = c^n$  , with n an integer strictly greater than 2

**Lemma 1 :**

If n, a, b and c are a non-zero integers with and  $a^n + b^n = c^n$  then:

$$\int_0^b x^{n-1} - \left( \frac{c-a}{b}x + a \right)^{n-1} \frac{c-a}{b} dx = 0$$

Proof :

$$a^n + b^n = c^n \Leftrightarrow \int_0^a n x^{n-1} dx + \int_0^b n x^{n-1} dx = \int_0^c n x^{n-1} dx$$

But as :

$$\int_0^c n x^{n-1} dx = \int_0^a n x^{n-1} dx + \int_a^c n x^{n-1} dx$$

So :

$$\int_0^b n x^{n-1} dx = \int_a^c n x^{n-1} dx$$

And as by changing variables we have :

$$\int_a^c n x^{n-1} dx = \int_0^b n \left( \frac{c-a}{b}y + a \right)^{n-1} \frac{c-a}{b} dy$$

Then :

$$\int_0^b x^{n-1} dx = \int_0^b \left( \frac{c-a}{b} y + a \right)^{n-1} \frac{c-a}{b} dy$$

It results:

$$\int_0^b x^{n-1} - \left( \frac{c-a}{b} x + a \right)^{n-1} \frac{c-a}{b} dx = 0$$

**Corollary 1 :** If  $N, n, a, b$  and  $c$  are a non-zero integers with and  $a^n + b^n = c^n$  then :

$$\int_0^{\frac{b}{N}} x^{n-1} - \left( \frac{c-a}{b} x + \frac{a}{N} \right)^{n-1} \frac{c-a}{b} dx = 0$$

**Proof :** It results from the proof of lemma 1 by replacing  $a, b$  and  $c$  respectively by  $\frac{a}{N}$ ,  $\frac{b}{N}$  and  $\frac{c}{N}$ .

**Lemma 2 :**

If  $a^n + b^n = c^n$ , where  $n, a, b$  and  $c$  are a non-zero integers with  $n > 2$  and  $a \leq b \leq c$ . Then for an integer  $N$  big enough we have :

$$f(x, a, b, c, N) = x^{n-1} - \left( \frac{c-a}{b} x + \frac{a}{N} \right)^{n-1} \frac{c-a}{b} \leq 0 \quad \forall x \in \left[ 0, \frac{b}{N} \right]$$

**Proof :**

We have :  $\frac{\partial f}{\partial x} = (n-1)x^{n-2} - (n-1) \left( \frac{c-a}{b} x + \frac{a}{N} \right)^{n-2} \left( \frac{c-a}{b} \right)$ ,  $f(0, a, b, c, N) < 0$  and

$\frac{\partial f}{\partial x} \Big|_{x=0} < 0$  so, by continuity,  $\forall y \in \left[ 0, \frac{b}{N} \right]$  with  $N$  an integer big enough, we have

$\frac{\partial f}{\partial x} \Big|_{x=y} < 0$ . So the function  $f$  is decreasing in  $\left[ 0, \frac{b}{N} \right]$  and we have :

$$f(x, a, b, c, N) \leq 0 \quad \forall x \in \left[ 0, \frac{b}{N} \right]$$

**Proof of Theorem:**

If  $a^n + b^n = c^n$ , where  $n, a, b$  and  $c$  are a non-zero integers with  $n > 2$  and  $a \leq b \leq c$ . Then for an integer  $N$  big enough, it results from the **lemma 2** that we have :

$$f(x, a, b, c, N) = x^{n-1} - \left( \frac{c-a}{b} x + \frac{a}{N} \right)^{n-1} \frac{c-a}{b} \leq 0 \quad \forall x \in \left[ 0, \frac{b}{N} \right]$$

And by using the **corollary 1**, we have  $\int_0^{\frac{b}{N}} x^{n-1} - \left( \frac{c-a}{b}x + \frac{a}{N} \right)^{n-1} \frac{c-a}{b} dx = 0$  .

So :  $x^{n-1} - \left( \frac{c-a}{b}x + \frac{a}{N} \right)^{n-1} \frac{c-a}{b} = 0 \quad \forall x \in \left[ 0, \frac{b}{N} \right]$

And therefore  $\frac{c-a}{b} = 1$  because  $f(x, a, b, c, N)$  is a null polynomial as it have more than  $n$  zeros. So  $c = a + b$  and  $a^n + b^n \neq c^n$  which is absurde .

### 3- The proof of Beal conjecture :

#### Corollary : [Beal conjecture]

If  $a^x + b^y = c^z$  where  $a, b, c, x, y$  and  $z$  are positive integers with  $x, y, z > 2$ , then  $a, b$ , and  $c$  have a common prime factor.

Equivalently, there are no solutions to the above equation in positive integers  $a, b, c, x, y, z$  with  $a, b$  and  $c$  being pairwise coprime and all of  $x, y, z$  being greater than 2.

#### Proof :

Let  $a^x + b^y = c^z$

If  $a, b$  and  $c$  are not pairwise coprime, then by posing  $a = ka'$  ,  $b = kb'$  , and  $c = kc'$  .

Let  $a' = u'^{yz}$  ,  $b' = v'^{xz}$  ,  $c' = w'^{xy}$  and  $k = u'^{yz}$  ,  $k = v'^{xz}$  ,  $k = w'^{xy}$

As  $a^x + b^y = c^z$  , we deduce that  $(uu')^{xyz} + (vv')^{xyz} = (ww')^{xyz}$  .

So :  $k^x u'^{xyz} + k^y v'^{xyz} = k^z w'^{xyz}$

This equation does not look like the one studied in the first theorem. But if  $a, b$  and  $c$  are pairwise coprime, we have  $k = 1$  and  $u = v = w = 1$  and we will have to solve the equation :

$$u'^{xyz} + v'^{xyz} = w'^{xyz}$$

The equation  $u'^{xyz} + v'^{xyz} = w'^{xyz}$  have a solution if at least one of the equations :

$$(u'^{xy})^z + (v'^{xy})^z = (w'^{xy})^z , (u'^{xz})^y + (v'^{xz})^y = (w'^{xz})^y , (u'^{yz})^x + (v'^{yz})^x = (w'^{yz})^x , \text{ have a solution .}$$

So by the proof given in the proof of the first Theorem we must have :  $z \leq 2$  or  $y \leq 2$  , or  $x \leq 2$  .

We therefore conclude that if  $a^x + b^y = c^z$  where  $a, b, c, x, y$ , and  $z$  are positive integers with  $x, y, z > 2$  , then  $a, b$ , and  $c$  have a common prime factor.

#### 4- Important notes :

1- If a, b, and c are not pairwise coprime, someone, by applying the proof given in the corollary like this :  $a=u^{yz}$  ,  $b=v^{xz}$  ,  $c=w^{xy}$  will have  $u^{xyz}+v^{xyz}=w^{xyz}$  , and could say that all the x,y and z are always smaller than 2. What is false:  $7^3+7^4=14^3$  .

The reason is sipmle: it is the common factor k which could increase the power, for example if  $k=c^{r'}$  in the proof, then  $c^z=(kc')^z=c^{(r+1)z}$  . You can take the example :  $2^r+2^r=2^{r+1}$  where  $k=2^r$  .

2- These techniques do not say that the equation  $a^n+b^n=c^n$  where  $a,b,c \in \mathbb{R} \cap ]0,+\infty[$  has no solution since in the proof the equation  $X^2+Y^2=Z^2$  can have a sloution. We take  $a=X^{\frac{2}{n}}$  ,  $b=Y^{\frac{2}{n}}$  and  $C=Z^{\frac{2}{n}}$  .

3 – In [3] I proved the abc conjecture which implies only that the equation  $a^x+b^y=c^z$  has only a finite number of solution with a, b, c, x, y, z a positive integers, a, b and c being pairwise coprime and all of x, y, z being greater than 2.

#### 5- Conclusion :

The techniques used in this article have allowed to prove both the Fermat' last theorem and the Beal' conjecture and have shown that the Beal conjecture is only a corollary of the Fermat' last theorem.

#### Bibliography :

[1] [https://en.wikipedia.org/wiki/Beal\\_conjecture](https://en.wikipedia.org/wiki/Beal_conjecture)

[2] [https://en.wikipedia.org/wiki/Fermat\\_last\\_theorem](https://en.wikipedia.org/wiki/Fermat_last_theorem)

[3] M. Mghiar , la preuve de la conjecture abc, iosr journal of mathematics (iosr-jm), e-issn: 2278-5728, p-issn: 2319-765x. volume 14, issue 4 ver. i (jul - aug 2018), pp 22-26.

[4] Andrew Wiles, Modular elliptic curves and Fermat's last Théorème, Annal of mathematics, volume 10,142, pages 443-551, september-december, 1995.