

Estimation and Prediction of Neutrino Mass Based on the Kinetic Theory of Gases

Hejie Lin¹ and Tsung-Wu Lin²

¹*Department of Mechanical Engineering,
Oakland University, Rochester Hills, Michigan, USA*

²*Department of Civil Engineering,
National Taiwan University, Taipei, Taiwan*

Abstract

Neutrinos and molecules co-exist in space. Because both have physical properties of mass and speed, it is a logical assumption that neutrinos can interact with molecules according to the kinetic theory of gases. If neutrinos interact with gases such as nitrogen and oxygen, the mass of neutrino can be estimated using the kinetic theory of gases. According to the kinetic theory of gases, the estimated mass of a neutrino is 6.27×10^{-38} kg or $0.0352 \text{ eV}/c^2$. The goal of this estimation is to predict an approximate mass of neutrinos.

1. Introduction

Since 1998, when the Super-Kamiokande neutrino detector discovered the non-zero mass of neutrinos [1], there have been great efforts to measure the exact mass of neutrinos by various groups, including but not limited to, KamLAND [2,3], MINOS [4], and OPERA [5]. Instead of measuring the exact mass of neutrinos, we can estimate the mass of neutrinos based on the kinetic theory of gases.

Neutrinos and molecules co-exist in space. Because both have physical properties of mass and speed, it is a logical assumption that neutrinos can interact with molecules according to the kinetic theory of gases. If neutrinos interact with gases such as nitrogen and oxygen, the mass of a neutrino can be estimated using the kinetic theory of gases.

For this calculation, oxygen molecules and neutrino particles are assumed to be interacting in a well-defined closed environment. Based on the kinetic theory of gases, the RMS (root mean square) speeds of oxygen molecules and neutrino particles are used.

Using the given approximate information,

u_{oxygen} : RMS speed of oxygen molecules at 20 °C, 1 atm (326 m/s)

m_{oxygen} : mass of one oxygen molecule (5.31×10^{-26} kg)

$u_{neutrino}$: RMS speed of neutrinos (speed of light: 3.00×10^8 m/s)

the mass of a neutrino can be estimated using the following mass-speed relationship.


$$m_{oxygen}(u_{oxygen})^2 = m_{neutrino}(u_{neutrino})^2$$

Based on the above equation, the estimated mass of a neutrino is 6.27×10^{-38} kg or 0.0352 eV/ c^2 . The goal of this estimation is to predict an approximate mass of neutrinos.

The mass-speed relationship is derived, in the following three sections, from the kinetic theory of gases [6-10] and Avogadro's law [11] using a hypothesized setup of two boxes containing separated oxygen molecules and neutrino particles.

2. Separated Oxygen Molecules and Neutrinos

A hypothesized setup is designed to demonstrate the interactions between oxygen molecules and neutrino particles. Let each oxygen molecule have a mass m_1 and RMS speed u_1 , and each neutrino have a mass m_2 and RMS speed u_2 . Additionally, let the number of oxygen molecules be n_1 and the number of neutrinos be n_2 . Assume that the oxygen molecules and neutrino particles are initially separated in two adjacent boxes, as illustrated below.



Assume that the two boxes have the same pressure, volume, and temperature:

$$\begin{aligned} P_1 &= P_2 = P \\ V_1 &= V_2 = V \\ T_1 &= T_2 = T \end{aligned}$$

3. Kinetic Theory of Gases

Based on the kinetic theory of gases, the amounts, masses, and RMS speeds of the particles in Box 1 and Box 2 have the following relationship.

$$n_1 m_1 u_1^2 = n_2 m_2 u_2^2$$

The above relationship can be derived from the fundamental relationship between pressure, density, and RMS speed of particles:

$$P = \frac{1}{3} \rho u^2$$

Since density is defined by the ratio of mass m to volume V , the above equation can be modified into

$$P = \frac{1}{3} \frac{n m}{V} u^2$$

or, equivalently.

$$n m u^2 = 3PV$$


Because Box 1 and Box 2 have the same pressure and volume, the relationship between the amounts, masses, and RMS speeds of the particles in Box 1 and Box 2 can be formulated as

$$n_1 m_1 u_1^2 = n_2 m_2 u_2^2$$

4. Avogadro's Law

Avogadro's law states that "equal volumes of all gases, at the same temperature and pressure, have the same number of molecules". Because Box 1 and Box 2 have the same temperatures and pressures, it can be concluded from Avogadro's law that the number of oxygen molecules, n_1 , and neutrino particles, n_2 , are the same:

$$n_1 = n_2 = n$$



By removing the common factor n from the last equation in the last section, the relationship between the masses and RMS speeds of the particles in Box 1 and Box 2 can be formulated as


$$m_1 u_1^2 = m_2 u_2^2$$

or rearranged as

$$m_2 = m_1 \frac{u_1^2}{u_2^2}$$

5. Mixed Oxygen Molecules and Neutrinos

Because Box 1 and Box 2 have the same pressures, removing the divider between two boxes will not change RMS speeds (u_1, u_2) of the particles.



6. Neutrino Mass

Let m_1 and u_1 be the mass and RMS speed of oxygen molecules respectively; m_2 and u_2 be the mass and RMS speed of neutrinos respectively, the above equation can be represented as

$$m_{neutrino} = m_{oxygen} \left(\frac{u_{oxygen}}{u_{neutrino}} \right)^2$$

The RMS speed of neutrinos is close to the speed of light c

$$u_{neutrino} \cong 3.00 \times 10^8 \left[\frac{m}{s} \right]$$

The mass of an oxygen molecule is

$$m_{oxygen} \cong 5.31 \times 10^{-26} [kg]$$

The RMS speed of oxygen molecules at 20 °C is

$$u_{oxygen} \cong 326 \left[\frac{m}{s} \right]$$

Hence the mass of a neutrino is given by

$$\begin{aligned} m_{neutrino} &\cong 5.31 \times 10^{-26} [kg] \left(\frac{326 \left[\frac{m}{s} \right]}{3.00 \times 10^8 \left[\frac{m}{s} \right]} \right)^2 \cong 6.27 \times 10^{-38} [kg] \\ &\cong 6.27 \times 10^{-38} [kg] \left(\frac{3.00 \times 10^8 \left[\frac{m}{s} \right]}{c} \right)^2 \frac{1 [eV]}{1.60 \times 10^{-19} [J]} \cong 0.0352 \left[\frac{eV}{c^2} \right] \end{aligned}$$

This is the estimated mass of a neutrino based on the kinetic theory of gases.

7. Comparison to Measurements of Neutrino Mass

In June of 2008, KamLAND-Zen reported a lowest upper limit of neutrino mass of 0.06 – 0.161 eV/c², based on their measured data from their neutrino mass detector [12]. The estimated neutrino mass of 0.0352 eV/c² is about half of the KamLAND-Zen's lowest upper limit of neutrino mass of 0.06 eV/c².

8. Conclusions

By comparing neutrinos to molecules, the mass of a neutrino is estimated to be 6.27×10^{-38} kg or 0.0352 eV/c² according to Avogadro's law and the kinetic theory of gases. This mass agrees with the KamLAND-Zen's lowest upper limit of neutrino mass of 0.06 – 0.161 eV/c². This classical analysis using the kinetic theory of gases thus predicts an approximate mass of neutrinos in the range of what has previously been reported.

References

1. Fukuda, Y.; Hayakawa, T.; Ichihara, E.; Inoue, K.; Ishihara, K.; Ishino, H.; et al. (1998), "Measurements of the Solar Neutrino Flux from Super-Kamiokande's First 300 Days", *Physical Review Letters*, 81 (6): 1158–1162, arXiv:hep-ex/9805021, doi:10.1103/PhysRevLett.81.1158.
2. Araki, T.; Eguchi, K.; Enomoto, S.; Furuno, K.; Ichimura, K.; Ikeda, H.; et al. (2005), "Measurement of Neutrino Oscillation with KamLAND: Evidence of Spectral Distortion", *Physical Review Letters*, 94 (8): 081801, arXiv:hep-ex/0406035, doi:10.1103/PhysRevLett.94.081801.
3. "MINOS experiment sheds light on mystery of neutrino disappearance". <http://news.fnal.gov/2006/03/minos-experiment-sheds-light-mystery-neutrino-dis-appearance/> Fermilab. 30 March 2006.
4. Amsler, C.; Doser, M.; Antonelli, M.; Asner, D.M.; Babu, K.S.; Baer, H.; et al. (2008), "Review of Particle Physics". doi:10.1016/j.physletb.2008.07.018.
5. Agafonova, N.; Aleksandrov, A.; Altinok, O.; Ambrosio, M.; Anokhina, A.; Aoki, S.; et al. (2010), "Observation of a first ν_τ candidate event in the OPERA experiment in the CNGS beam", *Physics Letters B*, 691 (3): 138–145, arXiv:1006.1623, doi:10.1016/j.physletb.2010.06.022.
6. Waterston, J. J. (1892), "On the physics of media that are composed of free and perfectly elastic molecules in a state of motion", *Philosophical Transactions of the Royal Society of London, A*. Vol.183, pp.1-79.
7. Herapath, J. (1847), *Mathematical Physics*, London.
8. Maxwell, J.C. (1860), "V. Illustrations of the dynamical theory of gases, Part I. On the motions and collisions of perfectly elastic spheres", *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, 4th Series, vol.19. pp.19-32.
9. Maxwell, J.C. (1860), "II, Illustrations of the dynamical theory of gases, Part II. On the process of diffusion of two or more kinds of moving particles among one another", *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, 4th Series, vol.20. pp.21-37.
10. Maxwell, J.C. (1868), "XV, On the dynamical theory of gases", *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, 4th Series, vol.35. pp.129-145 and pp.185-217.
11. Avogadro, A. (1811), "Essai d'une maniere de determiner les masses relatives des molecules elementaires des corps, et les proportions selon lesquelles elles entrent dans ces combinations", *Journal de Physique*, vol.73, pp.58-76, English translation.
12. Gando, A.; et al. (KamLAND-Zen Collaboration) (11 May 2016), "Search for Majorana neutrinos near the inverted mass hierarchy region with KamLAND-Zen", *Physical Review Letters*, 117 (8): 082503, arXiv:1605.02889, doi:10.1103/PhysRevLett.117.082503.