

# The Alternative Schrödinger's Equation

**Michael Tzoumpas**

Mechanical and Electrical Engineer  
National Technical University of Athens  
Irinis 2, 15234 Athens, Greece

E-mail: m.tzoumpas@gmail.com

December 2018

**Abstract.** According to the unified theory<sup>1,2</sup> of dynamic space the inductive-inertial phenomenon<sup>3</sup> and its forces has been developed. These forces act on the electric units<sup>4</sup> of the dynamic space, forming the grouping<sup>3</sup> units (namely electric charges or forms of the electric field). So, by this inductive phenomenon and the phenomenon of motion<sup>5</sup> the wave function will be calculated. This wave function, which essentially interprets the phenomena of motion waves,<sup>6</sup> replaces the Schrödinger's equation.

*Keywords:* Inductive phenomenon; grouping units; wave function.

PACS numbers: 03.50.Kk, 12.10.-g

## 1. Magnetic forces are Coulomb's electric ones

The magnetic forces are described as electric ones created by grouping units<sup>3</sup> of the moving electrons (Fig. 1), due to the inductive-inertial phenomenon.<sup>3</sup> If  $Q$  is a moving electric charge at speed  $u$ , while  $Q_1$  is the respective electric charge of its grouping units, then it is obvious that

$$Q_1 = KQu, \quad (1)$$

where  $K$  is a ratio constant. We put in Eq. 1 the speed

$$u = u_a C_0, \quad (2)$$

where  $u_a$  is the respective timeless speed,<sup>7</sup> then


$$Q_1 = KQu_a C_0. \quad (3)$$

As  $u_a$  is dimensionless value then, due to Eq. 3, it should obviously apply

$$KC_0 = 1 \Rightarrow K = \frac{1}{C_0} \quad (4)$$

and so the Eq. 3 becomes

$$Q_1 = Qu_a \Rightarrow u_a = \frac{Q_1}{Q} \Rightarrow u_a^2 = \frac{Q_1^2}{Q^2}. \quad (5)$$



**Figure 1.** The grouping units and their first pair of reproduction extra grouping units

However, the timeless speed<sup>7</sup> has been found as a function of the pressure difference<sup>6</sup>  $\Delta P$  on both sides of the formation of the first grouping unit and of the cohesive pressure<sup>4</sup>  $P_0$ , namely it is

$$u_a = \sqrt{\frac{\Delta P}{P_0}} \Rightarrow u_a^2 = \frac{\Delta P}{P_0}. \quad (6)$$

Therefore, due to Eqs 5 and 6, it is

$$u_a^2 = \frac{Q_1^2}{Q^2} = \frac{\Delta P}{P_0} \Rightarrow \Delta P = P_0 u_a^2. \quad (7)$$

## 2. The wave function replaces Schrödinger's equation

The time and spatial fluctuation of the spherical formation of the first grouping unit implies a harmonic change in the difference<sup>6</sup>  $\Delta P$  of the cohesive pressure. Therefore, the first amplitude  $A_1$  (Fig. 2) of the pressure fluctuation  $\Delta P = P_0 u_a^2$  (Eq. 7) will be

$$A_1 = \frac{\Delta P}{2} = \frac{P_0}{2} u_a^2 \Rightarrow A_1 = \frac{P_0}{2} u_a^2 \quad (8)$$

and for  $u_a^2 = Q_1^2/Q^2$  (Eq. 5), then Eq. 8 becomes

$$A_1 = \frac{P_0}{2} \cdot \frac{Q_1^2}{Q^2}. \quad (9)$$

The electric charge of the second grouping unit, due to Eq. 5, becomes


$$Q_2 = Q_1 u_a. \quad (10)$$

The fluctuation amplitude  $A_2$  decreases, keeping in denominator the accelerated electric charge  $Q$  (Eq. 9) as the operative cause of the phenomenon, that is

$$A_2 = \frac{P_0}{2} \cdot \frac{Q_2^2}{Q^2}. \quad (11)$$

By replacing the electric charge  $Q_2 = Q_1 u_a$  (Eq. 10) of the second grouping unit in Eq. 11, the fluctuation amplitude  $A_2$  becomes

$$A_2 = \frac{P_0}{2} \cdot \frac{Q_2^2}{Q^2} = \frac{P_0}{2} \cdot \frac{Q_1^2 u_a^2}{Q^2} \Rightarrow A_2 = \frac{P_0}{2} \cdot \frac{Q_1^2}{Q^2} u_a^2. \quad (12)$$



**Figure 2.** Descending change of pressure difference  $\Delta P$  as motion arrow<sup>6</sup> of the electron with a motion formation diameter  $d = \lambda/2$ , where  $\lambda$  the wavelength of the harmonic fluctuation amplitude  $A$  of motion wave ( $A_1 = P_0 u_a^2/2$ ,  $A_2 = P_0 u_a^4/2$ ,  $A_3 = P_0 u_a^6/2$ , where  $u_a < 1$  the timeless speed<sup>7</sup> of the electron)

However, due to Eq. 9, the Eq. 12 becomes

$$A_2 = A_1 u_a^2, \quad (13)$$

which results in

$$A_2 = A_1 u_a^2, A_3 = A_2 u_a^2, A_4 = A_3 u_a^2, \dots, A_n = A_{n-1} u_a^2, \quad (14)$$

where  $A_n$  is the amplitude on either side of the formation and, due to Eq. 8, the Eq. 14 becomes


$$A_1 = \frac{P_0}{2} u_a^{2 \cdot 1}, A_2 = \frac{P_0}{2} u_a^{2 \cdot 2}, A_3 = \frac{P_0}{2} u_a^{2 \cdot 3}, \dots, A_n = \frac{P_0}{2} u_a^{2n}. \quad (15)$$

Therefore, we conclude that the fluctuation amplitude decreases with geometrical progress and more pronounced for low speeds, since the timeless speed<sup>7</sup> is  $u_a < 1$ . If  $u_a = 1$  (the timeless speed of light<sup>7</sup>), Eq. 15 becomes

$$A_n = \frac{P_0}{2} = \frac{\Delta P}{2} \Rightarrow \Delta P = P_0, \quad (16)$$

that is, the E/M wave<sup>8</sup> uses the entire cohesive pressure<sup>4</sup>  $P_0$  of constant amplitude, while its electric charge of grouping units are equivalent, since  $Q_1 = Q u_a$  (Eq. 5) or in general  $Q_n = Q_{n-1} u_a$  and for  $u_a = 1$  it is  $Q_n = Q_{n-1}$ .

The wavelength of the formation (Fig. 2) is  $\lambda = 2d$  and, of course, the first fluctuation amplitude of  $\Delta P$  is  $A_1 = P_0/2u_a^2$  (Eq. 8) observed at the ends of the half-wave  $\lambda/2$ . This fluctuation decreases by geometrical progress, as mentioned above.



**Figure 3.** Horizontal component  $F_x$  of kinetic force<sup>9</sup>  $F$  causes a sinusoidal change in the pressure difference  $\Delta P$  or the motion arrow<sup>7</sup> of the particle

The pressure difference  $\Delta P$  is a sinusoidal function of the distance  $x$  from the particle. Fig. 3 shows that the horizontal component  $F_x$  of the kinetic force<sup>9</sup>  $F$  causes a sinusoidal change in the pressure difference  $\Delta P$  or else in the motion arrow of the particle. Using the projection theorem (Fig. 3) we calculate the force  $F_x = F_s/\pi \cdot \sin\omega$  (Eq. 17) as follow:

The arc  $2\pi R/4 = \pi R/2$  corresponds to an accumulated force<sup>9</sup>  $F_s/4$  and the length  $x$  corresponds to a horizontal force  $F_x/2$ . So, for  $x/R = \sin\omega$ , it is

$$x = \frac{\pi R}{2} \cdot \frac{F_x/2}{F_s/4} = \pi R \frac{F_x}{F_s} \Rightarrow F_x = \frac{F_s}{\pi} \cdot \frac{x}{R} = \frac{F_s}{\pi} \sin\omega \quad (17)$$

as a sinusoidal function. The pressure difference  $\Delta P$ , caused by the  $F_x$  vertical to the cross-section  $\pi R^2$ , becomes

$$\Delta P = \frac{F_x}{\pi R^2} = \frac{F_s}{\pi^2 R^2} \cdot \frac{x}{R} = \frac{F_s}{\pi^2 R^2} \sin\omega \Rightarrow \Delta P = \frac{F_s}{\pi^2 R^2} \sin\omega \quad (18)$$

and so, it is a sinusoidal function. So, the pressure difference  $\Delta P$  can be written as

$$\Delta P = A \sin \frac{2\pi x}{\lambda}, \quad (19)$$

where  $A$  the fluctuation amplitude of wavelength  $\lambda = \lambda/2 + \lambda/2$  corresponding to the diameter  $d = \lambda/2$  of the grouping unit (Fig. 2). By replacing the general equation of amplitude  $A_n = P_0 u_a^{2n}/2$  (Eq. 15) in Eq. 19, the pressure difference  $\Delta P$  becomes

$$\Delta P = \frac{P_0}{2} u_a^{2n} \sin \frac{2\pi x}{\lambda}. \quad (20)$$

This wave function applies (Fig. 2) for

$$x = \frac{\lambda}{4}, \frac{3\lambda}{4}, \frac{5\lambda}{4}, \dots, \frac{(2n-1)\lambda}{4}. \quad (21)$$

From the general equation  $x = (2n-1)\lambda/4$  (Eq. 21) and for the absolute value of  $x$  it is found

$$n = \frac{2|x| + \lambda/2}{\lambda}. \quad (22)$$

Therefore, due to Eq. 22, the Eq. 20 becomes

$$\Delta P = \frac{P_0}{2} u_a^{\frac{4|x|+\lambda}{\lambda}} \sin \frac{2\pi x}{\lambda}, \quad (23)$$

which for  $|x| > \lambda/4$  decreases continuously.

This wave function, which essentially interprets the phenomena of motion waves, replaces Schrödinger's equation, where  $-\lambda/4 < x < +\lambda/4$ ,  $u_a = u/C_0$  the timeless speed<sup>7</sup> of the particle,  $u$  its time speed,  $C_0$  the light speed,  $\lambda = h/mu$  the so-called de Broglie's wave length,<sup>6</sup>  $h$  the Planck's constant,<sup>10</sup>  $m$  the mass<sup>6</sup> of the particle and  $P_0$  the cohesive pressure<sup>4</sup> of space.

### 3. References

- [1] N.I.Gosdas, *The Unified Theory of Dynamic Space*, Greek Edition (Trohalia, Athens, 1999).
- [2] M.Tzoumpas, *Hubble's Law and antigravity - Higgs boson and gravity*, <http://viXra.org/abs/1710.0082> [Quantum Gravity and String Theory].
- [3] M.Tzoumpas, *What is de Broglie's wave-particle?*, <http://viXra.org/abs/1806.0248> (section 1) [Quantum Gravity and String Theory].
- [4] M.Tzoumpas, *Hubble's Law and antigravity - Higgs boson and gravity*, <http://viXra.org/abs/1710.0082> (subsection 2.1) [Quantum Gravity and String Theory].
- [5] M.Tzoumpas, *Time as motion phenomenon - Physics Laws do not apply to inertial systems*, <http://viXra.org/abs/1802.0372> (section 4) [Relativity and Cosmology].
- [6] M.Tzoumpas, *What is de Broglie's wave-particle?*, <http://viXra.org/abs/1806.0248> (section 2) [Quantum Gravity and String Theory].
- [7] M.Tzoumpas, *The timeless Universe*, <http://viXra.org/abs/1804.0408> (section 1) [Astrophysics].
- [8] M. Tzoumpas, *The structure of autonomous motion - E/M waves*, <http://viXra.org/abs/1511.0025> (section 1) [Quantum Gravity and String Theory].
- [9] M.Tzoumpas, *Time as motion phenomenon - Physics Laws do not apply to inertial systems*, <http://viXra.org/abs/1802.0372> (section 5) [Relativity and Cosmology].
- [10] M.Tzoumpas, *Time as motion phenomenon - Physics Laws do not apply to inertial systems*, <http://viXra.org/abs/1802.0372> (section 3) [Relativity and Cosmology].