

On the compactification of the gravitational field

S. Halayka*

February 24, 2019

Abstract

The emission of gravitons by a mass is considered.

1 On graviton emission

It is assumed that the number of gravitons emitted from a kilogram of mass per second is some finite, countable number. In other words, it is assumed that the gravitational field is quantized, and that the quanta are gravitons. It is assumed that these gravitons propagate at the speed of light in vacuum c .

In *gravito-hydrodynamically-bound* systems like the Sun, a mass is an omnidirectional graviton emitter. However, if one is to increasingly gravitationally stimulate a mass, then the stimulation will eventually turn that mass from an omnidirectional graviton emitter into a unidirectional graviton emitter – a mass reciprocates gravitons toward the gravitational stimulation, *in lieu of* the mass's usual omnidirectional graviton emission. For a perfectly unidirectional graviton emitter, the strength of the gravitational interaction would increase by a factor of c^2 , because the gravitational field (a bunch of gravitons) would be compactified from a (3+1)D field down to a (1+1)D beam. A perfectly unidirectional graviton emitter would be a GASER (the gravitational analogue of the electromagnetic LASER).

In *gravitationally-bound* systems like the Milky Way, it is the inherently anisotropic gravitational interaction of the galaxy that makes the gravitational interaction stronger than that found in *gravito-hydrodynamically-bound* systems. For instance, the galaxy is a (3+1)D ball at the centre, and is more and more like a (2+1)D disk the further from the centre one goes. The strength of the gravitational interaction would increase as the shape goes from (3+1)D to (2+1)D, and in the case of a perfectly flat (2+1)D disk, this compactification of the gravitational field would make the gravitational interaction stronger by a factor of c . From this perspective, it is seen that dark matter is not absolutely required to explain the galactic dynamics. Likewise, the dynamics of cosmic filaments do not absolutely require dark matter if one idealizes a cosmic filament as a (1+1)D beam, where in perfect case the strength of the gravitational interaction would increase by a factor of c^2 .

*sjhalayka@gmail.com

References

- [1] Fontana G. Possible Graviton Transitions and Gaser Action in High-Tc Superconductors – <https://arxiv.org/abs/cond-mat/0208276>
- [2] Fontana G. Design of a Quantum Source of High-Frequency Gravitational Waves (HFGW) and Test Methodology – <https://arxiv.org/abs/physics/0410022>
- [3] Binney J. and Merrifield M. Galactic Astronomy – ISBN: 0-691-02565-7
- [4] Binney J. and Tremaine S. Galactic Dynamics, Second Edition – ISBN: 978-0-691-13027-9
- [5] Misner C.W., Thorne K.S., and Wheeler J.A. Gravitation – ISBN: 978-0716703440
- [6] Nicolson I. Dark Side of the Universe – ISBN: 978-0801885921