

THEORY OF ELECTRON

WU SHENG-PING

ABSTRACT. This article try to unified the four basic forces by Maxwell equations, the only experimental theory. Self-consistent Maxwell equation with the e-current from matter current is proposed, and is solved to four kinds of electrons and the structures of particles. The static properties and decay are reasoned, all meet experimental data. The equation of general relativity sheerly with electromagnetic field is discussed as the base of this theory. In the end the conformation elementarily between this theory and QED and weak theory is discussed.

CONTENTS

1. Unit Dimension of sch	1
2. Self-consistent Electrical-magnetic Fields	2
3. Calculation of Recursive Re-substitution	3
4. Solution	3
5. Electrons and Their Symmetries	4
6. Propagation and Movement	7
7. Balance Formula	9
8. Muon	9
9. Pion	9
10. Pion Neutral	10
11. Tauon	10
12. Proton	10
13. Neutron	11
14. Mesons	11
15. Mechanical Quantification	11
16. Great Unification	13
17. Conclusion	13
References	13

1. UNIT DIMENSION OF sch

A rebuilding of units and physical dimensions is needed. Time s is fundamental.

We can define:

The unit of time: s (second)

The unit of length: cs (c is the velocity of light)

Date: Mar 9, 2019.

Key words and phrases. Maxwell equations; Decay; Electron function; Recursive re-substitution.

The unit of energy: \hbar/s (\hbar is Plank constant)

The unit dielectric constant ϵ is

$$[\epsilon] = \frac{[Q]^2}{[E][L]} = \frac{[Q]^2}{\hbar c}$$

The unit of magnetic permeability μ is

$$[\mu] = \frac{[E][T]^2}{[Q]^2[L]} = \frac{\hbar}{c[Q]^2}$$

The unit of Q (charge) is defined as

$$c\epsilon = c\mu = 1$$

then

$$\begin{aligned} [Q] &= \sqrt{\hbar} \\ [H] &= [Q]/[L]^2 = [cD] = [E] \end{aligned}$$

Then

$$\sqrt{\hbar} : C = (1.0546 \times 10^{-34})^{1/2}$$

C is charge SI unit Coulomb.

For convenience, new base units by unit-free constants are defined,

$$c = 1, \hbar = 1, [Q] = \sqrt{\hbar}$$

then all physical units are power of second s^n , the units are reduced.

Define

$$\begin{aligned} \text{UnitiveElectricalCharge} : \sigma &= \sqrt{\hbar} \\ \sigma &= 1.03 \times 10^{-17} C \approx 64e \\ e_{/\sigma} &= e/\sigma = 1.57 \times 10^{-2} \approx 1/64 \end{aligned}$$

The system is redefined and rebuilt as:

$$s \rightarrow Cs : 1 = m_e/e_{/\sigma} =: \beta$$

$s \rightarrow Cs$ means the value of the redefined second becomes c seconds. We always use this units system.

2. SELF-CONSISTENT ELECTRICAL-MAGNETIC FIELDS

Try so-called expanded Maxwell equation for the free E-M field in the previously defined units

$$(2.1) \quad \begin{aligned} \partial' \cdot \partial A_\nu &= iA^{\mu*} \partial_\nu A_\mu / 2 + cc. = J_\nu \\ \partial_\nu \cdot A^\nu &= 0 \end{aligned}$$

with definition

$$\begin{aligned} (A^i) &:= (V, \mathbf{A}), (A_i) := (V, -\mathbf{A}) \\ (J^i) &= (\rho, \mathbf{J}), (J_i) = (\rho, -\mathbf{J}) \\ \partial &:= (\partial_i) := (\partial_t, \partial_{x_1}, \partial_{x_2}, \partial_{x_3}) \\ \partial' &:= (\partial^i) := (\partial_t, -\partial_{x_1}, -\partial_{x_2}, -\partial_{x_3}) \end{aligned}$$

It's deduced by using momentum to express e-current in a electron: the mass and charge have the same movement in electron. The equation 2.1 have symmetry

$$CPT, cc.PT$$

The following is the energy of a piece of field A :

$$(2.2) \quad \varepsilon := \frac{1}{2}(\langle E, E \rangle + \langle H, H \rangle)$$

$$(2.3) \quad \varepsilon|_{\infty}^t = \frac{1}{2}(\langle V|\partial \cdot \partial'|V \rangle - \langle \mathbf{A}|\partial \cdot \partial'|\mathbf{A} \rangle)|_{\infty}^t$$

The time-variant part is neglected, as a convention.

3. CALCULATION OF RECURSIVE RE-SUBSTITUTION

The solution by recursive re-substitution (RRS) for the two sides of the equation is proposed. For the equation

$$\hat{P}'B = \hat{P}B$$

Its algorithm is that (It's approximate, the exact solution needs a rate on the start state in the re-substitution for the normalization condition)

$$\hat{P}'\left(\sum_{k \leq n} B_k + B_{n+1}\right) = \hat{P}\sum_{k \leq n} B_k$$

A function is initially set and is corrected by re-substitution. Here is the initial state

$$V = V_i e^{-ikt}, A_i = V, \partial_{\mu} \partial^{\mu} A_i = 0$$

It's substituted into equation 2.1. The fields' correction A_n with n degrees of A_i is called the n degrees correction.

4. SOLUTION

Firstly

$$\nabla^2 A = -k^2 A$$

is solved. Exactly, it's solved in spherical coordinate

$$0 = r^2(\nabla^2 f + k^2 f) = k^2 r^2 f + (r^2 f_r)_r + \frac{1}{\sin \theta}(\sin \theta f_{\theta})_{\theta} + \frac{1}{\sin^2 \theta}(f_{\varphi})_{\varphi}$$

We use this term to replace the former. Its solution is

$$\begin{aligned} f &= R\Theta\Phi = R_l Y_{lm} \\ \Theta &= P_l^m(\cos \theta), \Phi = \cos(\alpha + m\varphi) \\ R_l &= N j_l(kr) \end{aligned}$$

$j_l(r)$ is spherical Bessel function.

$$\begin{aligned} j_1(r) &= \frac{\sin(r)}{r^2} - \frac{\cos r}{r} \\ j_1(0) &= 0 \end{aligned}$$

$$\int_0^{\infty} dr \cdot r^2 j_1(kr) j_1(k'r) = C k^{-2} \delta(k - k')$$

Define

$$\begin{aligned} F(x) &:= N R_1(r) Y_{1,1}(\theta, \varphi) \\ F^*(x) * F(x) &= \frac{\delta(r)}{4\pi r^2} \end{aligned}$$

with consideration

$$(4.1) \quad \nabla^2(F^*(x) * F(kx))$$

$F^*(x) * F(kx)$ have singularity at grid origin O .

$$\begin{aligned}
F^*(x) * F(x) &= \frac{1}{4\pi} \int_I \sin \theta' d\theta' d\varphi' dr' \cdot r' F^*(x') \cdot r' F(x - x') \\
&\quad \lim_{r' \rightarrow \infty} r' F^*(x') \cdot r' F(x - x') \\
&= C(\theta, \varphi, \theta', \varphi') \left(\frac{\sin(r')}{r'} - \cos(r') \right) \left((1 + C') \frac{\sin(r')}{r'} - \cos(r') \right) + O(r'^{-2}) \\
&\quad k^{3/2} F^*(kx) * k^{3/2} F(kx) \\
&= \frac{1}{4\pi} \int_I \sin \theta' d\theta' d\varphi' d(kr') \cdot kr' F^*(kx') \cdot kr' F(kx - kx') = \frac{k\delta(kr)}{4\pi k^2 r^2}
\end{aligned}$$

$\sin(r')/r'$ has fourier transform on variable r' .

Because the function F hasn't Fourier transform, and $\langle F, F \rangle$ is infinite, the limit of its forms truncated is then considered. It's defined as

$$\Omega_k(x) := NR_1(kx)Y, \langle \Omega_k(x) | \Omega_k(x) \rangle = 1$$

These set the following valid

$$\begin{aligned}
\langle \Omega(x) | \nabla | \Omega_k(x) \rangle &= \lim_{l \rightarrow \infty} \langle \Omega^l(x) | \nabla | \Omega_k^l(x) \rangle \\
\Omega_k^l(x) &:= \frac{\Omega_k(x)(h(x+l) - h(x-l))}{\langle \Omega_k(x)(h(x+l) - h(x-l)) | \Omega_k(x) \rangle^{1/2}}
\end{aligned}$$

and also through the limits

$$\nabla(\Omega(x) * \Omega_k(x)) = \lim_{l \rightarrow \infty} \nabla(\Omega^l(x) * \Omega_k^l(x)) = (\nabla\Omega(x)) * \Omega_k(x)$$

and so on. This is explained by a finite cosmos.

It's easy to prove

$$\langle \Omega_1^n(x), \Omega_1^n(x) \rangle = 1$$

We have the calculation

$$\begin{aligned}
\langle \nabla\Omega_k * \Omega_1 | \nabla\Omega_k * \Omega_1 \rangle &= \langle \nabla\Omega_k * \nabla\Omega_k^* | \Omega_1^* * \Omega_1 \rangle \\
&= \langle \nabla\Omega_k * \nabla\Omega_k^* | \langle \Omega_1 | \Omega_1 \rangle \rangle = - \langle (\nabla^2\Omega_k) * \Omega_k^* | \langle \Omega_1 | \Omega_1 \rangle \rangle
\end{aligned}$$

The solution of $l = 1, m = 1, Q = e/\sigma$ is calculated or tested for electron,

$$V = NR_1(-k_e r)Y_{1,-1}e^{ik_e t}$$

5. ELECTRONS AND THEIR SYMMETRIES


Some states of electrical field A are defined as the core of the electron, which's the start function $A_i = V$ that is electrical, for the RRS to get the whole electron function e :

$$\begin{aligned}
e_r^+ &: NR_1(k_e r)Y_{1,1}e^{-ik_e t}, \\
e_l^+ &: NR_1(k_e r)Y_{1,-1}e^{-ik_e t} \\
e_r^- &: NR_1(-k_e r)Y_{1,-1}e^{ik_e t} \\
e_l^- &: NR_1(-k_e r)Y_{1,1}e^{ik_e t}
\end{aligned}$$

These symbols is defined:

$$\begin{aligned}
(-e_l^-) &:= -e_r^+(-t), (-e_l^+) := -e_r^-(-t) \\
(-e_r^-) &:= -e_l^+(-t), (-e_r^+) := -e_l^-(-t)
\end{aligned}$$

r, l is the direction of the spin.

FIGURE 1. The function of j_1

The electron function with charge and mass is normalized, using the equation 2.1

$$\begin{aligned} \langle e_c^\mu | i\partial_t | e_{\mu c} \rangle &= Q_c \\ \langle \nabla e_\mu | \nabla e^\mu \rangle &\approx m_e \end{aligned}$$

Then

$$\begin{aligned} k_{ec} &\approx m_e / Q_c \\ \langle e_{c\mu} | e_c^\mu \rangle &\approx Q_c \end{aligned}$$

The magnetic dipole moment of electron is calculated as the first rank of proximation

$$\begin{aligned} \mathbf{r} \times i\partial \cdot \partial' A / 4 + cc. \\ \mu_z &= \langle A_i^\nu | i\partial_\phi | A_{i\nu} \rangle / 4 + cc. \\ &= \frac{Q_e}{2m_e} \end{aligned}$$

By the discussion in the section 2 the spin is

$$S_z = \mu_z k_e / e = 1/2$$

The correction in RRS of the equation 2.1 is calculated as

$$\begin{aligned} (5.1) \quad A_n &= A_{n-1} i\partial(A_i - A_i^*) / 2 * u \\ &= (A_i^* (i\partial_t A_i)) * u \cdot (i\partial_t(A_i - A_i^*) / 2 * u)^{n-3} i\partial(A_i - A_i^*) / 2 * u \\ &= (A_i^* (i\partial_t A_i)) (i\partial_t(A_i - A_i^*) / 2)^{n-3} (i\partial(A_i - A_i^*) / 2) |_{t \rightarrow t-r} \\ u &= \delta(t-r) / (4\pi r) \end{aligned}$$

with

$$\begin{aligned} (x', t') &:= (x, t-r), r'^2 = x' \cdot x' \\ \partial_x \cdot \partial_x - \partial_t^2 &= \partial_{x'} \cdot \partial_{x'} \\ f(x) e^{ict} * \frac{\delta(t-r)}{4\pi r} &= f(x) e^{ict} * \frac{1}{4\pi r} * \delta(t-r) \end{aligned}$$

and we have

$$\delta(t-a) *' \delta(t-b) \cdot \delta(t-c) = \delta(t-a) *' \delta(t-c) \cdot \delta(t-b)$$

with definition

$$f(x) *' g(x) := \int_{-\infty}^{\infty} f(x+t)g(t)dt$$

hence

$$(5.2) \quad f_0(t) *' f_1(t) \cdot f_2(t) = f_0(t) *' f_2(t) \cdot f_1(t)$$

$$f_i(t) := g_i(t) \sum_{n=-\infty}^{\infty} \delta(t-kn)$$

The first degree term of electron function is corrected with magnetic part

$$A_1^\nu = (\Omega_k(x)e^{-ikt}, i\nabla\Omega_k(x)e^{-ikt}/k)$$

then

$$(5.3) \quad \langle e^\nu | \partial \cdot \partial' | e_\nu \rangle = 0$$

$$\partial_\nu \cdot e^\nu = 0$$

$$\partial_\nu \cdot J_e^\nu = 0$$

These are Lorentz gauge and current constraint.

With the discussion of the section 7, it's valid that:

$$\langle (e_r^+)^\nu | \partial \cdot \partial' | (-e_l^+) \rangle = 0$$

$$\langle (e_r^+)^\nu | \partial \cdot \partial' | (-e_r^-) \rangle = 0$$

$$\langle (e_r^+)^\nu | \partial \cdot \partial' | (-e_l^-) \rangle = 0$$

The function of e_r^+ is decoupled with e_l^+

$$\langle (e_r^+)^\nu | \partial \cdot \partial' | (e_l^+) \rangle = 0$$

The following is the increment of field energy ε on the coupling of e_r^+ , e_r^- mainly between A_2

$$\varepsilon_e = \langle (e_r^+)^\nu | \partial \cdot \partial' | (e_r^-) \rangle \approx -e_{/\sigma}^3 m_e = -\frac{1}{1.66 \times 10^{-16} s}$$

The conditions 2.2, 2.3 is used. The following are the value of increments on the coupling of electrons

ε_e	e_r^+	e_r^-	e_l^+	e_l^-
e_r^+	+	-	0	0
e_r^-	-	+	0	0
e_l^+	0	0	+	-
e_l^-	0	0	-	+

The following is the increment of field energy ε on the coupling of e_r^+ , e_l^- mainly between A_4

$$\varepsilon_x = \langle (e_r^+)^\nu | \partial \cdot \partial' | (e_l^-) \rangle \approx -\frac{1}{2} e_{/\sigma}^7 m_e = -\frac{1}{1.09 \times 10^{-8} s}$$

The part other of these are counteracted, including the degree 8. The following are the value of increments on the coupling of electrons

$$\begin{array}{ccccc} \varepsilon_x & e_r^+ & e_r^- & e_l^+ & e_l^- \\ e_r^+ & + & 0 & 0 & - \\ e_r^- & 0 & + & - & 0 \\ e_l^+ & 0 & - & + & 0 \\ e_l^- & - & 0 & 0 & + \end{array}$$

6. PROPAGATION AND MOVEMENT

The propagation is:

$$f *_3 \sum_i e_i$$

The convolution is on space. The particle number normalization is:

$$\langle f, f \rangle = 1$$

The following are stable propagation:

<i>particle</i>	<i>electron</i>	<i>photon</i>	<i>neutino</i>
<i>notation</i>	e_r^\pm	γ_r	ν_r
<i>structure</i>	e_r^\pm	$(e_r^+ + e_r^-)$	$(e_r^+ + e_l^-)$

With the condition 5.3, their static mass is zero except the couplings.

The movement of the propagation is called *Movement*, ie. the third level wave:

$$f * \sum f_i * e_{ij}$$

The following coupling system of particle x is calculated

$$A_x = \sum_i e_i = \sum_c n_c e_c$$

c differ in charge, spin, and being negative or positive:

$$c = (r, l)(+, -)(p, n)$$

It's the initial and start state for RRS

$$A_i = e_x *_4 \sum_i e_i$$

$$e_x := \Omega_{k_x} f(t) \delta(t)$$

$$f(t) := e^{-ih(t)}, h \in \mathbf{R}$$

It meets the normalization of particle number. By referencing to the identity 5.2

$$f(t) \delta(t) * g(t) = f(t) g(-t) * \delta(-t)$$

f is chosen to obtain:

$$\partial \cdot \partial' A_i = 0$$

or equivalently

$$A_{i\nu}^* *' (\partial \cdot \partial' A_{i\nu}) = 0$$

then

$$\begin{aligned} f(t) &\approx e^{-ik_x t} \\ A_i &\approx \Omega_{k_x} e^{-ik_x t} *_3 \sum_i e_i \end{aligned}$$

The normalization of charge is

$$\langle A_i^\nu | i\partial_t A_{i\nu} \rangle = Q_x$$

then

$$k_x \approx n/Q_x, n := \sum_c n_c^2$$

The initial EM energy is

$$\varepsilon_i = \langle A_i^\nu | \partial \cdot \partial' | A_{i\nu} \rangle / 2 \approx n/e_\sigma$$

The initial MDM is

$$\mu_z \approx \langle A_i^\nu | -(\mathbf{x} + \mathbf{y}) \times i\nabla | A_{i\nu} \rangle / 4 + cc. \approx \frac{1}{2|k_x|e_\sigma} \sum_c Q_c n_c^2$$

The main part is generated from the crossing between the first degree term of the naked electron. And the current moment

$$|\mathbf{L}| \approx | \langle A_i^\nu | i\nabla | A_{i\nu} \rangle / 2 + cc. | \approx \left| \frac{m_e}{k_x} \sum_c Q_c n_c^2 \right|$$

The main current moment is generated from the crossing between the second degree correction and the first degree term of the naked electron.

The corrected field A meets the equation 2.1 with

$$A = A_i + \sum_{n=2}^{\infty} A_n$$

We notice that the terms in this equation are all delta functions of space, so that the interaction between the decayed blocks are released suddenly all at once, but this decaying process doesn't appear in the solution. Its solution is the one with electrons and their propagations coupled wholly in grid origin.

The following are the second degree correction of the equation 2.1:

$$\partial' \cdot \partial A_\nu = iA^{\mu*} \partial_\nu A_\mu / 2 + cc. = -I_s - I_c$$

For the light decay the self-effect at the initial time is

$$\begin{aligned} -I_{sc} &:= (e_x * n_c e_c)^{\nu*} \cdot i\partial(e_x * n_c e_c)_\nu / 2 + cc. \\ &= n_c^2 (e_x^* * e_x) (e_c^{\nu*} * i\partial e_{c\nu}) / 2 + cc. = n_c^2 e_x^* * e_x \cdot (\partial \cdot \partial') e_c \\ I_s &= \sum_c I_{sc} \end{aligned}$$

The following is the crossing effects at the initial time

$$\begin{aligned} -I_c &:= cc. + (n_{r+p} n_{r-p} - n_{r+n} n_{r-n}) (e_x * e_r^+)^{\nu*} \cdot i\partial(e_x * e_r^-)_\nu \\ &\quad + (n_{l+p} n_{l-p} - n_{l+n} n_{l-n}) (e_x * e_l^+)^{\nu*} \cdot i\partial(e_x * e_l^-)_\nu \\ &\quad + (n_{r+p} n_{l-p} - n_{r+n} n_{l-n}) (e_x * e_r^+)^{\nu*} \cdot i\partial(e_x * e_l^-)_\nu \\ &\quad + (n_{l+p} n_{r-p} - n_{l+n} n_{r-n}) (e_x * e_l^+)^{\nu*} \cdot i\partial(e_x * e_r^-)_\nu \end{aligned}$$

It's easy to use the coupling of these currents to calculate the emission of a decay mode. It's similar to calculate of strong effect.

7. BALANCE FORMULA

The reaction

$$A_{1i} - A_{2i} \rightarrow A_{1f} - A_{2f}$$

is equivalent to the following of the same energy emission

$$A_{1i} + A_{2f}(x, -t) \rightarrow A_{1f} + A_{2i}(x, -t)$$

by their forms as sums of odd function and even function of t . This means we can shift electron to the other side with the same emission of EM energy.

8. MUON

The core of muon is

$$\mu_r^- : e_\mu * (e_r^- - e_r^+ - e_l^-)$$

With the equation 15.1, μ is approximately with mass $3m_e/e_{j\sigma} = 3 \times 64m_e$ [3.2][1] (The data in bracket is experimental by the referenced lab), spin S_e (electron spin), MDM $\mu_B k_e/k_\mu$.

The main channel of decay is

$$\mu_r^- \rightarrow e_r^- - \nu_r, \quad e_r^- \rightarrow -e_r^- + \nu_l$$

with balance formula

$$\begin{aligned} & e_\mu * e_r^- + \delta^{1/2}(x + v_2 t) * \nu_r \\ & \rightarrow \delta^{1/2}(x - v_1 t) * e_r^- + e_\mu(-t) * \nu_r \end{aligned}$$

Referencing to the discussion of the section 6 to find the main EM emission as

$$\frac{\varepsilon_x m_e}{k_\mu} = -\frac{1}{2.18 \times 10^{-6} s} \quad [2.1970 \times 10^{-6} s][1]$$

9. PION

The core of pion is

$$\pi_l^+ : e_\pi(\varphi) * (e_r^+ \pm e_l^+) + e_\pi(-\varphi) * e_l^-$$

It's approximately with mass $3 \times 64m_e$ [4.2][1], spin S_e and MDM $\mu_B k_e/k_{\pi^+}$.

Decay Channels:

$$\pi_l^+ \rightarrow e_l^+ + \nu_r, \quad e_l^+ \rightarrow -e_l^+ + \nu_l$$

The mean life approximately is

$$-\varepsilon_x/2 = \frac{1}{2.2 \times 10^{-8} s} \quad [(2.603 \times 10^{-8} s)[1]$$

The precise result is calculated with successive decays.

10. PION NEUTRAL

The core of pion neutral is like a atom

$$\pi^0 : ((e_r^+ + e_l^+), (e_r^- + e_l^-))$$

It's with mass approximately $4 \times 64m_e$ [4.2][1], zero spin, and zero MDM. It's the main decay mode as

$$\pi^0 \rightarrow \gamma_r + \gamma_l$$

The loss of energy is

$$-2\varepsilon_e = \frac{1}{8.3 \times 10^{-17}s} \quad [8.4 \times 10^{-17}s][1]$$

The following particle is similar to π^0

$$(e_x(-\varphi) * (ne_r^+ - n'e_l^-) + e_x * (e_r^+ + e_l^+), \\ e_x(-\varphi) * (-ne_l^- + n'e_r^+) + e_x * (e_r^- + e_l^-))$$

11. TAUON

The core of tauon maybe

$$\tau_r^- : e_\tau(-\varphi) * (ne_r^+ - ne_l^-) + e_\tau * (e_r^- - e_l^- - e_l^+)$$

Its mass approximately $53 \times 64m_e$ [54][1] ($n = 5$), spin S_e and MDM $\mu_B m_e / k_\tau$. It has decay mode with a couple of neutrinos counteracted

$$e_r^- - e_l^- - e_l^+ \rightarrow e_r^- - \gamma_r$$

$$e_\tau * e_r^- + \delta^{1/2}(x + v_2 t) * \gamma_l \rightarrow \delta^{1/2}(x - v_1 t) * e_r^- + e_\tau(-t) * \gamma_l$$

The main EM emission is

$$\frac{\varepsilon_e m_e}{k_\tau} = -\frac{1}{5.5 \times 10^{-13}s} \quad [2.91 \times 10^{-13}s; BR. \quad 0.17][1]$$

Perhaps, it's a mixture with distinct coefficients n . The following particle is similar to τ

$$e_\tau(-\varphi) * (-ne_r^+ - e_l^+ - e_l^- + e_r^-) + e_\tau * (ne_l^- - e_r^+ - e_r^- - e_l^+ - e_l^-)$$

12. PROTON

The core of proton may be like

$$p_r^- : e_p * (-4e_r^+ - 3e_r^- - 2e_l^-)$$

The mass is $29 \times 64m_e$ [29][1] that's very close to the real mass. The MDM is calculated as $3\mu_N$, spin is S_e . The proton thus designed is eternal.

13. NEUTRON

Neutron is the atom of a proton and a electron and a neutrino,

$$n = (p_r^+, -\nu_r, e_l^-)$$

Neutrino circles around proton with

$$m_\nu \omega r^2 = 1$$

The interaction is between their (proton and neutrino) magnetic fields of A_2 (gross current)

$$m_\nu \omega^2 r = \frac{1}{4\pi} \cdot 3 \cdot 2 \cdot e_\sigma^2 k_e / (m_p r^2)$$

$$m_\nu = -e_{/\sigma}^7 m_e / 2$$

The EM energy emitted by neutrino is approximately

$$\frac{1}{2} \cdot \left(\frac{6}{4\pi}\right)^2 e_{/\sigma}^{7+2 \cdot 2} m_e^3 / (m_p^2 e_{/\sigma}^2) = \frac{1}{673s}$$

14. MESONS

We can define kinds of energy decreases of decay

interaction	EM	side EM	weak	side weak	strong
abbreviation	L	LS	W	WS	S
emission unit	$-\varepsilon_e$	$-\varepsilon_e m_e / k_x $	$-\varepsilon_x$	$-\varepsilon_x m_e / k_x $	m_e

We analyze the mesons [1] as the following

name	mass(MeV)	emission	type	ratio	construction
η	547	1.3keV	L	—	π^0 - class, perhaps
K^\pm	493	$1.24 \times 10^{-8}s$	W	0.88	$e_x(\varphi) * (ne_r^+ - e_l^- + e_l^+)$ $+e_x(-\varphi) * (-ne_l^- + e_r^-)$
K_S^0	—	$10^{-10}s$	—	—	unclear
K_L^0	—	$5 \times 10^{-8}s$	—	—	unclear
D^0	1865	$4100 \times 10^{-16}s$	LS	1.5	$((-ne_r^+ + n'e_r^- - e_l^+ - e_l^-)$
B^0	5279	$1.52 \times 10^{-12}s$	LS	1	$, (ne_l^- - n'e_l^+ - e_r^+ - e_r^-))$
D_S^\pm	1968	$5000 \times 10^{-16}s$	LS	1.3	τ - class
D^\pm	1869	$10400 \times 10^{-16}s$	LS	0.6	τ
B_S^\pm	5279	$1.6 \times 10^{-12}s$	LS	1	τ - class
Υ	9460	54keV	—	—	unclear
J/φ	3096	93keV	—	—	unclear
others	—	—	S	—	—

15. MECHANICAL QUANTIFICATION

It's the mechanic feature of the electromagnet fields as

$$T_{ij} = F_i^{k*} F_{kj} - g_{ij} F_{\mu\nu} F^{\mu\nu*} / 4$$

T is stress-energy tensor,

$$T_{ij} = \sum mu_i u_j, u = dx/ds$$

T_{00} is quantum expression of the energy, by Lorentz transform it's easy to get the quantum expression of momentum:

$$A^{\nu*} i \partial A_\nu + cc. \propto \sum mu$$

The observed mass in mass-center frame is

$$M = \int dV T_{00}$$

$$(15.1) \quad M = \varepsilon$$

It's obtained by moving the frame and then the energy is checked.

The decayed part is independent (uncoupled) of the undecayed part for the equation 2.1, hence the undecayed part A' meets the same condition

$$\partial' \cdot \partial A'_\nu = iA'^{\mu*} \partial_\nu A'_\mu / 2 + cc.$$

In principle, we can solve this equation to find the decay process. For the partial field A_n of n independent particles decaying to stable state,

$$F = \langle A'_n | \partial \cdot \partial' | A_{n\nu} \rangle = 2n\Gamma e^{-C\Gamma t} = \langle A_n^\mu | iA_n^{\nu*} \partial_\mu A_{n\nu} / 2 + cc. \rangle$$

The conditions 2.2, 2.3 are used. For some number n and charge Q

$$(15.2) \quad 1 = \int_0^\infty dt \langle A'_1 | \partial \cdot \partial' | A_{1\nu} \rangle = \frac{Q}{\sigma} \left(\int_0^\infty dt \langle A_1^\mu | iA_1^{\nu*} \partial_\mu A_{1\nu} / 2 + cc. \rangle \right), Q = 1$$

$$1 = \int_0^\infty dt \langle A'_n | \partial \cdot \partial' | A_{n\nu} \rangle = \int_0^\infty dt \langle A_n^\mu | iA_n^{\nu*} \partial_\mu A_{n\nu} / 2 + cc. \rangle, \sigma = 1$$

then

$$1 = \left(\int_0^\infty dt \langle A'_n | \partial \cdot \partial' | A_{n\nu} \rangle \right)^3 = \left(\langle A_n^\mu | iA_n^{\nu*} \partial_\mu A_{n\nu} / 2 + cc. \rangle \right)^2$$

$$1 = n^3 \left(\int_0^\infty dt \langle A'_1 | \partial \cdot \partial' | A_{1\nu} \rangle \right)^3 = n^2 \left(\int_0^\infty dt \langle A_1^\mu | iA_1^{\nu*} \partial_\mu A_{1\nu} / 2 + cc. \rangle \right)^2$$

$$\left(\int_0^\infty dt \langle A'_1 | \partial \cdot \partial' | A_{1\nu} \rangle \right)^3 = \left(\int_0^\infty dt \langle A_1^\mu | iA_1^{\nu*} \partial_\mu A_{1\nu} / 2 + cc. \rangle \right)^2$$

$$\frac{Q}{\sigma} = n^{-1/2}$$

They are substituted into the equation 15.2 to lead to

$$n = 1$$

hence

$$(15.3) \quad 1 = \int_0^\infty dt \langle A^\nu | \partial \cdot \partial' | A_\nu \rangle / 2$$

It's the normalization of one decaying particle, and it leads to the result between decay life and EM emission (also the interaction potential):

$$C = 1$$

The distribution shape of decay can be explain as

$$e^{-\Gamma t/2} e_x * \sum_i e_i \approx \Omega_x * \sum_i e_i \cdot e^{-\Gamma t/2 - ik_x t}, 0 < t < \Delta$$

It's the real wave of the particle x near the initial time and expanded in that time span

$$\approx \Omega_x * \sum_i e_i \cdot \int_{-\infty}^\infty dk \frac{C e^{-ikt}}{k - k_x - i\Gamma/2}$$

16. GREAT UNIFICATION

The General Theory of Relativity is

$$(16.1) \quad R_{ij} - \frac{1}{2}Rg_{ij} = 8\pi GT_{ij}/c^4$$

Firstly the unit second is redefined as S to simplify the equation 16.1

$$R_{ij} - \frac{1}{2}Rg_{ij} = T_{ij}$$

Then

$$R_{ij} - \frac{1}{2}Rg_{ij} = F_i^{k*}F_{kj} - g_{ij}F_{\mu\nu}F^{\mu\nu*}/4$$

We observe that the co-variant curvature is

$$R_{ij} = F_{ik}^*F_j^k + g_{ij}F_{\mu\nu}^*F^{\mu\nu}/8$$

17. CONCLUSION

Fortunately, this model explained all the effects in the known world: strong, weak and electromagnetic effects, and even subclassify them further if not being to add new ones. In this model the only field is electromagnetic field, and this stands for the philosophical that the unified world is from an unique source, all that depend on a simple hypothesis: the current of matter in a system can be devised to analysis the e-charge current.

My description of particles is compatible with QED elementarily, and only contributes to it with theory of consonance state in fact. In some way, the electron function is a good promotion for the experimental models of proton and electron that went up very early.

Underlining my calculations a fact is that the electron has the same phase (electron resonance), which the BIG BANG theory would explain, all electrons are generated in the same time and place, the same source.

REFERENCES

- [1] K. Nakamura et al. (Particle Data Group), JPG 37, 075021 (2010) (URL: <http://pdg.lbl.gov>)
E-mail address: sunylock@139.com

8 HANBEI RD, TIANMEN, HUBEI PROVINCE, THE PEOPLE'S REPUBLIC OF CHINA. POSTCODE: 431700