

# Expanding Universe from Weyl Cosmology and Asymptotic Safety in Quantum Gravity \*

Carlos Castro Perelman  
Center for Theoretical Studies of Physical Systems  
Clark Atlanta University, Atlanta, GA. 30314  
Ronin Institute, 127 Haddon Pl., NJ. 07043  
perelmanc@hotmail.com

June 2019

## Abstract

A study of the simplest Jordan-Brans-Dicke-like action within the context of Weyl geometry, combined with the findings of Weinberg's Asymptotic Safety program in quantum gravity, leads to a plethora of nice numerical results : (i) like singling out the quartic potential from all the others; (ii) having (Anti) de Sitter space as the most natural solution; (iii) furnishing the value of the observed vacuum energy density at the Hubble scale  $\frac{3}{8\pi G_N R_H^2} \sim 10^{-122} M_P^4$ ; (iv) and a value of  $\frac{3}{8\pi} M_P^4$  for the vacuum energy density at the Planck scale; (v) interpreting the "Bang" of the Big Bang as the *singularity* of the Weyl gauge field of dilations at  $t = 0$  ushering in the era of inflation, and (vi) allowing the possibility that our universe is a Black Hole whose horizon coincides with the cosmological Hubble horizon. It is warranted to explore deeper the interplay among Weyl geometry, Asymptotic safety and Maldacena's *AdS/CFT* correspondence (holographic renormalization group flow). Also relevant is the work by Wetterich on the role of dilatation symmetry in higher dimensions and the vanishing of the cosmological constant. Last, but not least, we should also consider the implications of Penrose' Conformal Cyclic Cosmology and Nottale's Scale Relativity Theory with the key findings of this work.

Keywords: Weyl Geometry; Cosmology; Asymptotic Safety; Brans-Dicke-Jordan Gravity; Cosmological Constant.

---

\*Dedicated to the loving memory of Irina Novikova, a brilliant and heavenly creature who met a tragic death at a young age.

# 1 A Brief Introduction : Why Weyl Geometry

The problem of dark energy and the solution to the cosmological constant problem is one of the most challenging problems facing Cosmology today. There are a vast numerable proposals for its solution. We refer to the monograph [13], [12], [6], [30] and many references therein for details. There have been many previous proposals [11] to explain dark matter (instead of dark energy) in terms of Brans-Dicke gravity [23]. One purpose of this work is to show that it is not only necessary to include the Jordan-Brans-Dicke scalar field  $\phi$ , but it is essential to include a Weyl geometric extension and generalization of Riemannian geometry (ordinary gravity ). In doing so we shall see how one can recover a plethora of key numerical results in Cosmology. The literature on Weyl geometry [1] is quite large, see [2], [3], [4], [11], [8], [24], [29], among many others, for references.

Given the Lorentzian signature  $(-, +, +, +)$ , let us begin with an action in a curved *Riemannian* background

$$S = \int d^4x \sqrt{|g|} \left( \frac{R}{16\pi G_o} - \frac{g^{\mu\nu}}{2} (\partial_\mu \Phi) (\partial_\nu \Phi) - V(\Phi) \right) \quad (1.1)$$

and associated with a canonical real scalar field  $\Phi$  with a known prescribed potential  $V(\Phi)$ . Varying the action with respect to the two fields  $g_{\mu\nu}, \Phi$  yields

$$R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R = 8\pi G_o \left( (\partial_\mu \Phi)(\partial_\nu \Phi) - \frac{1}{2} g_{\mu\nu} g^{\alpha\beta} (\partial_\alpha \Phi)(\partial_\beta \Phi) - g_{\mu\nu} V(\Phi) \right) \quad (1.2)$$

$$\frac{1}{\sqrt{|g|}} \partial_\mu \left( \sqrt{|g|} g^{\mu\nu} \partial_\nu \Phi \right) - \frac{\partial V(\Phi)}{\partial \Phi} = 0 \quad (1.3)$$

The two equations (1.2, 1.3) (Einstein-Klein-Gordon system) are now *coupled* and induce a *nonlinear* Klein-Gordon-like equation for  $\Phi$  after solving eq-(1.2) for the metric  $g_{\mu\nu}$  in terms of  $\Phi$ . Namely, a substitution of the form  $g_{\mu\nu}[\Phi]$  into (1.3) yields a *nonlinear* Klein-Gordon-like equation.

The authors [27] have shown that the nonrelativistic limit of the two coupled equations (1.2,1.3) furnish the *nonlinear* Newton-Schrödinger equation

$$i\hbar \frac{\partial \Psi(\vec{r}, t)}{\partial t} = - \frac{\hbar^2}{2m} \nabla^2 \Psi(\vec{r}, t) + V \Psi(\vec{r}, t) - \left( Gm^2 \int \frac{|\Psi(\vec{r}', t)|^2}{|\vec{r} - \vec{r}'|} d^3r' \right) \Psi(\vec{r}, t) \quad (1.4)$$

which is obtained after solving the Poisson equation

$$\nabla^2 U = 4\pi G_o m \rho = 4\pi G_o m \Psi^* \Psi \quad (1.5)$$

for the Newtonian potential  $U = U(\Psi, \Psi^*)$  and substituting its value into the Schrödinger equation.

The immediate advantage of recurring to Weyl geometry, is that it will allow us to find exact solutions to the very complicated coupled system of equations (1.2, 1.3). And, in doing so, it will furnish the value of the observed vacuum energy density at the Hubble scale  $\frac{3}{8\pi G_N R_H^2} \sim 10^{-122} M_P^4$ ; and a value  $\frac{3}{8\pi} M_P^4$  for the vacuum energy density at the Planck scale.

It is essential to emphasize that the Weyl geometric approach undertaken in this work is *not* the same as working with Conformal gravity based on the full Conformal group of translations, Lorentz boosts, dilations, and conformal boosts. The standard approach in Conformal gravity to recover ordinary Einstein gravity is based on the action corresponding to the kinetic terms of a real scalar field  $\varphi$

$$S = \int d^4x \sqrt{|g|} \frac{1}{2} [ \varphi D_\mu^c D_c^\mu \varphi ] \quad (1.6)$$

the conformal Laplacian can be rewritten

$$D_\mu^c D_c^\mu \varphi = \frac{1}{\sqrt{|g|}} \partial_\mu \left( \sqrt{|g|} g^{\mu\nu} D_\nu^c \varphi \right) + A_\mu D_c^\mu \varphi + \frac{\varphi}{6} R \quad (1.7)$$

One can fix, firstly, the conformal boosts transformations by choosing the gauge  $A_\mu = 0$ . And, secondly, one can then fix the scaling symmetry by gauging  $\varphi$  to a constant  $\varphi_o^2 = (16\pi G_N)^{-1}$ . In this way one recovers the Einstein-Hilbert action from the last term of eq.(1.7).

Weyl's geometry main feature is that the norm of vectors under parallel infinitesimal displacement going from  $x^\mu$  to  $x^\mu + dx^\mu$  change as follows  $\delta||V|| \sim ||V|| A_\mu dx^\mu$  where  $A_\mu$  is the Weyl gauge field of scale calibrations that behaves as a connection under Weyl transformations :

$$A'_\mu = A_\mu - \partial_\mu \Omega(x). \quad g_{\mu\nu} \rightarrow e^{2\Omega} g_{\mu\nu}. \quad (1.8)$$

involving the Weyl scaling parameter  $\Omega(x^\mu)$ . The Weyl covariant derivative operator acting on a tensor  $\mathbf{T}$  is defined by  $D_\mu \mathbf{T} = ( \nabla_\mu + \omega(\mathbf{T}) A_\mu ) \mathbf{T}$ ; where  $\omega(\mathbf{T})$  is the Weyl weight of the tensor  $\mathbf{T}$  and the derivative operator  $\nabla_\mu = \partial_\mu + \Gamma_\mu$  involves a connection  $\Gamma_\mu$  which is comprised of the ordinary Christoffel symbols  $\{\rho_{\mu\nu}\}$  plus the  $A_\mu$  terms

$$\Gamma_{\mu\nu}^\rho = \{\rho_{\mu\nu}\} + \delta_\mu^\rho A_\nu + \delta_\nu^\rho A_\mu - g_{\mu\nu} g^{\rho\sigma} A_\sigma \quad (1.9)$$

The Weyl gauge covariant operator  $\partial_\mu + \Gamma_\mu + w(\mathbf{T})A_\mu$  obeys the condition

$$D_\mu (g_{\nu\rho}) = \nabla_\mu (g_{\nu\rho}) + 2 A_\mu g_{\nu\rho} = 0. \quad (1.10)$$

where  $\nabla_\mu (g_{\nu\rho}) = -2 A_\mu g_{\nu\rho} = Q_{\mu\nu\rho}$  is the non-metricity tensor. Torsion can be added [3] if one wishes but for the time being we refrain from doing so. The connection  $\Gamma_{\mu\nu}^\rho$  is Weyl *invariant* so that the geodesic equation in Weyl spacetimes is Weyl-covariant under Weyl gauge transformations (scalings)

$$ds \rightarrow e^\Omega ds; \frac{dx^\mu}{ds} \rightarrow e^{-\Omega} \frac{dx^\mu}{ds}; \frac{d^2x^\mu}{ds^2} \rightarrow e^{-2\Omega} \left[ \frac{d^2x^\mu}{ds^2} - \frac{dx^\mu}{ds} \frac{dx^\nu}{ds} \partial_\nu \Omega \right].$$

$$g_{\mu\nu} \rightarrow e^{2\Omega} g_{\mu\nu}; A_\mu \rightarrow A_\mu - \partial_\mu \Omega; A^\mu \rightarrow e^{-2\Omega} (A^\mu - \partial^\mu \Omega); \Gamma_{\mu\nu}^\rho \rightarrow \Gamma_{\mu\nu}^\rho. \quad (1.11)$$

The Weyl connection and curvatures scale as

$$\Gamma_{\mu\nu}^\rho \rightarrow \Gamma_{\mu\nu}^\rho, \quad \mathcal{R}_{\mu\nu\sigma}^\rho \rightarrow \mathcal{R}_{\mu\nu\sigma}^\rho, \quad \mathcal{R}_{\mu\nu} \rightarrow \mathcal{R}_{\mu\nu}, \quad \mathcal{R} \rightarrow e^{-2\Omega} \mathcal{R} \quad (1.12)$$

Thus, the Weyl covariant geodesic equation transforms under Weyl scalings as

$$\frac{d^2x^\rho}{ds^2} + \Gamma_{\mu\nu}^\rho \frac{dx^\mu}{ds} \frac{dx^\nu}{ds} - A_\mu \frac{dx^\mu}{ds} \frac{dx^\rho}{ds} = 0 \rightarrow$$

$$e^{-2\Omega} \left[ \frac{d^2x^\rho}{ds^2} + \Gamma_{\mu\nu}^\rho \frac{dx^\mu}{ds} \frac{dx^\nu}{ds} - A_\mu \frac{dx^\mu}{ds} \frac{dx^\rho}{ds} \right] = 0. \quad (1.13)$$

The Weyl weight of the metric  $g_{\nu\rho}$  is 2. The meaning of  $D_\mu(g_{\nu\rho}) = 0$  is that the angle formed by two vectors remains the same under parallel transport despite that their lengths may change. This also occurs in conformal mappings of the complex plane. The Weyl covariant derivative acting on a scalar  $\phi$  of Weyl weight  $\omega(\phi) = -1$  is defined by

$$D_\mu \phi = \partial_\mu \phi + \omega(\phi) A_\mu \phi = \partial_\mu \phi - A_\mu \phi. \quad (1.14)$$

The Weyl scalar curvature in  $D$  dimensions and signature  $(-, +, +, +, \dots)$  is <sup>1</sup>

$$\mathcal{R}_{Weyl} = R_{Riemann} - (d-1)(d-2)A_\mu A^\mu - 2(d-1)\nabla_\mu A^\mu. \quad (1.15)$$

Having reviewed very briefly the basics of Weyl geometry we shall embark with the crux of this work.

## 2 Weyl Cosmology and Asymptotic Safety

### 2.1 A Weyl invariant Jordan-Brans-Dicke Model

Having introduced the basics of Weyl geometry our starting action is the Weyl-invariant Jordan-Brans-Dicke-like action involving the scalar  $\phi$  field and the scalar Weyl curvature  $\mathcal{R}_{Weyl}$

$$S[g_{\mu\nu}, A_\mu, \phi] = S[g'_{\mu\nu}, A'_\mu, \phi'] \Rightarrow$$

<sup>1</sup>Some authors define their  $A_\mu$  field with the opposite sign as  $-A_\mu$  which changes the sign in the last term of the Weyl scalar curvature (1.15)

$$\int d^4x \sqrt{|g|} [ \phi^2 \mathcal{R}_{Weyl}(g_{\mu\nu}, A_\mu) - \frac{1}{2} g^{\mu\nu} (D_\mu \phi)(D_\nu \phi) - V(\phi) ] =$$

$$\int d^4x \sqrt{|g'|} [ (\phi')^2 \mathcal{R}_{Weyl}(g'_{\mu\nu}, A'_\mu) - \frac{1}{2} g'^{\mu\nu} (D'_\mu \phi')(D'_\nu \phi') - V(\phi') ] \quad (2.1)$$

where under Weyl scalings one has

$$\phi' = e^{-\Omega} \phi; \quad g'_{\mu\nu} = e^{2\Omega} g_{\mu\nu}; \quad \mathcal{R}_{Weyl}(g'_{\mu\nu}, A'_\mu) = e^{-2\Omega} \mathcal{R}_{Weyl}(g_{\mu\nu}, A_\mu)$$

$$V(\phi') = e^{-4\Omega} V(\phi), \quad \sqrt{|g'|} = e^{4\Omega} \sqrt{|g|}; \quad D'_\mu \phi' = e^{-\Omega} D_\mu \phi; \quad A'_\mu = A_\mu - \partial_\mu \Omega. \quad (2.2)$$

One could complicate matters by adding more terms to the most simple action (2.1) like

$$\int d^4x \sqrt{|g|} ( \mathcal{R}_{Weyl}^2 + F_{\mu\nu} F^{\mu\nu} + \dots ) \quad (2.3)$$

see [7], [2], [8]. However, it won't be necessary to modify the action (2.1) to recover fundamental results.

The effective Newtonian coupling  $G$  is related to  $\phi$  as follows  $(16\pi G)^{-1} \equiv \phi^2$ , and it is spacetime dependent in general and has a Weyl weight equal to 2. Despite that one has *not* introduced any explicit dynamics to the  $A_\mu$  field (there are no  $F_{\mu\nu} F^{\mu\nu}$  terms in the action (2.1)) one still has the equation obtained from the variation of the action in  $d = 4$  w.r.t to the  $A^\mu$  field and which leads to the pure-gauge configurations provided  $\phi \neq 0$

$$\frac{\delta S}{\delta A_\mu} = 0 \Rightarrow \phi^2 \frac{\delta \mathcal{R}_{Weyl}}{\delta A_\mu} + \frac{\delta(D_\mu \phi)}{\delta A_\mu} \frac{\delta S_{matter}}{\delta(D_\mu \phi)} = 0 \Rightarrow$$

$$g^{\mu\nu} D_\nu \phi^2 = 0 \Rightarrow D_\mu \phi = 0 \Rightarrow A_\mu = \partial_\mu \ln(\phi). \quad (2.4)$$

Hence, a variation of the action w.r.t the  $A_\mu$  field leads to the pure gauge solutions (2.4) which is tantamount to saying that the scalar  $\phi$  is Weyl-covariantly *constant*  $D_\mu \phi = 0$  in any gauge  $D_\mu \phi = 0 \rightarrow e^{-\Omega} D_\mu \phi = D'_\mu \phi' = 0$  (for non-singular gauge functions  $\Omega \neq \pm\infty$ ).

Therefore, the scalar  $\phi$  does not have true local dynamical degrees of freedom from the Weyl spacetime perspective. Since the gauge field is a total derivative, under a local gauge transformation with a gauge function  $\Omega = \ln(\phi/\phi_o)$ , one can gauge away (locally) the gauge field  $A_\mu$  and have  $A'_\mu = 0$  in the new gauge. Globally, however, this may not be the case because there may be *topological* obstructions. Therefore, the gauge  $A'_\mu = 0$ , implies that  $\phi' = \phi_o = \text{constant}$ . Consequently  $16\pi G' = \phi'^{-2}$  can be fixed to a constant, and one may set  $G' = G_N$  where  $G_N$  is the observed Newtonian gravitational coupling.

The pure-gauge configurations leads to the Weyl integrability condition  $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu = 0$  when  $A_\mu = \partial_\mu \Omega$ , and means physically that if we parallel transport a vector under a closed loop in a *flat* spacetime, as we come back to the starting point, the *norm* of the vector has not changed; i.e, the rate at

which a clock ticks does not change after being transported along a closed loop back to the initial point; and if we transport a clock from  $A$  to  $B$  along different paths, the clocks will tick at the same rate upon arrival at the same point  $B$ . This will ensure, for example, that the observed spectral lines of identical atoms will not change when the atoms arrive at the laboratory after taking different paths ( histories ) from their coincident starting point. In this way on can bypass Einstein's objections to Weyl. If  $F_{\mu\nu} \neq 0$  the Weyl geometry is no longer integrable. This can occur if one adds explicit  $F_{\mu\nu}F^{\mu\nu}$  terms to the action which may lead to true dynamical degrees of freedom for the gauge field  $A_\mu$ .

This result  $D_\mu\phi = 0$  also follows in other dimensions. Substituting

$$A_\mu = \frac{2}{d-2} \partial_\mu \ln\phi \quad (2.5)$$

into

$$\mathcal{R}_{Weyl} = R_{Riemann} - (d-1)(d-2)A_\mu A^\mu - 2(d-1)\nabla_\mu A^\mu \quad (2.6)$$

gives

$$\mathcal{R}_{Weyl} = R_{Riemann} - 4 \frac{d-1}{d-2} \frac{\nabla_\mu \nabla^\mu \phi}{\phi} \quad (2.7)$$

The last term in (2.7) has a similar functional form as Bohm's quantum potential [21], [22]. From now we shall denote  $R$  for the Riemannian scalar curvature  $R_{Riemann}$ . The covariant derivative  $\nabla_\mu$  appearing in (2.7) is the one defined in terms of the Christoffel connection  $\{\}$ , and *not* based on the Weyl connection  $\Gamma$ .

Given an action (2.1) in  $d = 4$  the field equations are obtained after the variations of the action with respect to the 3 fields  $g_{\mu\nu}, A_\mu, \phi$ , respectively

$$\begin{aligned} \phi^2 \left( \mathcal{R}_{\mu\nu}^{Weyl} - \frac{1}{2} g_{\mu\nu} \mathcal{R}_{Weyl} \right) + D_\mu D_\nu \phi^2 + g_{\mu\nu} g^{\alpha\beta} D_\alpha D_\beta \phi^2 = \\ \frac{1}{2} (D_\mu \phi)(D_\nu \phi) - \frac{1}{4} g_{\mu\nu} g^{\alpha\beta} (D_\alpha \phi)(D_\beta \phi) - \frac{1}{2} g_{\mu\nu} V(\phi) \end{aligned} \quad (2.8)$$

$$D_\mu \phi^2 = 2 D_\mu \phi = 0 \Rightarrow A_\mu = \partial_\mu \ln(\phi) \quad (2.9)$$

$$2\phi \mathcal{R}_{Weyl} - \frac{\partial V(\phi)}{\partial \phi} + D_\mu D^\mu \phi = 0 \quad (2.10)$$

As stated earlier, the field equation  $D_\mu\phi = 0$  just states the  $\phi$  is Weyl-covariantly constant. This result when followed by taking the trace of (2.8) gives  $\phi^2 \mathcal{R}_{Weyl} = 2V(\phi)$  which allows to eliminate  $\mathcal{R}_{Weyl} = 2\phi^{-2}V(\phi)$ , and inserting it in eq (2.10) yields  $4\phi^{-1}V(\phi) - V'(\phi) = 0$ , *singling* out the quartic potential  $V(\phi) = \kappa\phi^4$  in  $d = 4$ , out of an infinity of possible choices for the potential. For example, one could have potentials of the form  $V = \sum_n c_n M^{4-n}\phi^n$  where  $M$  is mass-like

parameter (a scalar moduli parameter) which scales as  $M \rightarrow e^{-\Omega} M$  in order to render the action Weyl invariant. To sum up, in this Weyl geometric approach the choice for the potential  $V = \kappa\phi^4$  is *not* ad hoc but can be inferred from the field equations themselves.

Eq-(2.10) in  $d = 4$  can be rewritten in terms of the Riemannian scalar curvature, after using  $D_\mu\phi = 0$ , as

$$2 R \phi - \frac{12}{\sqrt{|g|}} \partial_\mu \left( \sqrt{|g|} g^{\mu\nu} \partial_\nu \phi \right) - \frac{\partial V(\phi)}{\partial \phi} = 0 \quad (2.11)$$

Upon inserting the derived expression for  $V(\phi) = \kappa\phi^4$  above, it gives

$$R \phi - \frac{6}{\sqrt{|g|}} \partial_\mu \left( \sqrt{|g|} g^{\mu\nu} \partial_\nu \phi \right) - 2 \kappa \phi^3 = 0 \quad (2.12)$$

It remains now to solve eq-(2.8) given  $D_\mu\phi = 0$  and  $V(\phi) = \kappa\phi^4$ . After factoring out  $\phi^2$  and substituting  $\mathcal{R}_{Weyl} = 2\kappa\phi^2$  leads to

$$\mathcal{R}_{\mu\nu}^{Weyl} = \frac{1}{2} g_{\mu\nu} \kappa\phi^2 \quad (2.13)$$

with

$$\mathcal{R}_{\mu\nu}^{Weyl} = R_{\mu\nu} - 2 \nabla_\mu A_\nu - g_{\mu\nu} g^{\alpha\beta} \nabla_\alpha A_\beta - 2 A_\mu A_\nu + 2 g_{\mu\nu} g^{\alpha\beta} A_\alpha A_\beta \quad (2.14)$$

Since the Weyl weight of  $\mathcal{R}_{\mu\nu}^{Weyl}$  is 0, after some straightforward lengthy algebra, one can infer the transformation law of the Riemannian Ricci tensor  $R_{\mu\nu}$  in  $d = 4$  under scalings  $g_{\mu\nu} \rightarrow e^{2\Omega} g_{\mu\nu}$

$$R'_{\mu\nu} = R_{\mu\nu} - 2 \nabla_\mu \nabla_\nu \Omega - g_{\mu\nu} g^{\alpha\beta} (\nabla_\alpha \nabla_\beta \Omega) + 2 (\nabla_\mu \Omega) (\nabla_\nu \Omega) - 2 g_{\mu\nu} g^{\alpha\beta} (\nabla_\alpha \Omega) (\nabla_\beta \Omega) \quad (2.15)$$

From eq-(2.13) one arrives at the Weyl invariant field equation

$$\mathcal{R}_{\mu\nu}^{Weyl}(\mathbf{g}, \mathbf{A}) = \mathcal{R}_{\mu\nu}^{Weyl}(\mathbf{g}', \mathbf{A}') = \frac{1}{2} g_{\mu\nu} \kappa \phi^2 = \frac{1}{2} g'_{\mu\nu} \kappa \phi'^2 \quad (2.16)$$

with  $\mathbf{g} = g_{\mu\nu}$ ,  $\mathbf{A} = A_\mu, \dots$ . The dimensionless parameter  $\kappa$  is inert under scalings.

The zero gauge choice  $A'_\mu = 0$  leads to

$$A'_\mu = \partial_\mu \left[ \ln \left( \frac{\phi'}{\phi_o} \right) \right] = A_\mu - \partial_\mu \Omega = \partial_\mu \left[ \ln \left( \frac{\phi}{\phi_o} \right) \right] - \partial_\mu \Omega = 0 \Rightarrow$$

$$\phi' = \phi_o; \quad e^\Omega = \frac{\phi}{\phi_o} \quad (2.17)$$

which resulted from  $D'_\mu\phi' = D_\mu\phi = 0$ .

Consequently, one arrives at

$$\mathcal{R}_{\mu\nu}^{Weyl}(\mathbf{g}', \mathbf{A}' = \mathbf{0}) = R'_{\mu\nu} = \frac{1}{2} g'_{\mu\nu} \kappa \phi_o^2 \Rightarrow R' = 2\kappa \phi_o^2 \quad (2.18)$$

leading to a family of spacetime backgrounds which are all conformally equivalent to backgrounds of *constant* Riemannian scalar curvature : (Anti) de Sitter spaces. The solutions to the scalar  $\phi$  field equation (2.12) defined in spacetime backgrounds which are conformally equivalent to a (Anti) de Sitter background

$$g_{\mu\nu} = e^{2\Omega(x)} g_{\mu\nu}^{(A)dS} = e^{-2\Omega(x)} g_{\mu\nu}^{(A)dS}, \quad \Omega = -\Omega' \quad (2.19)$$

are of the form  $\phi = e^{-\Omega'(x)} \phi_o = e^{\Omega(x)} \phi_o$ ;  $\phi_o = (16\pi G_N)^{-1/2}$  is the constant directly related to the observed Newtonian coupling  $G_N$ .

Under Weyl scalings the constant Riemannian scalar curvature of (Anti) de Sitter space in  $d = 4$  transforms as

$$g_{\mu\nu} = e^{2\Omega(x)} g_{\mu\nu}^{(A)dS} = e^{-2\Omega(x)} g_{\mu\nu}^{(A)dS}, \Rightarrow$$

$$R = e^{2\Omega(x)} \left( R'_{(A)dS} + 6 (\nabla_\mu \nabla^\mu \Omega) - 6 (\nabla_\mu \Omega) (\nabla^\mu \Omega) \right) \quad (2.20)$$

such that

$$R \phi - \frac{6}{\sqrt{|g|}} \partial_\mu \left( \sqrt{|g|} g^{\mu\nu} \partial_\nu \phi \right) - 2 \kappa \phi^3 =$$

$$e^{3\Omega(x)} \left( R'_{(A)dS} \phi_o - 2 \kappa \phi_o^3 \right) = 0 \quad (2.21)$$

as expected.

To sum up, starting with a scalar-tensor theory within the context of Weyl's geometry, permits to *derive* the expression for the potential  $V(\phi) = \kappa \phi^4$ , instead of being introduced by hand, and find the solutions  $g_{\mu\nu}, A_\mu$  to the field equations (2.8, 2.9, 2.10)

$$g_{\mu\nu}[\phi] = e^{-2\Omega} g_{\mu\nu}^{(A)dS}[\phi_o] = \left(\frac{\phi_o}{\phi}\right)^2 g_{\mu\nu}^{(A)dS}[\phi_o], \quad A_\mu[\phi] = \partial_\mu \left[ \ln\left(\frac{\phi}{\phi_o}\right) \right] \quad (2.22)$$

The (Anti) de Sitter metric  $g_{\mu\nu}^{(A)dS}[\phi_o]$  has an explicit dependence on  $\phi_o$  via the cosmological constant  $\Lambda$  :  $R' = 4\Lambda = 2\kappa\phi_o^2$ .  $\kappa < 0$  for Anti de Sitter space;  $\kappa > 0$  for de Sitter space. The solutions with  $\kappa = 0$  lead, for example, to the Schwarzschild ( $R'_{\mu\nu} = R' = 0$ ) and Reissner-Nordstrom ( $R' = 0$ ) metrics corresponding to static spherically symmetric backgrounds.

The prime example of a de Sitter background of constant Riemannian scalar curvature is the observed accelerated-expanding universe  $R' = 12H_o^2 = \frac{12}{R_H^2}$  where  $R_H$  is the present day Hubble radius. Substituting  $\phi'^2 = \phi_o^2 = (16\pi G_N)^{-1}$  and  $R' = 12H_o^2$  into eq-(2.21) yields the numerical coefficient  $\kappa$  of the potential  $V(\phi') = \kappa \phi'^4$ ,

$$12H_o^2 = 2\kappa \phi_o^2 \Rightarrow \kappa = \frac{6}{\phi_o^2 R_H^2} \quad (2.23)$$

therefore, by evaluating the potential at  $\phi' = \phi_o$

$$V(\phi_o) = \kappa \phi_o^4 = \frac{6}{\phi_o^2 R_H^2} \phi_o^4 = 6 \frac{\phi_o^2}{R_H^2} = \frac{6}{16\pi G_N R_H^2} = \frac{3}{8\pi G_N R_H^2} = \rho_{cr} \quad (2.24)$$

one recovers, in a straightforward fashion, the Universe's observed critical mass density with the precise numerical factor, and which agrees also with the observed vacuum energy density  $\rho_{vac}$ .

The boundary conditions at  $t \rightarrow \infty$  for  $\phi$  are  $\phi \rightarrow \phi_o$ , and  $(d\phi/dt) \rightarrow 0$ . Because there are an infinite number of choices<sup>2</sup> for the scaling factor  $\Omega(t)$ , the physics and experimental evidence should dictate what are the suitable expressions one should have for  $\Omega(t)$ . In other words, by choosing the specific gauge  $A'_\mu = 0 \Rightarrow A_\mu = (A_t = \partial_t \Omega(t), 0, 0, 0)$  one will break the Weyl scale invariance by fixing  $\phi' = \phi_o = (16\pi G_N)^{-1/2}$  to a constant. In order to find  $\Omega(t)$  we shall be guided by the results of the Asymptotic Safety program in quantum gravity [13] that there is a (nontrivial) interacting ultraviolet fixed point  $G_*$  for the dimensionless running gravitational coupling in the  $k \rightarrow \infty$  (infinite energy) limit (short distance limit) defined as :  $G_* = \lim_{k \rightarrow \infty} G_k(r) k^2(r) = 0 \times \infty \neq 0$ .

The limiting value of the running gravitational coupling  $G_{k=\infty}(r)$  obtained in the dynamical renormalization of black-hole spacetimes turned out to be [15]

$$G_\infty(r) = G_N (1 - e^{-r^3/r_s l_{cr}^2}) \quad (2.25)$$

$r_s$  is the classical Schwarzschild radius  $2G_N M$ , and  $l_{cr} \sim L_P$  (Planck scale) represents a critical length scale below which the modifications by the running Newtonian coupling become important [15]. The end result of this running gravitational coupling is a "renormalized" black hole spacetime of the Dymnikova-type which is free from singularities [17] and given by the metric

$$(ds)^2 = - \left(1 - \frac{2G_\infty(r)M_o}{r}\right)(dt)^2 + \left(1 - \frac{2G_\infty(r)M_o}{r}\right)^{-1}(dr)^2 + r^2(d\Omega_2)^2 \quad (2.26)$$

The  $r \rightarrow 0$  limit of  $G_\infty(r)/r$  is  $G_o r^2/r_s l_{cr}^2$  leading to a *repulsive* de Sitter core at very short distances. Similar repulsive de Sitter core were found later on in [16] by using a Gaussian profile mass density function, and in infinite derivative gravity [19].

We shall borrow the physical implications of these results, giving the short distance  $G_\infty(r=0) = 0$ , and large distance  $G_\infty(r \rightarrow \infty) \rightarrow G_N$  behavior, and apply them to the *temporal* flow of  $G(t) \sim \phi^{-1/2}(t)$ , as  $t$  runs from 0 to  $\infty$ ; i.e. as the Universe expands from the big-bang singularity (point) to the present size and beyond. In this fashion we will be able to find a judicious choice for the Weyl scaling function  $\Omega(t)$  and which is tantamount of determining the

<sup>2</sup>One could speculate that the infinite choices for  $\Omega$  bears a connection to the String Landscape paradigm. Conformal invariance is a key symmetry in String theory

expression for  $A_\mu$ , since the gauge field does not have any true dynamics (it is pure gauge).

There are many different expressions to describe the de Sitter metric depending on the coordinates being used. A flat slicing of the four-dim de Sitter space is

$$(ds)_{dS}^2 = -(dt)^2 + e^{2H_0 t} \sum_{i=1}^3 (dy_i)^2 = e^{2H_0 t} \left( -e^{-2H_0 t} (dt)^2 + \sum_{i=1}^3 (dy_i)^2 \right) \quad (2.27)$$

After performing the change of coordinates

$$\int_{\xi_\infty}^{\xi} d\xi = \int_{\infty}^t e^{-Ht} dt \Rightarrow \xi_\infty - \xi = R_H e^{-H_0 t}, \quad R_H \equiv H_0^{-1} \quad (2.28)$$

yields a conformally flat metric

$$(ds)_{dS}^2 = \frac{R_H^2}{(\xi_\infty - \xi)^2} \left( - (d\xi)^2 + \sum_{i=1}^3 (dy_i)^2 \right) \quad (2.29)$$

The next step is to select the appropriate conformal scaling factor of the above metric as indicated by eq-(2.22). Inspired by the functional form of  $G_\infty(r)$  in eq-(2.25), we can deduce the temporal dependence  $G(t)$  from the following correspondence ( $c = 1$ )

$$r \leftrightarrow t, \quad l_{cr} = L_P \leftrightarrow t_P, \quad r_s = 2G_N M \leftrightarrow R_H \quad (2.30)$$

where the Schwarzschild radius  $r_s$  (black hole horizon) corresponds now to the cosmological horizon (Hubble radius), and the Planck scale  $L_P$  corresponds to the Planck time  $t_P$ . At the end of this work we will say more about the above correspondence within the framework of black hole Cosmology [20].

Therefore, the choice of the scaling factor  $\Omega(t)$  which follows from the correspondence (2.30) associated to  $G_\infty(r)$  (2.25) is given by

$$e^{2\Omega(t)} = \left( \frac{\phi}{\phi_o} \right)^2 \equiv \frac{1}{1 - e^{-t^3/R_H t_P^2}} \Rightarrow \Omega(t) = -\frac{1}{2} \ln(1 - e^{-t^3/R_H t_P^2}) \quad (2.31)$$

from the latter expression one can deduce the temporal flow of the Weyl's gauge field

$$A_t[\phi(t)] = \partial_t \left[ \ln \left( \frac{\phi(t)}{\phi_o} \right) \right] = \partial_t \Omega(t) = -\frac{3}{2} \frac{(t^2/R_H t_P^2) e^{-t^3/R_H t_P^2}}{1 - e^{-t^3/R_H t_P^2}}, \quad (2.32)$$

and  $A_i = 0 (i = 1, 2, 3)$ , giving  $A_t(t) \rightarrow 0$ , as  $t \rightarrow \infty$ , and  $A_t \rightarrow -3/2t \rightarrow -\infty$ , as  $t \rightarrow 0$ . By construction, the flow of the gravitational coupling is

$$G(t) = G_N (1 - e^{-t^3/R_H t_P^2}), \quad G(t=0) = 0, \quad G(t \rightarrow \infty) \rightarrow G_N \quad (2.33)$$

which is compatible with the results [15] in the ultraviolet and infrared regimes.

An important remark is in order. If one were to view the net result of the temporal-dependent Weyl scaling factor in (2.31) as the introduction of an *effective* decreasing Hubble function  $H(t)$  of the form

$$H(t) = H_o + \frac{3}{2} \frac{(t^2/R_H t_P^2) e^{-t^3/R_H t_P^2}}{1 - e^{-t^3/R_H t_P^2}} \quad (2.34)$$

one encounters a problem with the integral

$$2 \int_0^t H(t') dt' = 2H_o t + \ln[1 - e^{-t^3/R_H t_P^2}] - \ln[0] = \infty \quad (2.35a)$$

because it *diverges* at  $t = 0$ . Hence, the Weyl scaling factor  $e^{-2\Omega}$  should be viewed as a “regularization” of  $\exp[2 \int_0^t (H(t') - H_o) dt']$  involving the effective Hubble function  $H(t)$  (2.34). This fact simply follows from the expression for the Weyl gauge field

$$A_t(t) = \partial_t \Omega(t) \Rightarrow \int_0^t A_{t'}(t') dt' = \Omega(t) - \Omega_o \quad (2.35b)$$

such that the divergence of the integral at  $t' = 0$  (due to the divergence of  $A_{t'}$  at  $t' = 0$ ) corresponds precisely with the *divergent* value of the constant  $\Omega_o = \Omega(t = 0) = \infty$ . Consequently, one can interpret the “Bang” of the Big Bang as the *singularity* of the Weyl gauge field at  $t = 0$ , ushering in the era of inflation.

Finally, from eqs-(2.23, 2.31) one can extract the temporal behaviour of the potential as the scalar field  $\phi$  rolls down

$$V(\phi(t)) = \kappa \phi^4(t) = \frac{6}{\phi_o^2 R_H^2} \frac{\phi_o^4}{(1 - e^{-t^3/R_H t_P^2})^2} \quad (2.36)$$

One finds then that as  $t \rightarrow \infty$ ,  $V(\phi) \rightarrow V(\phi_o) = (3/8\pi G_N R_H^2) = \rho_{vac}$ , recovering the observed vacuum energy density. Whereas at Planck’s time  $t = t_P$ , one finds after a Taylor expansion of the exponential, in  $c = 1$  units, the expected very large result

$$V(\phi(t_P)) = \frac{6}{\phi_o^2 R_H^2} \frac{\phi_o^4}{(1 - e^{-t_P^3/R_H})^2} \simeq \frac{6}{\phi_o^2 R_H^2} \frac{\phi_o^4}{(t_P/R_H)^2} = \frac{3}{8\pi} M_P^4 \quad (2.37)$$

simply by substituting  $16\pi\phi_o^2 = G_N^{-1} = M_P^2$  ( $\hbar = c = 1$ ). As the bubble expands it borrows energy from the vacuum, thus depleting its energy density to the extremely low value currently observed. The non-zero value of  $\phi = \phi_o \neq 0$  as  $t \rightarrow \infty$ , is an “attractor” point. The scalar field  $\phi$  does not reach the minimum

$\phi = 0$  of the potential. If we were to set  $\phi_o = 0 \Rightarrow G_o = G_N = \infty$  which would have been unphysical.

Note that other choices for  $\Omega(t)$  like

$$e^{-2\Omega(t)} = \left(\frac{\phi_o}{\phi}\right)^2 = 1 - e^{-(t^3/R_H t_P^2)^\beta}, \quad \beta > 1 \quad (2.38)$$

would lead to a huge vacuum energy at the Planck scale (Planck time)

$$\rho = \frac{3}{8\pi} \left(\frac{R_H}{L_P}\right)^{2\beta-2} M_P^4 \gg M_P^4 \quad (2.39)$$

Therefore,  $\beta = 1$  is the right value consistent with  $\rho \sim M_P^4$ , and the  $G_\infty(r) \leftrightarrow G(t)$  correspondence.

The scalar curvature  $R \sim e^{2\Omega}(4\Lambda) + \dots$  blows up at  $t = 0$  because  $e^{2\Omega}$  (2.31) diverges at  $t = 0$ , and which is consistent with the Big-Bang singularity. The fact that the universe emerges from a “point” is also consistent with the fact that the metric  $g_{\mu\nu} = e^{-2\Omega} g_{\mu\nu}^{dS}$  degenerates/pinches-off to *zero* at  $t = 0$ . In the late time period  $t \rightarrow \infty$ , the scaling factor  $e^{-2\Omega} \rightarrow 1$ , such that one recovers the observed de Sitter metric, with  $R = 4\Lambda = 12H_o^2 = 12(R_H)^{-2}$ , and the Newtonian gravitational coupling  $G_N$ .

This behavior should be contrasted with the Conformal Cyclic Cosmology (CCC) proposal by [31] where our Big Bang singularity corresponds to the future timelike singularity of the previous phase (cycle) of the Universe. The infinite conformal factor that pinches-off the metric to zero at  $t = 0$  in our universe, is what “squashes down” the future time like singularity of the previous phase of the universe and turns it into the spacelike singularity of our Big Bang.

Concluding, the most salient feature of all these results is that they relied solely on Weyl’s geometry and the short/large distance behavior of the running gravitational coupling in the Asymptotic Safety program of quantum gravity. The Renormalization group flow of the cosmological constant in Asymptotic Safety was studied by [14]. The scale dependence of  $\Lambda(k)$  in the de Sitter case was found to be [14]

$$\Lambda(k) = \Lambda_0 + \frac{b G(k)}{4} k^4, \quad \Lambda_0 > 0 \quad (2.40)$$

where  $b$  is positive numerical constant. In  $d = 4$ , the dimensionless gravitational coupling has a nontrivial fixed point  $g = G(k)k^2 \rightarrow g_*$  in the  $k \rightarrow \infty$  limit, and the dimensionless variable  $\lambda = \Lambda(k)k^{-2}$  has also a nontrivial ultraviolet fixed point  $\lambda_* \neq 0$  [14]. The infrared limits are  $\Lambda(k \rightarrow 0) = \Lambda_0 > 0$ ,  $G(k \rightarrow 0) = G_N$ . Where the ultraviolet limit are  $\Lambda(k = \infty) = \infty$ ;  $G(k = \infty) = 0$ .

To summarize the main result of this section, the temporal flow of the potential  $V(\phi(t))$  recaptures the same effects as the above Renormalization group flow of the cosmological constant and leads to the flow of the vacuum energy density, from the large value of  $M_{Planck}^4$  to the extremely small present value  $10^{-122} M_{Planck}^4$ , and which could provide important clues to the resolution of the cosmological constant problem.

## 2.2 Static Spherically Symmetric Solutions and Black Hole Cosmology

Having analyzed the deep cosmological implications of the *temporal* behavior of the solutions to the field equations associated with the simplest Weyl invariant action, and corresponding to a Jordan-Brans-like scalar field  $\phi$ , a metric  $g_{\mu\nu}$  and Weyl's field  $A_\mu$ , in this section we shall focus on the *spatial* behavior of the solutions, in the static spherically symmetric case.

Let us start with the renormalization-group improved black-hole metric [14]

$$(ds)^2 = - \left(1 - \frac{2G(r)M_o}{r}\right)(dt)^2 + \left(1 - \frac{2G(r)M_o}{r}\right)^{-1}(dr)^2 + r^2(d\Omega_2)^2 \quad (2.41)$$

based on the Renormalization group flow of  $G(r)$  in the Asymptotic Safety program [13],

The Einstein field equations corresponding to the above metric are  $G_\nu^\mu = 8\pi G(r)T_\nu^\mu$ , where  $T_\nu^\mu$  is in this case the effective stress energy tensor associated to the graviton quantum-self energy resulting from vacuum polarizations effects [18]. Note the presence of the running Newtonian coupling  $G(r)$  in the right hand side. A small variation of Newton's constant triggers a ripple effect of successive back reactions of the semiclassical background spacetime [15].

The limiting value of the running gravitational coupling  $G_{k=\infty}(r)$  obtained in the dynamical renormalization of the black-hole spacetime represented by the metric (2.41) turned out to be [15]

$$G_\infty(r) = G_o (1 - e^{-r^3/r_s l_{cr}^2}) \quad (2.42)$$

It *is* this value of  $G_\infty(r) \sim \phi^{-2}(r)$  that provides the radial dependence of the scalar field  $\phi(r)$ , which in turn, will select the expression for the Weyl scaling as follows

$$e^{-2\Omega(r)} = \left(\frac{\phi_o}{\phi(r)}\right)^2 = \frac{G_\infty(r)}{G_o} = 1 - e^{-r^3/r_s l_{cr}^2} \quad (2.42)$$

Given the de Sitter metric in static coordinates

$$(ds)^2 = - \left(1 - \frac{\Lambda}{3} r^2\right)(dt)^2 + \left(1 - \frac{\Lambda}{3} r^2\right)^{-1}(dr)^2 + r^2 (d\Omega_2)^2 \quad (2.43)$$

the rescaled de Sitter metric is  $g_{\mu\nu} = e^{-2\Omega(r)} g_{\mu\nu}^{dS}$ . And by construction, eq-(2.21) will be satisfied for the scalar field  $\phi(r)$  whose functional dependence can be read directly from eq-(2.42), and where the de Sitter metric is displayed in eq-(2.43).

As  $r \rightarrow 0$ ,  $G_\infty(r) \rightarrow 0$ . As  $r \rightarrow \infty$ ,  $G_\infty(r) \rightarrow G_o = G_N$ . The scaling factor vanishes at  $r = 0$ , so the rescaled de Sitter metric pinches off to zero, to a "point", and the scalar curvature blows up, like in the Big-Bang at  $t = 0$ . At  $r = \infty$  one recovers the de Sitter metric (2.43) with a constant scalar

Riemannian curvature  $R = 4\Lambda$ . In this fashion one obtains similar results as in section **2.1**.

In passing, we deem it important to point out that the metric (2.43) is not the only one which furnishes a constant scalar curvature. To proceed let us focus on the following spacetime background (in natural  $c = 1$  units)

$$(ds)^2 = - \left(1 - \frac{2G_o\mathcal{M}(r)}{r}\right)(dt)^2 + \left(1 - \frac{2G_o\mathcal{M}(r)}{r}\right)^{-1}(dr)^2 + r^2(d\Omega_2)^2 \quad (2.44)$$

related to an anisotropic self gravitating fluid droplet [16].

The energy-momentum tensor corresponding to the Einstein equations  $G_\nu^\mu = 8\pi G_o T_\nu^\mu$ , and associated to the metric (2.44) is given by

$$T_\nu^\mu \equiv \text{diag} (-\rho(r), p_r(r), p_\theta(r), p_\varphi(r)) \quad (2.45)$$

physically it represents a self-gravitating anisotropic fluid (bubble) whose mass density and pressure components are

$$\rho = - p_r = \frac{1}{4\pi r^2} \frac{d\mathcal{M}}{dr}, \quad p_\theta = p_\varphi = - \frac{1}{8\pi r} \frac{d^2\mathcal{M}}{dr^2} \quad (2.46)$$

From eqs-(2.44,2.45) one learns that

$$p_\theta = p_\varphi = - \rho - \frac{r}{2} \frac{d\rho}{dr} \quad (2.47)$$

and which is consistent with the local conservation law  $\nabla_\mu T_\nu^\mu = 0$

In the case of a point mass  $M_o$  the above equations lead to

$$\rho = \frac{M_o\delta(r)}{4\pi r^2} = - p_r, \quad p_\theta = p_\varphi = - \frac{M_o}{8\pi r} \delta'(r) = \frac{M_o\delta(r)}{8\pi r^2} \quad (2.48)$$

after using the relation  $r\delta'(r) = -\delta(r)$ . The scalar curvature is

$$R = - 2G_o \left( \frac{(d^2\mathcal{M}/dr^2)}{r} + 2 \frac{(d\mathcal{M}/dr)}{r^2} \right) \quad (2.49)$$

consistent with taking the trace of the field equations  $R = -8\pi G_o T$ .

When the metric background is provided by (2.44), the scalar curvature is

$$R[\mathcal{M}(r)] = - 2G_o \left( \frac{(d^2\mathcal{M}(r)/dr^2)}{r} + 2 \frac{(d\mathcal{M}(r)/dr)}{r^2} \right) \quad (2.50)$$

and the differential equation to solve for  $\mathcal{M}(r)$  is the following Euler-Cauchy equation

$$\frac{(d^2\mathcal{M}(r)/dr^2)}{r} + 2 \frac{(d\mathcal{M}(r)/dr)}{r^2} = C = - \frac{4\Lambda}{2G_o} \quad (2.51)$$

after setting the scalar curvature to a constant equal to  $4\Lambda$ .

The solution to (2.51) is

$$\mathcal{M}(r) = \frac{C}{12} r^3 + \frac{A}{r} + B \quad (2.52)$$

and the metric (2.44) ends up by having the same functional *form* as the Schwarzschild-(Anti)de Sitter-Reissner-Nordstrom metric, whose temporal  $g_{tt}$  and radial components  $g_{rr}$  are

$$g_{tt} = -\left(1 - \frac{2G_o M_o}{r} + \frac{Q^2 G_o}{r^2} \pm \frac{\Lambda}{3} r^2\right), \quad g_{rr} = -g_{tt}^{-1} \quad (2.53)$$

after the following identification among the parameters is made

$$B = M_o, \quad -\frac{G_o C}{2} = \pm \Lambda, \quad -2A = Q^2 \quad (2.54)$$

Since there is no mass nor charge in the real scalar-tensor field action (2.1), by setting  $M_o = 0, Q^2 = 0$  one recovers the (Anti) de Sitter metric. The existence of a cosmological de Sitter horizon at the Hubble scale  $1 - \frac{\Lambda}{3} R_H^2 = 0$  fixes then the value of  $\Lambda = \frac{3}{R_H^2}$ . Had  $M_o \neq 0$ , and/or  $Q^2 \neq 0$ , the value of  $\Lambda$  would have been different from  $\frac{3}{R_H^2}$  if  $R_H$  still remained as the cosmological horizon.

To finalize, we will discuss the connection that the Dymnikova-type of metric [17] has with black hole cosmology. Although there is a mathematical equivalence in writing  $G_\infty(r)M_o \leftrightarrow G_o\mathcal{M}(r)$ , when the mass function is  $\mathcal{M}(r) = M_o(1 - \exp(-r^3/r_s L_P^2))$ , there is a clear physical difference. In the former, one has a point mass  $M_o$  at  $r = 0$  and a running gravitational coupling [15]. In the latter [17] one has the Newtonian coupling  $G_o = G_N$  and a mass profile distribution as if the point mass  $M_o$  were smeared all over space. The importance of the Dymnikova solution is that one can rewrite

$$\frac{2G_o\mathcal{M}(r)}{r} = \frac{\Lambda(r)}{3} r^2 \quad (2.55)$$

in terms of a running cosmological ‘‘constant’’  $\Lambda(r)$ . At the Planck scale, one has a de Sitter core with a Planck size throat size and a with a very large value of  $\Lambda(L_P) \sim \frac{1}{L_P^2}$ . Whereas at the Hubble scale, one may rewrite the asymptotic Schwarzschild behavior of the metric in the form

$$\frac{2G_o M_o}{R_H} = \frac{2G_o M_o}{R_H} \frac{1}{R_H^2} R_H^2 = \frac{\Lambda(R_H)}{3} R_H^2 \quad (2.56)$$

with  $\Lambda(R_H) = \frac{3}{R_H^2}$ , if, and only if,  $M_o$  obeys  $\frac{2G_o M_o}{R_H} = 1$ , and which is tantamount to viewing the Universe as a black hole whose mass  $M_U = M_o$  is enclosed inside the cosmological horizon  $R_H$ , that matches also its Schwarzschild radius  $r_s = 2G_o M_o = R_H$ . When this occurs, the uniform density over a spherical ball of radius  $R_H$  given by  $M_o/(4\pi/3)R_H^3 = \frac{3}{8\pi G_o R_H^2}$  coincides precisely with

the observed critical density (also vacuum density) of our universe. For more details of black hole cosmology see [20].

Concluding, although the numerical findings of Asymptotic safety have been confined to  $d = 4$ , in principle we should expect the results of this work to admit an extension to other dimensions. If not this would signal four dimensions  $d = 4$  as special. In a few words, the study of a Jordan-Brans-Dicke-like action (2.1) within the context of Weyl geometry, combined with the findings of Asymptotic Safety in quantum gravity, leads to a plethora of nice numerical results :

(i) like singling out the quartic potential from all the others; (ii) having (Anti) de Sitter space as the most natural solution; (iii) furnishing the value of the observed vacuum energy density at the Hubble scale  $10^{-122} M_P^4$ ; (iv) a  $M_P^4$  vacuum energy density at the Planck scale; (v) interpreting the “Bang” of the Big Bang as the *singularity* of the Weyl gauge field of dilations at  $t = 0$  ushering in the era of inflation, and (vi) allowing the possibility that our universe is a black hole whose horizon coincides with the cosmological Hubble horizon.

Since (Anti) de Sitter space was an instrumental solution to the field equations (2.8-2.10), it is warranted to explore deeper the interplay among Weyl geometry, Asymptotic safety and the *AdS/CFT* correspondence (holographic renormalization group flow). Also relevant is the work by [5] on the role of dilatation symmetry in higher dimensions and the vanishing of the cosmological constant. Last, but not least, we should also consider the implications of Conformal Cyclic Cosmology [31] and Scale Relativity Theory [26] with the key findings of this work.

### Acknowledgements

We thank M. Bowers for assistance and to Raymond Aschheim for performing some numerical solutions of a nonlinear differential equation.

## References

- [1] H. Weyl, 1918a. “Gravitation und Elektrizitat.” Sitzungsberichte der Koniglich Preussischen Akademie der Wissenschaften zu Berlin pp. 465. GA II, 29-42, [31]. In (Weyl 1968, II, 29-42) [31], English in (O’Raifeartaigh 1997, 24-37).  
H. Weyl, 1918b. Raum, - Zeit - Materie. Berlin etc.: Springer. Weitere Auflagen: 21919, 31919, 41921, 51923, 61970, 71988, 81993. English and French translations from the 4th ed. in 1922.  
H. Weyl, 1918c. “Reine Infinitesimalgeometrie”. Mathematische Zeitschrift 2:384-411. In (Weyl 1968, II, 1-28).
- [2] E. Scholz, “Weyl Geometry in late 20-th Century Physics”, arXiv : 1111.3220 [math.HO]

- E. Scholz, “On the geometry of cosmological model building”, arXiv:grqc/0511113.
- E. Scholz, “Cosmological spacetimes balanced by a Weyl geometric scale covariant scalar field”, *Foundations of Physics* **39** (2009) 45.
- [3] M. Israelit and N. Rosen, “Cosmic dark matter and Dirac gauge function. *Foundations of Physics* **25** (1995) 763.
- M. Israelit and N. Rosen, “A Weyl-Dirac geometric particle. *Foundations of Physics* **26** (1996) 585.
- M. Israelit, “Primary matter creation in a Weyl-Dirac cosmological model, *Foundation of Physics* **32** (2002) 295.
- M. Israelit, “Quintessence and dark matter created by Weyl-Dirac geometry, *Foundation of Physics* **32** (2002) 945.
- [4] W. Kao, “Inflationary solution in Weyl invariant theory”, *Phys. Letts A* **149** (1990) 76.
- [5] C. Wetterich, “Quantum scale symmetry”, arXiv : 1901.04741.
- C. Wetterich “Dilatation symmetry in higher dimensions and the vanishing of the cosmological constant” *Phys. Rev. Lett.* **102** (2009) 141303.
- [6] S. Cree, T. Davis, T. Ralph, Qi. Wang, Z. Zhu, and W. Unruh “Can the fluctuations of the quantum vacuum solve the cosmological constant problem?” *Phys. Rev. D* **98**, 063506 (2018)
- [7] L. Smolin, “Towards a theory of spacetime structure at very short distances”, *Nuclear Physics B* **160** (1979) 253.
- [8] P. Mannheim, and D. Kazanas, “Exact vacuum solution to conformal Weyl gravity and galactic rotation curves”, *Astrophysical Journal* **342** (1989) 635.
- [9] C. Castro, “The cosmological constant and Pioneer anomaly from Weyl geometry and Mach’s principle”, *Physics Letters B* **675** (2009) 226.
- [10] S. Weinberg, *Cosmology* (Oxford University Press, 2008).
- [11] H. Kim, ”Can the Brans-Dicke gravity possibly with  $\Lambda$  be a theory of Dark matter ? ” astro-ph/0604055.
- H. Kim, “Brans-Dicke theory as a unified model for dark matter dark energy”, *Monthly Notices Royal Astronomical Society* 364 (2005) 813.
- M. Arik and M. Calik, ”Can Brans-Dicke scalar field account for dark energy and dark matter ? gr-qc/0505035.
- Chao-Jun Feng, ”Ricci Dark Energy in Brans-Dicke Theory” arXiv : 0806.0673.

- [12] S. Capozziello, V. Cardone and A. Troisi, " Jour. of Cosmology and Astroparticle Physics **08** (2006) 001.  
S. Capozziello, S. Nojiri and S. Odintsov, Phys. Letts **B 634** (2006) 93.
- [13] S. Weinberg, "Ultraviolet divergences in quantum theories of gravitation" *General Relativity : An Einstein centenary survey*, Eds. S. Hawking and W. Israel, Cambridge University Press (1979) p. 790.
- [14] A. Bonano and Reuter. "Renormalization group improved black hole spacetimes" Phys. Rev. **D62** (2000) 043008.  
M. Reuter and F. Saueressig, "Quantum Einstein Gravity" arXiv: 1202.2274  
R. Percacci, "A short introduction to Asymptotic Safety" arXiv: 1110.6389.  
D. Litim, "Renormalization group and the Planck scale" arXiv: 1102.4624.  
S. Nagy, "Lectures on Renormalization and Asymptotic Safety", arXiv : 1211.4151.  
M. Niedermaier, "The Asymptotic Safety Scenario in Quantum Gravity-An Introduction" , gr-qc/0610018.
- [15] A. Platania, "Dynamical renormalization of black-hole spacetime" arXiv : 1903.10411.
- [16] E. Spallucci and A. Smailagic, "Regular Black Holes from semi-classical down to Planck size", Int. J. Mod. Phys. **D 26**, n.7, 1730013, (2017) [27 pages]  
P. Nicolini, E. Spallucci, M. F. Wondrak, "Quantum Corrected Black Holes from String T-Duality", arXiv : 1902.11242.
- [17] I. Dymnikova, "Spherically Symmetric Spacetime with Regular de Sitter Center", Gen. Relat. Grav **24** (1992) 235.
- [18] E. Poisson and W. Israel, Class. Quan. Grav. **5** (1988) L 201.
- [19] A. S. Cornell, G. Harmsen, G. Lambiase, A. Mazumdar, "Rotating metric in Non-Singular Infinite Derivative Theories of Gravity" arXiv : 1710.02162.
- [20] R. Pathria, "The Universe as a Black Hole". Nature. **240** (5379): 298-299.  
I. Goody, "Chinese universes". Physics Today. **25** (1972) 7.  
P. Landsberg, "Mass Scales and the Cosmological Coincidences". Annalen der Physik. **496** (2) (1984) 88-92.  
Black Hole Cosmology, [http://en.wikipedia.org/wiki/Black\\_hole\\_cosmology](http://en.wikipedia.org/wiki/Black_hole_cosmology)
- [21] E. Santamato, "Geometric derivation of the Schrödinger equation from classical mechanics in curved Weyl spaces", Phys. Rev. **D 29**, (1984) 216.

- [22] C. Castro, “Bohm’s Potential, Classical/Quantum Duality and Repulsive Gravity” *Phys. Letters B* (2018), doi.org/10.1016/j.physletb.2018.11.013.
- [23] V. Faraoni, *Cosmology in Scalar-Tensor Gravity* (Kluwer Academic Publishers, 2003). Y. Fujii and K. Maeda, *The Scalar-Tensor Theory of Gravitation* (Cambridge University Press 2003).
- [24] F. Hehl, J. McCrea, E. Mielke and Y. Ne’eman, “Metric-affine gauge theory of gravity: Field equations, Noether identities, world spinors, and breaking of dilation invariance. *Phys. Reports* **258** (1995) 1.
- [25] C. Castro, “Developments of the Relativistic Bohm-Poisson Equation and Dark Energy” *Theoretical Physics* **Vol. 4**, no. 1, (March 2019), 26-39; <https://dx.doi.org/10.22606/tp.2019.41003>.
- [26] L. Nottale, *Scale Relativity And Fractal Space-Time: A New Approach to Unifying Relativity and Quantum Mechanics* (World Scientific 2011)  
L. Nottale, *Fractal Space-Time and Micro-physics* (World Scientific 1993).
- [27] D. Giulini and A. Grossardt, “The SchrodingerNewton equation as a non-relativistic limit of self-gravitating KleinGordon and Dirac fields. *Classical and Quantum Gravity* **29**, no. 21 (2012) 215010.  
S. Khanapurkar, A. Pradhan, V. Dhruv, T. P. Singh “Non-relativistic limit of Einstein-Cartan-Dirac equations” *Phys. Rev. D* **98**, 104027 (2018)
- [28] A. Guth, ”Inflationary universe: A possible solution to the horizon and flatness problems ” *Phys. Rev. D.* **23** (2) (1981) 347.  
E. Fahri and A. Guth, *Phys. Letts B* **183** (1987) 149.
- [29] A. Maeder, “Dynamical effects of the scale invariance of the empty space: The fall of dark matter ? ” arXiv : 1711.07740.  
”An Alternative to the  $\Lambda$ CDM Model: The Case of Scale Invariance”. *The Astrophysical Journal.* **834** (2) (2017) 194.
- [30] F. K. Anagnostopoulos, D. Benisty, S. Basilakos, E. I. Guendelman “Dark energy and dark matter unification from dynamical space time: observational constraints and cosmological implications”, arXiv : 1904.05762.
- [31] V.G. Gurzadyan, and R. Penrose, “On CCC-predicted concentric low-variance circles in the CMB sky”. *Eur. Phys. J. Plus.* **128** (2) (2013) 22.