

On the Emergence of Spacetime Dimensions from Kolmogorov Entropy

Ervin Goldfain

Abstract

This short pedagogical report is based on a couple of premises. First, it was recently shown that the long run of non-equilibrium Renormalization Group flows is prone to end up on strange attractors. As a result, multifractals are likely to provide the proper framework for the characterization of effective field theories. Secondly, it is known that multifractal analysis uses the Kolmogorov entropy (K-entropy) to quantify the degree of disorder in chaotic systems and turbulent flows. Building on these premises, the report details the remarkable connection between K-entropy, multifractal sets and spacetime dimensions. It also supports the proposal that near and above the Fermi scale, spacetime is defined by continuous and arbitrarily small deviations from four-dimensions.

Key words: Kolmogorov entropy, multifractals, minimal fractal manifold, effective field theory.

We have conjectured in [1, 2] that the flow from the ultraviolet (UV) to the infrared (IR) sector of any multidimensional nonlinear field theory approaches chaotic dynamics in a universal way. This result stems from several independent routes to aperiodic behavior and implies that the IR attractor of effective field theories is likely to replicate the properties of *strange attractors* and *multifractal sets*. In particular, the chaotic behavior of the Renormalization Group flow near or above the Fermi scale suggests that phenomena on or above this scale mimic the dynamics on a strange attractor [3-5].

Let a generic UV to IR trajectory be described by the n -dimensional phase-space flow $x(\tau)$. Here, τ denotes the evolution parameter (“time”) corresponding to the Renormalization Group scale μ

$$\tau = \log\left(\frac{\mu}{\mu_0}\right) \quad (1)$$

The random behavior of the flow near the strange attractor can be characterized by dividing the phase-space into n - dimensional hypercubes of side r , which are sampled at discrete time intervals $\Delta\tau$. The generalized K -entropy of order $q \neq 1$ is given by the equation [6]

$$K_q(X) = -\lim_{r \rightarrow 0} \lim_{\Delta\tau \rightarrow 0} \lim_{N \rightarrow \infty} \frac{1}{N\Delta\tau} \frac{1}{q-1} \ln \sum_{i_1, i_2, \dots, i_M}^{M(r)} p_{i_1, i_2, \dots, i_M}^q \quad (2)$$

where $X = x_i$ is the discrete random variable, that is, $x_i = x(\tau = i\Delta\tau)$, and p_{i_1, i_2, \dots, i_M} stands for the joint probability that the trajectory $x(\tau = \Delta\tau)$ is in i_1 , $x(\tau = 2\Delta\tau)$ is in i_2 and $x(\tau = M\Delta\tau)$ is in i_M . The K-entropy defines the asymptotic scenario where $r \rightarrow 0$ and the phase-space is sampled with an infinite number of steps ($N \rightarrow \infty$) at vanishing time intervals ($\Delta\tau \rightarrow 0$). In the special case $N\Delta\tau = 1$ and when the joint probability is constant across all hypercubes ($M(r) = \text{const.}$, $p_{i_1, i_2, \dots, i_M} = p_i$), (2) turns into the *Rényi entropy* in the logarithm base b , which assumes the form

$$S_q(X) = \frac{1}{1-q} \log_b \left(\sum_{i=1}^M p_i^q \right) \quad (3)$$

Furthermore, (3) reduces to the familiar *thermodynamic entropy* when $q \rightarrow 1$ and Boltzmann constant is set to $k_B = 1$ [7]

$$S(X) = -\sum_{i=1}^M p_i \ln p_i \quad (4)$$

Finally, the concept of *generalized dimension* of order q is introduced in conjunction with (3) according to

$$D_q = \lim_{r \rightarrow 0} \frac{1}{1-q} \frac{\log_b \left(\sum_{i=1}^M p_i^q \right)}{\log r} \quad (5)$$

A particularly straightforward expression of (3) is obtained for the null order $q=0$ and the natural logarithm base. It is referred to as *topological entropy* and is given by

$$S_0(r) = \ln \sum_{i=1}^M p_i^0 = \ln M \quad (6)$$

It is known that the *box-counting dimension* of a fractal object of normalized size r is defined as

$$D_0 \approx \frac{\ln M}{\ln r} \Rightarrow M \approx r^{D_0} = \varepsilon^{-D_0} \quad (7)$$

in which M denotes the number of covering boxes and $\varepsilon = r^{-1}$ is the normalized size of the box. The dimension of ordinary Euclidean space corresponds to integer and positive-definite values of the box-counting dimension, $D_0 = k$, $k = 0, 1, 2, \dots$.

Comparing (6) to (7) leads to the connection between the box-counting dimension and topological entropy via

$$\boxed{\varepsilon^{-D_0} = \exp[S_0(r)]} \quad (8)$$

Two straightforward conclusions may be drawn from (8):

- Maximal topological entropy ($S_0(r) \rightarrow \infty$) matches the limit $\varepsilon \rightarrow 0$ and corresponds to the four-dimensional continuum of both General Relativity and Quantum Field Theory.
- The steady increase of topological entropy along the flow implies that, near or above the Fermi scale, spacetime is endowed with a *continuous* spectrum of dimensions ($\varepsilon = 4 - D_0 \ll 1$), asymptotically reaching $D_0 = 4$ as $\varepsilon \rightarrow 0$ [8].

References:

[1] <https://www.prespacetime.com/index.php/pst/article/view/1244>

A copy of this article can be found at:

[https://www.academia.edu/38764569/Multifractal Analysis and the Dynamics of Effective Field Theories](https://www.academia.edu/38764569/Multifractal_Analysis_and_the_Dynamics_of_Effective_Field_Theories)

[2] Available at the following site (*in progress*):

[https://www.academia.edu/38852586/The Strange Attractor Structure of Turbulence and Effective Field Theories fourth draft](https://www.academia.edu/38852586/The_Strange_Attractor_Structure_of_Turbulence_and_Effective_Field_Theories_fourth_draft)

[3] Available at the following site:

[https://www.academia.edu/38735370/Chaotic Dynamics of the Renormalization Group Flow and Standard Model Parameters](https://www.academia.edu/38735370/Chaotic_Dynamics_of_the_Renormalization_Group_Flow_and_Standard_Model_Parameters)

[4] Available at the following site (*in progress*):

[https://www.academia.edu/38744111/Bifurcations and the Dynamic Content of Particle Physics](https://www.academia.edu/38744111/Bifurcations_and_the_Dynamic_Content_of_Particle_Physics)

[5] <https://arxiv.org/pdf/hep-th/0304178.pdf>

[6] <https://www.sciencedirect.com/science/article/pii/S0898122113000345>

[7] <https://arxiv.org/pdf/cond-mat/0207707.pdf>

[8] <http://www.aracneeditrice.it/index.php/pubblicazione.html?item=9788854889972>

Also available at the following site:

<https://www.researchgate.net/publication/278849474> Introduction to Fractional Field Theory consolidated version