

# A four circle problem and division by zero

HIROSHI OKUMURA  
 Maebashi Gunma 371-0123, Japan  
 e-mail: hokmr@yandex.com

**Abstract.** We generalize a problem involving four circles and a triangle, and consider some limiting cases of the problem by division by zero.

**Keywords.** four circle problem, division by zero.

**Mathematics Subject Classification (2010).** 01A27, 51M04

## 1. INTRODUCTION

We generalize the following problem involving four circles and a triangle in [20]. The same sangaku problem was proposed in 1826 and cited in [19] and [1] with no solution. Some limiting cases of the problem will be considered by division by zero [6].




Figure 1.

**Problem 1.** For a triangle  $EFG$  with incircle  $\alpha$ ,  $\delta$  is the circle passing through  $E$  and  $F$  and touching  $\alpha$ ,  $\gamma$  is the incircle of the curvilinear triangle made by  $\delta$  and the sides  $EF$  and  $GE$ , and  $\beta$  is the circle touching  $\delta$  and  $FG$  at the midpoint from the side opposite to  $\alpha$ . Let  $a$ ,  $b$  and  $c$  be the radii of  $\alpha$ ,  $\beta$  and  $\gamma$ , respectively. Show  $a^2 = 4bc$ .

A similar sangaku problem considering the case  $|EF| = |GE|$  can be found in [2, p. 302].

## 2. GENERALIZATION

The problem assumes that  $\alpha$  is the incircle of  $EFG$ , but we show that the condition is inessential. We consider the following figure (see Figure 2): For a chord  $FG$  of a circle  $\delta$ ,  $M$  is the midpoint of  $FG$ ,  $\beta$  is a circle touching  $\delta$  and  $FG$  at  $M$ ,  $\alpha$  is a circle touching  $\delta$  and the chord  $FG$  from the side opposite to  $\beta$ ,  $f$  and  $g$  are the tangents of  $\alpha$  from the points  $F$  and  $G$ , respectively,  $\gamma$  is the circle lying on the same side of  $FG$  as  $\alpha$  and touching  $\delta$  externally and  $f$  and  $g$  from the same side

as  $\alpha$ . Let  $a$ ,  $b$ ,  $c$  and  $d$  be the radii of  $\alpha$ ,  $\beta$ ,  $\gamma$  and  $\delta$ , respectively. We denote this configuration by  $\mathcal{S}$ .




Figure 2: The configuration  $\mathcal{S}$ .




Figure 3:  $4b = a = c$

We use a rectangular coordinate system with origin  $M$  such that the center of  $\alpha$  has coordinate  $(x_a, a)$  for a real number  $x_a$ . Firstly we consider a special case in which  $f$  and  $g$  are parallel (see Figure 3).

**Theorem 1.** *The following statements are equivalent for  $\mathcal{S}$ .*

- (i) *The lines  $f$  and  $g$  are parallel.*
- (ii) *The center of  $\alpha$  lies on the circle of diameter  $FG$ .*
- (iii)  *$a = 4b$ .*

*Proof.* We may assume that the point  $G$  has coordinates  $(k, 0)$ , and  $f$  and  $g$  have equations  $x + m_1y + k = 0$  and  $x + m_2y - k = 0$ , respectively for real numbers  $m_1$  and  $m_2$ . Since  $f$  and  $g$  touch  $\alpha$ , we have

$$(1) \quad m_1 = \frac{a^2 - (k + x_a)^2}{2a(k + x_a)}, \quad m_2 = -\frac{a^2 - (k - x_a)^2}{2a(k - x_a)}.$$

Notice that  $k^2 - x_a^2 \neq 0$ , since  $k^2 - x_a^2 = 0$  implies that  $\alpha$  touches  $FG$  at  $F$  or  $G$ . The lines  $f$  and  $g$  are parallel if and only if  $m_1 = m_2$ , which is equivalent to

$$(2) \quad a^2 + x_a^2 = k^2.$$

This proves the equivalence of (i) and (ii), since the left side equals the square of the distance between the center of  $\alpha$  and  $M$  (see Figure 3). While the square of the distance between the centers of  $\delta$  and  $\alpha$  equals

$$(3) \quad x_a^2 + (d - 2b - a)^2 = (d - a)^2.$$

And the power of the origin with respect to  $\delta$  equals

$$(4) \quad -2b(2d - 2b) = -k^2.$$

Eliminating  $d$  from (3) and (4), we get  $xa^2 + 4ab = k^2$ , which implies

$$a^2 + xa^2 - k^2 = a(a - 4b).$$

Hence (2) and  $a = 4b$  are equivalent, i.e., (i) and (iii) are equivalent.  $\square$

**Corollary 1.** *One of the three relations  $4b < a < c$ ,  $4b = a = c$ ,  $4b > a > c$  holds for  $\mathcal{S}$ .*

Figures 2, 3 and 4 show the cases  $4b > a > c$ ,  $4b = a = c$  and  $4b < a < c$ , respectively. The next theorem is a generalization of Problem 1.

**Theorem 2.** *The following statements hold.*

- (i) *The relation  $a^2 = 4bc$  holds.*
- (ii) *One of the internal common tangents of  $\alpha$  and  $\gamma$  is parallel to  $FG$ .*

*Proof.* We use the same notation as in the proof of Theorem 1. If  $f$  and  $g$  are parallel, we get  $a = c$ . Therefore we get  $a^2 = 4bc$  by Theorem 1. We assume that  $f$  and  $g$  intersect. We denote the point of intersection by  $E$ , which has coordinates

$$(5) \quad (x_e, y_e) = \left( \frac{k(m_1 + m_2)}{m_1 - m_2}, \frac{-2k}{m_1 - m_2} \right).$$

Substituting (1) in (5), we get

$$(6) \quad (x_e, y_e) = \left( x_a - \frac{2a^2 x_a}{a^2 - k^2 + x_a^2}, 2a - \frac{2a^3}{a^2 - k^2 + x_a^2} \right).$$

The square of the distance between the centers of  $\delta$  and  $\gamma$  equals

$$(7) \quad x_c^2 + (d - 2b - y_c)^2 = (c + d)^2,$$

where  $(x_c, y_c)$  are the coordinates of the center of  $\gamma$ . Since  $E$  is the external center of similitude of  $\alpha$  and  $\gamma$ , we get

$$(8) \quad \frac{-cx_a + ax_c}{a - c} = x_e, \quad \frac{-ca + ay_c}{a - c} = y_e.$$

Eliminating  $x_a, x_c, y_c, x_e, y_e$  and  $d$  from (3), (4), (6), (7), (8), we get

$$(a^2 - 4bc)j(k) = 0,$$

where  $j(k) = 4(a - 4b)b^2 - (4b - c)k^2$ . If  $j(k) = 0$ , we have  $k^2 = 4(a - 4b)b^2 / (4b - c) > 0$ . This implies  $a < 4b < c$  or  $c < 4b < a$ . However this contradicts Corollary 1. Therefore we get  $j(k) \neq 0$ , which implies  $a^2 = 4bc$ .

We prove (ii). If  $f$  and  $g$  are parallel, the centers of  $\alpha, \gamma$  and  $M$  are collinear, i.e.,  $x_a/a = x_c/y_c$ . Eliminating  $b, c, k, x_a, x_c$  from the equations  $x_a/a = x_c/y_c, a = c, a = 4b$ , (3), (4) and (7), we get

$$(3a - y_c)((a + 4d)a + (4d - a)y_c) = 0.$$

Therefore we get  $y_c = 3a = 2a + c$ , since  $(4d - a)y_c > 0$ . If  $f$  and  $g$  intersect, we eliminate  $b, k, x_a, x_c, x_e, y_e$  from (3), (4), (6), (7), (8). Then we get

$$(2a + c - y_c)((a + 4d)c + (4d - a)y_c) = 0.$$

Therefore we get  $y_c = 2a + c$ . This proves (ii). □




Figure 4: The configuration  $\mathcal{S}$  in the case  $4b < a < c$ .

There are several sangaku problems stating the next corollary [2, p. 312, p. 317, p. 419] (see Figure 5).

**Corollary 2.** For a semicircle  $\delta$  with diameter  $FG$ , let  $\alpha$  be the circle of radius  $a$  touching  $\delta$  and  $FG$  at the midpoint. If  $c$  is the inradius of the curvilinear triangle made by  $\delta$  and the tangents of  $\alpha$  from the points  $E$  and  $F$ , then  $a = 4c$ .




Figure 5.




Figure 6.

The next corollary can be found in the sangaku hung in 1830 [3, p. 40], which is incorrectly cited in [1, p. 34] (see Figure 6).

**Corollary 3.** For the configuration  $\mathcal{S}$ , let  $\alpha'$  be a circle of radius  $a'$  touching the circle  $\delta$  and its chord  $FG$  from the side opposite to  $\alpha$ . If the inradius of the curvilinear triangle made by  $\delta$  and the tangents of  $\alpha'$  from  $F$  and  $G$  equals  $c'$ , then  $a^2 a'^2 = cc' |FG|^2$ .

*Proof.* Let  $b'$  be the radius of the circle touching  $\delta$  and  $FG$  at the midpoint from the side opposite to  $\alpha'$ . Then we have  $a'^2 = 4b'c'$ , while  $|FG|^2 = 16bb'$  and  $a^2 = 4bc$ . Eliminating  $b$  and  $b'$  from the three equations, we get  $a^2 a'^2 = |FG|^2 cc'$ .  $\square$

## 3. LIMITING CASES WITH DIVISION BY ZERO

In this section we fix the circle  $\delta$  for  $\mathcal{S}$  and consider the case where one of the circles  $\alpha$  and  $\beta$  has radius 0 with the definition of division by zero [6]:

$$(9) \quad \frac{z}{0} = 0 \text{ for a complex number } z.$$

Notice that the definition implies that lines have radius 0 as circles [17].

We now consider a simple case in which the centers of  $\alpha$ ,  $\beta$  and  $\gamma$  are collinear for  $\mathcal{S}$  and use a rectangular coordinate system with origin at the point of tangency of  $\beta$  and  $\delta$  such that the center of  $\delta$  has coordinates  $(0, d)$ . The point of tangency of  $\gamma$  and  $\delta$  and the tangent of  $\delta$  at the point are denoted by  $D$  and  $t$ , respectively (see Figure 7). Notice that  $d = a + b$ .




Figure 7.

**3.1. The case  $b = 0$ .** Firstly we consider the case  $b = 0$ . Then  $\beta$  is a point or a line. The circle  $\alpha$  has an equation  $x^2 + (y - (b + d))^2 = (b - d)^2$ , which is arranged as

$$(10) \quad f_a(x, y) = (x^2 + (y - d)^2 - d^2) + 2b(2d - y) = 0.$$

From  $f_a = 0$ , we get  $x^2 + (y - d)^2 = d^2$  in the case  $b = 0$ . Also from  $f_a/b = 0$  we get  $y = 2d$  in the case  $b = 0$  by (9). Hence  $\alpha$  coincides with the circle  $\delta$  or the line  $t$  in the case  $b = 0$ .

The circle  $\beta$  has an equation

$$f_b(x, y) = (x^2 + y^2) - 2by = 0.$$

From  $f_b = 0$  we get  $x^2 + y^2 = 0$  in the case  $b = 0$ . Also from  $f_b/b = 0$  we get  $y = 0$  in the case  $b = 0$  by (9). Hence  $\beta$  coincides with the origin or the  $x$ -axis in this case.

The circle  $\gamma$  has an equation  $x^2 + (y - 2d - c)^2 = c^2$ , where  $c = (d - b)^2/(4b)$ , which is arranged as.

$$f_c(x, y) = \frac{d^2}{2b}(2d - y) + \left( x^2 + \left( y - \frac{3d}{2} \right)^2 - \frac{d^2}{4} \right) + \frac{b}{2}(2d - y) = 0.$$

From  $f_c = 0$  we get  $x^2 + (y - 3d/2)^2 = (d/2)^2$  in the case  $b = 0$ . Also from each of  $f_c b = 0$  and  $f_c/b = 0$  we get  $y = 2d$  in the case  $b = 0$ . Hence  $\gamma$  coincides with the line  $t$  or the circle of radius  $d/2$  touching  $\delta$  at  $D$  in this case.

When  $\beta$  approaches to the origin, the circles  $\alpha$  and  $\gamma$  approach to  $\delta$  and  $t$ , respectively. Therefore we can consider that  $\alpha$  and  $\gamma$  coincide with  $\delta$  and  $t$ , respectively when  $\beta$  degenerates to the origin, (see Figure 8). The relation  $a^2 = 4bc$  does not holds in this case, but  $a^2/b = 4c$  and  $a^2/c = 4b$  hold by (9), since the radius of  $t$  equals 0.

When  $\beta$  coincides with the  $x$ -axis, we can thereby consider that  $\alpha$  and  $\gamma$  coincides with  $t$  and the circle of radius  $d/2$  touching  $\delta$  at  $D$ , respectively as the remaining case (see Figure 9). The relation  $a^2 = 4bc$  holds in this case.




Figure 8.




Figure 9.

case	$\alpha$	$\beta$	$\gamma$	relation of the radii
1	$\delta$	origin	$t$	$a^2/b = 4c, a^2/c = 4b$
2	$t$	$x$ -axis	circle of radius $d/2$ touching $\delta$ at $D$	$a^2 = 4bc$

Table 1:  $b = 0$ .

We summarize the results in Table 1. The case 1 described in Figure 8 is supposable without (9). But (9) enable us to get the case by algebraic manipulation. On the other hand, the case 2 described in Figure 9 can not be obtained without (9). In this case  $d = a + b$  does not hold, but still can be considered that  $\alpha$  and  $\beta$  touch. However we cannot attain a reasoned interpretation for this case at the current moment.

**3.2. The case  $a = 0$ .** We now consider the case  $a = 0$ . Substituting  $b = d - a$  in (10), we get

$$f_a = (x^2 + (y - 2d)^2) + 2a(y - 2d) = 0.$$

Hence we get  $x^2 + (y - 2d)^2 = 0$  or  $y = 2d$  in the case  $a = 0$ . Therefore  $\alpha$  coincides with  $D$  or  $t$  in this case. Similarly we have

$$f_b = (x^2 + (y - d)^2 - d^2) + 2ay = 0.$$

Therefore we get  $x^2 + (y - d)^2 = d^2$  or  $y = 0$  in the case  $a = 0$ . Hence  $\beta$  coincides with  $\delta$  or the  $x$ -axis in the case  $a = 0$ . Also we have

$$f_c = 2d(x^2 + (y - 2d)^2) + 2a(x^2 + (y - 2d)^2) + a^2(2d - y) = 0.$$

Therefore we get  $x^2 + (y - 2d)^2 = 0$  or  $y = 2d$  in the case  $a = 0$ . Hence  $\gamma$  coincides with  $D$  or  $t$  in this case.

When  $\alpha$  approaches to  $D$ ,  $\beta$  and  $\gamma$  approach to  $\delta$  and  $D$ , respectively. Hence we consider that  $\beta$  and  $\gamma$  coincide with  $\delta$  and  $D$ , respectively when  $\alpha$  coincides

with  $D$  (see Figure 10). As the remaining case  $\beta$  and  $\gamma$  coincide with the  $x$ -axis and  $t$ , respectively when  $\alpha$  coincides with  $t$  (see Figure 11).

We summarize the results in Table 2. The case 3 described in Figure 10 is supposable without (9). On the other hand, the case 4 described in Figure 11 can not be obtained without (9). However we cannot attain a reasoned interpretation for this case at the current moment.




Figure 10.




Figure 11.

case	$\alpha$	$\beta$	$\gamma$	relation of the radii
3	$D$	$\delta$	$D$	$a^2 = 4bc$
4	$t$	$x$ -axis	$t$	$a^2 = 4bc$

Table 2:  $a = 0$ .

For an extensive introduction of division by zero with Wasan geometry see [14], and its application to Wasan geometry see [4], [8], [9, 10, 11, 12, 13], [15].

#### 4. INCORRECT SANGAKU PROBLEMS

In [16] we have considered two incorrect sangaku problems in [5, p. 69, p. 123], each of which can also be found in [7] and [21], respectively.




Figure 12: The figures in [5], [21].




Figure 13: The figure in [5].

The problems and the answers are almost the same as Problem 1, i.e., they demand to show the relation  $a^2 = 4bc$  for three circles  $\alpha$ ,  $\beta$  and  $\gamma$  of radii  $a$ ,  $b$  and  $c$ , respectively. However the figures are slightly different as shown in Figures 12 and 13. The figure in [21] is also the same as Figure 12. It seems that those problems were correct and essentially the same as Problem 1 in the original context but the figures were incorrectly transcribed in [5] and [21]. While the figure in [7] is the same as Figure 1, therefore the problem is correct.

## REFERENCES

- [1] H. Fukagawa, D. Pedoe, Japanese Temple Geometry Problems, 1989, Charles Babbage Research Centre, Winnipeg Canada.
- [2] Fukushimaken Wasan Kenkyū Hozonkai (福島県和算研究保存会) ed., The Sangaku in Fukushima (福島の算額), Sōju Shuppan (蒼樹出版), 1989.
- [3] A. Hirayama (平山諦), M. Ohno (大野政治), A. Mitsuhashi (三橋愛子) ed., The sangaku in Chiba prefecture (千葉県の算額), Naritasan Shiryōkan (成田山資料館), 1970.
- [4] Y. Kanai, H. Okumura, A three tangent congruent circle problem, Sangaku J. Math., **1** (2017) 16–20.
- [5] Kinki Sūgakushigakukai (近畿数学史学会) ed., The Sangaku in Kinki (近畿の算額), Osaka Kyōiku Tosho (大阪教育図書), 1992.
- [6] M. Kuroda, H. Michiwaki, S. Saitoh, M. Yamane, New meanings of the division by zero and interpretations on  $100/0 = 0$  and on  $0/0 = 0$ , Int. J. Appl. Math., **27**(2) (2014) 191–198.
- [7] H. Kuwabara (桑原秀夫), R. Yoshida (吉田柳二), T. Yamaguchi (山口正) ed., The sangaku in Shiga (滋賀の算額), 1977.
- [8] T. Matsuura, H. Okumura, S. Saitoh, Division by zero calculus and Pompe’s theorem, Sangaku J. Math., **3** (2019) 36–40.
- [9] H. Okumura, Remarks on Archimedean circles of Nagata and Ootoba, Sangaku J. Math., **3** (2019) 119–122.
- [10] H. Okumura, The arbelos in Wasan geometry: Ootoba’s problem and Archimedean circles, Sangaku J. Math., **3** (2019) 91–97.98–104.
- [11] H. Okumura, A characterization of the golden arbelos involving an Archimedean circle, Sangaku J. Math., **3** (2019) 67–71.
- [12] H. Okumura, Wasan geometry with the division by 0, Int. J. Geom., **7**(1) (2018) 17–20.
- [13] H. Okumura, Solution to 2017-1 Problem 4 with division by zero, Sangaku J. Math., **2** (2018) 27–30.
- [14] H. Okumura, S. Saitoh, Wasan geometry and division by zero calculus, Sangaku J. Math., **2** (2018) 57–73.
- [15] H. Okumura and S. Saitoh, Applications of the division by zero calculus to Wasan geometry, Glob. J. Adv. Res. Class. Mod. Geom., **7**(2) (2018) 44–49.
- [16] H. Okumura (奥村博), On a problem in a sangaku in Kinki (近畿の算額の問題について), Wasan (和算), (72) (1992) 13–15.
- [17] S. Saitoh, Division by zero calculus (draft), 2019.
- [18] Saitama prefectural library (埼玉県立図書館) ed., The sangaku in Saitama (埼玉の算額), 1969, Saitama Prefectural Library (埼玉県立図書館).
- [19] Shiraishi (白石長忠) ed., Shamei Sampu (社盟算譜), 1826, Tohoku University Digital Collection.
- [20] Suzuki (鈴木順吉良茂) ed., Sampō Tenshōhō Yōran (算法天生法要覽) volume 1, 1788, Tohoku University Digital Collection.
- [21] I. Yamamoto (山本一郎) ed., The sangaku in Hyōgo (兵庫の算額), 1967.