

$\zeta(z) = X(z) - Y(z)$ A decomposition of the Riemann Zeta function for $Re(z) > 0, z \neq 1$

Pedro Caceres

Dallas, Texas (USA) – March 8th, 2020

Email: pcaceres@comcast.net

Phone: +1 (763) 412-8915

www.DNA-PRIMES.com

Abstract:

In this paper, we define the C-transformation as:

$$[1] \quad C_n\{f\} = \sum_{k=1}^n f(k) - \int f(n) \, dn$$

And the C-values as:

$$[2] \quad C\{f\} = \lim_{n \rightarrow \infty} C_n\{f\}$$

And we obtain a new representation for $\zeta(z)$ in the form $\zeta(z) = X(z) - Y(z)$ applying the C-transformation to the function $f(x) = \frac{1}{x^z}$ for $z \in C, Re(z) \geq 0, z \neq 1$.

Nomenclature and conventions

- a. $\zeta(z) = \sum_{k=1}^{\infty} k^{-z}$ is the Riemann Zeta function
- b. $\alpha = a = Re(z)$ is the real part of a complex number z
- c. $\beta = b = Im(z)$ is the imaginary part of a complex number z

1. C-Transformation of $f(x)$

The C-transformation of an integrable function $f(x)$ is defined by:

$$[3] \quad C_n\{f(x)\} = \sum_{k=1}^n f(k) - \int f(n) \, dn$$

And the C-values is the limit, if it exists, of the C-transformation when $n \rightarrow \infty$:

$$[4] \quad C\{f(x)\} = \lim_{n \rightarrow \infty} C_n\{f(x)\}$$

1.1. C-Transformation of $f(x) = \frac{1}{x}$ for $x \in R$:

$$[5] \quad C_n\left\{\frac{1}{x}\right\} = \sum_{k=1}^n \frac{1}{k} - \int \frac{dn}{n}$$

and

$$[6] \quad C\left\{\frac{1}{x}\right\} = \lim_{n \rightarrow \infty} \left(\sum_{k=1}^n \frac{1}{k} - \ln(n)\right) = \gamma$$

(γ = Euler-Mascheroni constant = 0.5772...)

1.2. C-Transformation of $f(x) = m$, for $m \in R$ constant:

$$[7] \quad C_n\{m\} = \sum_{k=1}^n m - \int m \, dn$$

$$[8] \quad C_n\{m\} = m * n - m * n = 0$$

and the C-values of $f(x) = m$ constant is:

$$[9] \quad C\{m\} = 0$$

1.3. C-Transformation of $f(x) = \sin(x)$ for $x \in R$:

$$[10] \quad C_n\{\sin(x)\} = \sum_{k=1}^n \sin(k) - \int \sin(n) \, dn$$

$$[11] \quad C_n\{\sin(x)\} = 1/2(\sin(n) - \cot\left(\frac{1}{2}\right)\cos(n) + \cot\left(\frac{1}{2}\right) + \cos(n))$$

And the C-values of $f(x) = \sin(x)$ are in the interval:

$$[12] \quad C\{\sin(x)\} \in \left[\frac{1}{2}\left(2 \cot\left(\frac{1}{2}\right) - 3\right), \frac{3}{2}\right]$$

One can also calculate that:

$$[13] \quad C\{\cos(x)\} \in \left[\frac{1}{2}\left(\cot\left(\frac{1}{2}\right) - 4\right), \frac{1}{2}\left(2 - \cot\left(\frac{1}{2}\right)\right)\right]$$

1.4. C-Transformation of $f(x) = e^{-x}$ for $x \in R$:

$$[14] \quad C_n\{e^{-x}\} = \sum_{k=1}^n e^{-k} - \int e^{-n} dn$$

$$[15] \quad C_n\{\sin(x)\} = \sum_{k=1}^n e^{-k} + \frac{e^{-n}}{n}$$

And the C-values of $f(x) = e^{-x}$ are:

$$[16] \quad C\{e^{-x}\} = \frac{1}{e-1}$$

1.5. C-Transformation of $f(x) = x^{-s}$ for $x, s \in R, s > 1$:

$$[17] \quad C_n\left\{\frac{1}{x^s}\right\} = \sum_{k=1}^n \frac{1}{k^s} - \int \frac{dn}{n^s}$$

$$[18] \quad C_n\left\{\frac{1}{x^s}\right\} = \sum_{k=1}^n \frac{1}{k^s} - \frac{n^{1-s}}{1-s}$$

and the C-value of $f(x) = \frac{1}{x^s}$ is the Riemann Zeta function for $s > 1$:

$$[19] \quad C\left\{\frac{1}{x^s}\right\} = \lim_{n \rightarrow \infty} \left(\sum_{k=1}^n \frac{1}{k^s} - \frac{n^{1-s}}{1-s} \right) = \lim_{n \rightarrow \infty} \left(\sum_{k=1}^n \frac{1}{k^s} \right) - \lim_{n \rightarrow \infty} \left(\frac{n^{1-s}}{1-s} \right) = \zeta(s) - 0 = \zeta(s)$$

1.6. C-Transformation of $f(z) = \frac{1}{x^z}$ for $z \in C, Re(z) \geq 0, z \neq 1$

$$[20] \quad C_n\left\{\frac{1}{x^z}\right\} = \sum_{k=1}^n \frac{1}{k^z} - \int \frac{dn}{n^z}$$

We will use Euler's identity:

$$[21] \quad e^x = \cos(x) + i * \sin(x)$$

To calculate [20] for $z = \alpha + \beta i$:

$$[22] \quad k^{-z} = k^{-\alpha} [\cos(\beta * \ln k) - i (\sin(\beta * \ln k))]$$

And:

$$[23] \quad \int dn/n^z = n^{(1-\alpha)} [\cos(\beta * \ln(n) - i \sin(\beta * \ln(n))) * \frac{[(1-\alpha) + i\beta]}{[(1-\alpha)^2 + \beta^2]}$$

One can now express the real and imaginary components of $C_n\{f\}$ as:

$$[24] \quad \text{Re}(C_n\{f\}) = \sum_{k=1}^n k^{-\alpha} (\cos(\beta * \ln(k)) + \frac{1}{[(1-\alpha)^2 + \beta^2]} (n^{(1-\alpha)} [(1-\alpha)*\cos(\beta*\ln(n)) + \beta* \sin(\beta*\ln(n))]))$$

$$[25] \quad \text{Im}(C_n\{f\}) = -\sum_{k=1}^n k^{-\alpha} (\sin(\beta * \ln(k)) + \frac{1}{[(1-\alpha)^2 + \beta^2]} (n^{(1-\alpha)} [\beta*\cos(\beta*\ln(n)) - (1-\alpha)*\sin(\beta*\ln(n))]))$$

One can calculate that, for $\alpha = \text{Re}(z) > 2$, and for any ϵ arbitrarily small, there is a value of $n=N$ such that for $n > N$, $C_N\{f\} - \zeta(z) < \epsilon$, as the following table shows:

α	β	$C_N\{f\}$ for $N=500$	$\zeta(z)$	$ C_N\{f\} - \zeta(z) $
2	0	1.644934068	1.654934067	$< 10^{-8}$
2	1	1.150355702 + 0.437530865 i	1.150355703 + 0.437530866 i	$< 10^{-8}$
3	0	1.202056903	1.202056903	$< 10^{-9}$

Table 1. Values of $C_n\{f(n) = k^{-z}\}$ for $\alpha = \text{Re}(z) > 1$ for $N=500$

The error $C_n\{f\} - \zeta(z)$ grows significantly in the critical strip for $0 \leq \alpha < 1$ as we can see in the following table:

A	β	$C_n\{f\}$	$\zeta(z)$	$ C_n\{f\} - \zeta(z) $
0.0	0	$C_N\{f\}$ for $N=500$	-0.5	0.5
0.2	2	0.399824505 + 0.322650799 i	0.360103 + 0.266246 i	> 0.05
0.7	0	-2.777900606	-2.7783884455	$> 10^{-4}$

Table 2. Values of $C_n\{f(n) = k^{-z}\}$ for $0 \leq \text{Re}(z) < 1$ for $N=500$

To understand better the value of the difference $C_n\left\{\frac{1}{k^z}\right\} - \zeta(z)$, one can plot the difference for $\alpha \in [0,1)$ and $\beta = 0$: (Similar exponential charts occur for all values of $\alpha \in [0,1)$ for any given value of β)




Figure 1 where $a = \alpha = \text{Re}(z)$ and $b = \beta = \text{Im}(z)$

And plot the difference for variable values of $\beta \in [0,1)$ and $\alpha = 0$: (Similar sine charts occur for all values of $\beta \in [0,1)$ for any given value of α)




Figure 2 where $a=\alpha=Re(z)$ and $b=\beta=Im(z)$

These charts lead to the following calculation of the difference $C_n \left\{ \frac{1}{kz} \right\} - \zeta(z)$:

$$[26] \quad \text{Re} \left[C_n \left\{ \frac{1}{kz} \right\} - \zeta(z) \right] = \frac{1}{2} n^{-a} * \cos (\beta * \ln (n)) + O\left(\frac{1}{n}\right)$$

$$[27] \quad \text{Im} \left[C_n \left\{ \frac{1}{kz} \right\} - \zeta(z) \right] = \frac{1}{2} n^{-a} * \sin (\beta * \ln (n)) + O\left(\frac{1}{n}\right)$$

With $O(1/n) \rightarrow 0$ when $n \rightarrow \infty$.

And one can finally write:

$$[28] \quad \begin{aligned} \text{Re}(C_n\{f\}) &= \sum_{k=1}^n k^{-a} (\cos (\beta * \ln (k)) + \\ &+ \frac{1}{[(1-\alpha)^2+\beta^2]} (n^{(1-\alpha)} [(1-\alpha)*\cos(\beta*\ln(n))+\beta*\sin(\beta*\ln(n))])) \\ &+ \frac{1}{2} n^{-a} * \cos(\beta * \ln(n)) \end{aligned}$$

$$[29] \quad \begin{aligned} \text{Im}(C_n\{f\}) &= -\sum_{k=1}^n k^{-a} (\sin (\beta * \ln (k)) + \\ &+ \frac{1}{[(1-\alpha)^2+\beta^2]} (n^{(1-\alpha)} [\beta*\cos(\beta*\ln(n)) - (1-\alpha)*\sin(\beta*\ln(n))])) \\ &+ \frac{1}{2} n^{-a} * \sin(\beta * \ln(n)) \end{aligned}$$

and the C-value of $f(x) = \frac{1}{x^z}$ for $z \in C, Re(z) \geq 0, z \neq 1$ is the Riemann Zeta function $\zeta(z)$.

1.7. A decomposition of $\zeta(z)$ based on the C-transformation of $f(x) = \frac{1}{x^z}$ for $z \in C, Re(z) \geq 0, z \neq 1$

One can rewrite [28] and [29] creating the X(z,n) and Y(z,n) functions:

$$[30] \quad \zeta(z) = \lim_{n \rightarrow \infty} [X(z,n) - Y(z,n)], \text{ where:}$$

$$[31] \quad X(z,n) = (\sum_{k=1}^n k^{-\alpha} (\cos(\beta * \ln(k)) + \frac{1}{2} n^{-\alpha} \cos(\beta \ln(n)) + i * (\sum_{k=1}^n k^{-\alpha} (\sin(\beta * \ln(k)) + \frac{1}{2} n^{-\alpha} \sin(\beta \ln(n))))$$

$$[32] \quad Y(z,n) = n^{(1-\alpha)} \frac{1}{[(1-\alpha)^2 + \beta^2]} [((1-\alpha) * \cos(\beta \ln(n)) + \beta * \sin(\beta \ln(n))) + i (\beta * \cos(\beta \ln(n)) - (1-\alpha) * \sin(\beta \ln(n)))]$$

If:

$$X(z) = \lim_{n \rightarrow \infty} X(z,n) \text{ and}$$

$$Y(z) = \lim_{n \rightarrow \infty} Y(z,n)$$

Then, one can write:

$$\zeta(z) = X(z) - Y(z)$$

The following table shows values for [30]:

z= 0 +j* 0 and n=500
Zeta(z) = -0.5 + i* 0.0 X(z)-Y(z) = -0.5 + i* 0.0 ---> Error = 0.0 + i* 0.0
z= 0.2 +j* 2 and n=500
Zeta(z) = 0.360102590022591 + i* -0.266246199765574 X(z)-Y(z) = 0.360102741838091 + i* -0.266246128959438 ---> Error= -1.5181550 e-7 + i* -7.080613 e-8
z= 0.4 +j* 0 and n=500
Zeta(z) = -1.13479778386698 + i* 0.0 X(z)-Y(z) = -1.1347977871726 + i* 0.0 ---> Error= 3.305619 e-9 + i* 0.0

Table 3. $\zeta(z)$ compared to $X(z) - Y(z)$

The highest error for $\alpha \in [0,1), \beta \in [0,100], n=1000$ is $8x10^{-6}$.

2. Representation of the function $X(z,n)$

The following chart represents $X(z,n)$ for $a=1/2$ and $b \in [1,6]$ and $n=250$

Function $X(z)$




Fig. 3: $X(z,n)$

The following chart represents $X(z,n)$ for $a \in [1,6]$ and $b = 1$ and $n = 250$

Function $X(z)$




Fig. 4: $X(z,n)$

3. Representation of the function $Y(z,n)$

The following chart represents $Y(z,n)$ for $a=1/2$ and $b \in [1,6]$ and $n=250$

Function $Y(z)$




Fig. 5: $Y(z,n)$

The following chart represents $Y(z,n)$ for $a \in [1,6]$ and $b = 1$ and $n = 250$

Function $Y(z)$




Fig. 6: $Y(z,n)$

4. Graphical representations of Square Modulus of $\zeta(z)$, $X(z)$, $Y(z)$, and $X(z) - Y(z)$

Representation of $|\zeta(z)|^2$ and $|X(z)-Y(z)|^2$. These two charts are the same as expected.




Fig. 7 : $|\zeta(z)|^2 = |X(z)-Y(z)|^2$

Representation of $|\zeta(z)|^2$, $|X(z)|^2$, $|Y(z)|^2$, and $|X(z)|^2 - |Y(z)|^2$




Fig. 8

Functions $|\zeta(z)|^2$, $|X(z)|^2$, $|Y(z)|^2$ and $|X(z)|^2 - |Y(z)|^2$
 $\alpha=0.5$




Fig. 9: if $|\zeta(z)|^2 = 0$ then $|X(z)|^2 = |Y(z)|^2$ when $Re(z)=1/2$

One can observe in the following chart that the zeros of $|X(z)|^2 - |Y(z)|^2$ happen on $z=1/2+i\beta_n$ with β_n the imaginary part of the nontrivial zeros of $\zeta(z)$. This shows that for a nontrivial zero of $\zeta(z)$:

$$|X(z)|^2 = |Y(z)|^2$$

$|\zeta(z)|^2$ and $|X(z)|^2 - |Y(z)|^2$




Fig. 10: if $|\zeta(z)|^2 = 0$ then $|X(z)|^2 = |Y(z)|^2$ when $Re(z)=1/2$

5. Conclusion

Using the defined C-transformation, one can write the Riemann Zeta function as the difference of two functions $X(z)$ and $Y(z)$ which will provide a new way of analyzing the zeros of the Zeta function.

$\zeta(z) = X(z) - Y(z)$, where:

$$X(z, n) = \left(\sum_{k=1}^n k^{-\alpha} (\cos(\beta * \ln(k)) + \frac{1}{2} n^{-\alpha} \cos(\beta \ln(n))) + \right. \\ \left. + i * \left(\sum_{k=1}^n k^{-\alpha} (\sin(\beta * \ln(k)) + \frac{1}{2} n^{-\alpha} \sin(\beta \ln(n))) \right) \right)$$

and: $X(z) = \lim_{n \rightarrow \infty} X(z, n)$

$$Y(z, n) = n^{(1-\alpha)} \frac{1}{[(1-\alpha)^2 + \beta^2]} \left[((1-\alpha) * \cos(\beta \ln(n)) + \beta * \sin(\beta \ln(n))) + \right. \\ \left. + i (\beta * \cos(\beta \ln(n)) - (1-\alpha) * \sin(\beta \ln(n))) \right]$$

and: $Y(z) = \lim_{n \rightarrow \infty} Y(z, n)$

The following chart (Fig. 1) shows the equivalence of $\zeta(z) = X(z) - Y(z)$:




Fig. 11: $\zeta(z) = X(z) - Y(z)$

The following chart (Fig. 2) shows that $|\zeta(z)|^2 = |X(z) - Y(z)|^2$:




Fig. 12: $|\zeta(z)|^2 = |X(z) - Y(z)|^2$

REFERENCES

- (1) B. Riemann, "Ueber die Anzahl der Primzahlen unter einer gegebenen Grösse", Monat. der Königl. Preuss. Akad. der Wissen. zu Berlin aus der Jahre 1859 (1860), 671–680; also, Gesammelte math. Werke und wissensch. Nachlass, 2. Aufl. 1892, 145–155.
- (2) Kopp, Konrad. "Theory and Application of Infinite Series." Blackie & Son Limited, London, 1928
- (3) Sondow, Jonathan and Weisstein, Eric W. "Riemann Zeta Function." From MathWorld--A Wolfram Web Resource. <http://mathworld.wolfram.com/RiemannZetaFunction.html>
- (4) WolframAlpha.com used for calculations, accessed March 2020.