

# A Proof of the Erdős-Straus Conjecture

Zhang Tianshu

Emails: chinazhangtianshu@126.com;

xinshijizhang@hotmail.com

Zhanjiang city, Guangdong province, China

## Abstract

In this article, we classify positive integers step by step, and use the formulation to represent a certain class therein until all classes.

First, divide all integers  $\geq 2$  into 8 kinds, and formulate each of 7 kinds therein into a sum of 3 unit fractions.

For the unsolved kind, again divide it into 3 genera, and formulate each of 2 genera therein into a sum of 3 unit fractions.

For the unsolved genus, further divide it into 5 sorts, and formulate each of 3 sorts therein into a sum of 3 unit fractions.

For two unsolved sorts  $\frac{4}{49+120c}$  and  $\frac{4}{121+120c}$  where  $c \geq 0$ , we use two unit fractions plus a proper fraction to replace each of them, then take out

these two unit fractions as  $\frac{1}{X}$  and  $\frac{1}{Y}$ . After that, prove that the remainder

can be identically converted to  $\frac{1}{Z}$ .

**AMS subject classification:** 11D72; 11D45; 11P81

**Keywords:** Erdős-Straus conjecture; Diophantine equation; unit fraction

## 1. Introduction

The Erdős-Straus conjecture relates to Egyptian fractions. In 1948, Paul Erdős conjectured that for any integer  $n \geq 2$ , there are

invariably  $\frac{4}{n} = \frac{1}{X} + \frac{1}{Y} + \frac{1}{Z}$ , where  $x, y$  and  $z$  are positive integers; [1].

Later, Ernst G. Straus further conjectured that  $x, y$  and  $z$  satisfy  $x \neq y, y \neq z$  and  $z \neq x$ , because there are the convertible formulas

$$\frac{1}{2r} + \frac{1}{2r} = \frac{1}{r+1} + \frac{1}{r(r+1)} \quad \text{and} \quad \frac{1}{2r+1} + \frac{1}{2r+1} = \frac{1}{r+1} + \frac{1}{(r+1)(2r+1)} \quad \text{where } r \geq 1; [2].$$

Thus, the Erdős conjecture and the Straus conjecture are equivalent from each other, and they are called the Erdős-Straus conjecture collectively.

As a general rule, the Erdős-Straus conjecture states that for every integer

$n \geq 2$ , there are positive integers  $x, y$  and  $z$ , such that  $\frac{4}{n} = \frac{1}{X} + \frac{1}{Y} + \frac{1}{Z}$ .

Yet it remains a conjecture that has neither is proved nor disproved; [3].

## 2. Divide integers $\geq 2$ into 8 kinds and formulate 7 kinds therein

First, divide integers  $\geq 2$  into 8 kinds, i.e.  $8k+1$  with  $k \geq 1$ , and  $8k+2, 8k+3, 8k+4, 8k+5, 8k+6, 8k+7, 8k+8$ , where  $k \geq 0$ , and arrange them as follows:

$\mathbb{N} \setminus \{1\}$ :  $8k+1, 8k+2, 8k+3, 8k+4, 8k+5, 8k+6, 8k+7, 8k+8$

0,	①,	2,	3,	4,	5,	6,	7,	8,
1,	9,	10,	11,	12,	13,	14,	15,	16,

2, 17, 18, 19, 20, 21, 22, 23, 24,  
 3, 25, 26, 27, 28, 29, 30, 31, 32,  
 ..., ..., ..., ..., ..., ..., ..., ..., ...

Excepting  $n=8k+1$ , formulate each of other 7 kinds into  $\frac{1}{X} + \frac{1}{Y} + \frac{1}{Z}$  :

(1) For  $n=8k+2$ , there are  $\frac{4}{8k+2} = \frac{1}{4k+1} + \frac{1}{4k+2} + \frac{1}{(4k+1)(4k+2)}$  ;

(2) For  $n=8k+3$ , there are  $\frac{4}{8k+3} = \frac{1}{2k+2} + \frac{1}{(2k+1)(2k+2)} + \frac{1}{(2k+1)(8k+3)}$  ;

(3) For  $n=8k+4$ , there are  $\frac{4}{8k+4} = \frac{1}{2k+3} + \frac{1}{(2k+2)(2k+3)} + \frac{1}{(2k+1)(2k+2)}$  ;

(4) For  $n=8k+5$ , there are  $\frac{4}{8k+5} = \frac{1}{2k+2} + \frac{1}{(8k+5)(2k+2)} + \frac{1}{(8k+5)(k+1)}$  ;

(5) For  $n=8k+6$ , there are  $\frac{4}{8k+6} = \frac{1}{4k+3} + \frac{1}{4k+4} + \frac{1}{(4k+3)(4k+4)}$  ;

(6) For  $n=8k+7$ , there are  $\frac{4}{8k+7} = \frac{1}{2k+3} + \frac{1}{(2k+2)(2k+3)} + \frac{1}{(2k+2)(8k+7)}$  ;

(7) For  $n=8k+8$ , there are  $\frac{4}{8k+8} = \frac{1}{2k+4} + \frac{1}{(2k+2)(2k+3)} + \frac{1}{(2k+3)(2k+4)}$  .

By this token,  $n$  as above 7 kinds of integers be suitable to the conjecture.

### 3. Divide the unsolved kind into 3 genera and formulate 2 genera therein

For the unsolved kind  $n=8k+1$  with  $k \geq 1$ , divide it by 3 and get 3 genera:

(1) the remainder is 0 when  $k=1+3t$ ; (2) the remainder is 2 when  $k=2+3t$ ;

(3) the remainder is 1 when  $k=3+3t$ , where  $t \geq 0$ , and *ut infra*.

k: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, ...

8k+1: 9, 17, 25, 33, 41, 49, 57, 65, 73, 81, 89, 97, 105, 113, 121, ...

The remainder: 0, 2, 1, 0, 2, 1, 0, 2, 1, 0, 2, 1, 0, 2, 1, ...

Excepting the genus (3), we formulate other 2 genera as follows:

(8) For  $\frac{8k+1}{3}$  where the remainder is equal to 0, there are

$$\frac{4}{8k+1} = \frac{1}{\frac{8k+1}{3}} + \frac{1}{8k+2} + \frac{1}{(8k+1)(8k+2)}$$

Due to  $k=1+3t$  and  $t \geq 0$ , there are  $\frac{8k+1}{3} = 8t+3$ , so we confirm that  $\frac{8k+1}{3}$

in the preceding equation is an integer.

(9) For  $\frac{8k+1}{3}$  where the remainder is equal to 2, there are

$$\frac{4}{8k+1} = \frac{1}{\frac{8k+2}{3}} + \frac{1}{8k+1} + \frac{1}{(8k+1)(8k+2)}$$

Due to  $k=2+3t$  and  $t \geq 0$ , there are  $\frac{8k+2}{3} = 8t+6$ , so we confirm that  $\frac{8k+2}{3}$

and  $\frac{(8k+1)(8k+2)}{3}$  in the preceding equation are two integers.

#### **4. Divide the unsolved genus into 5 sorts and formulate 3 sorts therein**

For the unsolved genus  $\frac{8k+1}{3}$  where the remainder is equal to 1 when  $k=3+3t$  and  $t \geq 0$ , then there are  $8k+1=25, 49, 73, 97, 121$  etc. So we divide

them into 5 sorts:  $25+120c$ ,  $49+120c$ ,  $73+120c$ ,  $97+120c$  and  $121+120c$  where  $c \geq 0$ , and *ut infra*.

$C \setminus n$ :	$25+120c$ ,	$49+120c$ ,	$73+120c$ ,	$97+120c$ ,	$121+120c$ ,
0,	25,	49,	73,	97,	121,
1,	145,	169,	193,	217,	241,
2,	265,	289,	313,	337,	361,
...	...	...	...	...	...

Excepting  $n=49+120c$  and  $n=121+120c$ , formulate other 3 sorts, they are:

(10) For  $n=25+120c$ , there are 
$$\frac{4}{25+120c} = \frac{1}{25+120c} + \frac{1}{50+240c} + \frac{1}{10+48c};$$

(11) For  $n=73+120c$ , there are

$$\frac{4}{73+120c} = \frac{1}{(73+120c)(10+15c)} + \frac{1}{20+30c} + \frac{1}{(73+120c)(4+6c)};$$

(12) For  $n=97+120c$ , there are

$$\frac{4}{97+120c} = \frac{1}{25+30c} + \frac{1}{(97+120c)(50+60c)} + \frac{1}{(97+120c)(10+12c)}.$$

For each of listed above 12 equations which express  $\frac{4}{n} = \frac{1}{X} + \frac{1}{Y} + \frac{1}{Z}$ , please each reader self to make a check respectively.

### 5. Prove the sort $\frac{4}{49+120c} = \frac{1}{X} + \frac{1}{Y} + \frac{1}{Z}$

For a proof of the sort  $\frac{4}{49+120c}$ , it means that when  $c$  is equal to each of

positive integers plus 0, there always are 
$$\frac{4}{49+120c} = \frac{1}{X} + \frac{1}{Y} + \frac{1}{Z}.$$

After  $c$  is given any value,  $\frac{4}{49+120c}$  can be substituted by each of infinite more a sum of two unit fractions plus a proper fraction, and that these fractions are different from one another, as listed below:

$$\begin{aligned} & \frac{4}{49+120c} \\ &= \frac{1}{13+30c} + \frac{1}{(13+30c)(49+120c)} + \frac{2}{(13+30c)(49+120c)} \\ &= \frac{1}{14+30c} + \frac{1}{(14+30c)(49+120c)} + \frac{6}{(14+30c)(49+120c)} \\ &= \frac{1}{15+30c} + \frac{1}{(15+30c)(49+120c)} + \frac{10}{(15+30c)(49+120c)} \\ & \dots \\ &= \frac{1}{13+\alpha+30c} + \frac{1}{(13+\alpha+30c)(49+120c)} + \frac{4\alpha+2}{(13+\alpha+30c)(49+120c)}, \text{ where } \alpha \geq 0 \\ & \text{and } c \geq 0 \end{aligned}$$

...

As listed above, we can first let  $\frac{1}{13+\alpha+30c} = \frac{1}{X}$  and

$$\frac{1}{(13+\alpha+30c)(49+120c)} = \frac{1}{Y}, \text{ then go to prove } \frac{4\alpha+2}{(13+\alpha+30c)(49+120c)} = \frac{1}{Z}$$

where  $c \geq 0$  and  $\alpha \geq 0$ , *ut infra*.

**Proof** First, let us compare the values of the numerator  $4\alpha+2$  and the denominator  $(13+\alpha+30c)(49+120c)$ .

Since there is  $c \geq 0$ , so  $13+\alpha+30c$  can always be greater than  $4\alpha+2$ . And then, let us just take  $13+\alpha+30c$  in the denominator, while reserve

49+120c for later.

In the fraction  $\frac{4\alpha + 2}{13 + \alpha + 30c}$ , since the numerator  $4\alpha + 2$  is an even number, such that the denominator  $13 + \alpha + 30c$  must be an even number. Only in this case, it can reduce the fraction to become possibly an unit fraction, so  $\alpha$  in the denominator  $13 + \alpha + 30c$  can only be each of positive odd numbers. After  $\alpha$  is assigned to odd numbers 1, 3, 5 and otherwise, and the

numerator and the denominator of the fraction  $\frac{4\alpha + 2}{13 + \alpha + 30c}$  after the values assignment divided by 2, then the fraction  $\frac{4\alpha + 2}{13 + \alpha + 30c}$  is turned into the

fraction  $\frac{3 + 4t}{k + 15c}$  identically, where  $c \geq 0$ ,  $t \geq 0$  and  $k \geq 7$ .

The noteworthy point above is that  $3 + 4t$  and  $k + 15c$  after the values assignment make up a fraction, they are on the same order of taking values of  $t$  and  $k$ , according to the order from small to large, i.e.

$$\frac{3 + 4t}{k + 15c} = \frac{3}{7 + 15c}, \frac{7}{8 + 15c}, \frac{11}{9 + 15c}, \dots$$

Such being the case, letting the numerator and the denominator of the

fraction  $\frac{3 + 4t}{k + 15c}$  divided by  $3 + 4t$ , then we get a temporary indeterminate

unit fraction, and its denominator is  $\frac{k + 15c}{3 + 4t}$ , and its numerator is 1.

Thus, we are necessary to prove that the denominator  $\frac{k + 15c}{3 + 4t}$  is able to

become a positive integer in the case where  $t \geq 0$ ,  $k \geq 7$  and  $c \geq 0$ .

In the fraction  $\frac{k+15c}{3+4t}$ , due to  $k \geq 7$ , the numerator  $k+15c$  after the values assignment are infinite more consecutive positive integers, while the denominator  $3+4t = 3, 7, 11$  and otherwise positive odd numbers.

The key above is that each value of  $3+4t$  after the values assignment can seek its integral multiples within infinite more consecutive positive integers of  $k+15c$ , in the case where  $t \geq 0$ ,  $k \geq 7$  and  $c \geq 0$ .

As is known to all, there is a positive integer that contains the odd factor  $2n+1$  within  $2n+1$  consecutive positive integers, where  $n=1, 2, 3, \dots$

Like that, there is a positive integer that contains the odd factor  $3+4t$  within  $3+4t$  consecutive positive integers of  $k+15c$ , no matter which odd number that  $3+4t$  is equal to, where  $t \geq 0$ ,  $k \geq 7$  and  $c \geq 0$ . It is obvious that a fraction that consists of such a positive integer as the numerator and  $3+4t$  as the denominator is an improper fraction.

Undoubtedly, every such improper fraction that is found in this way, via the reduction, it is surely a positive integer.

That is to say,  $\frac{k+15c}{3+4t}$  as the denominator of the aforesaid temporary indeterminate unit fraction can become a positive integer, and the positive

integer is represented by  $\mu$ . So, the fraction  $\frac{3+4t}{k+15c}$  is exactly  $\frac{1}{\mu}$ .



For the unit fraction  $\frac{1}{\mu}$ , multiply its denominator by  $49+120c$  reserved,

then we get the unit fraction  $\frac{1}{\mu(49+120c)}$ , and let  $\frac{1}{\mu(49+120c)} = \frac{1}{Z}$ .

Since  $\frac{1}{(13+\alpha+30c)(49+120c)} + \frac{4\alpha+2}{(13+\alpha+30c)(49+120c)}$  are equal to

$\frac{4\alpha+3}{(13+\alpha+30c)(49+120c)}$ , so if  $4\alpha+3$  serve as one numerator such that

$\frac{4\alpha+3}{(13+\alpha+30c)(49+120c)}$  is an unit fraction, then we can multiply the

denominator of the unit fraction by 2 to make a sum of two identical unit fractions, then again, convert them into the sum of two each other's -

distinct unit fractions by the formula  $\frac{1}{2r} + \frac{1}{2r} = \frac{1}{r+1} + \frac{1}{r(r+1)}$ .

Thus it can be seen, the fraction  $\frac{4\alpha+3}{(13+\alpha+30c)(49+120c)}$  is able to be expressed into a sum of two each other's -distinct unit fractions in the case where  $c \geq 0$  and  $\alpha \geq 0$ .

Let us synthesize the above results inferred, then get the equation

$$\frac{4}{49+120c} = \frac{1}{13+\alpha+30c} + \frac{1}{(13+\alpha+30c)(49+120c)} + \frac{1}{\mu(49+120c)}, \text{ where } \alpha \geq 0, \mu$$

is an integer and  $\mu = \frac{k+15c}{3+4t}$ ,  $t \geq 0, k \geq 7$  and  $c \geq 0$ .

In other words, we have proved  $\frac{4}{49+120c} = \frac{1}{X} + \frac{1}{Y} + \frac{1}{Z}$ .

**6. Prove the sort**  $\frac{4}{121+120c} = \frac{1}{X} + \frac{1}{Y} + \frac{1}{Z}$

The proof in this section is exactly similar to that in the section 5. Namely,

for a proof of the sort  $\frac{4}{121+120c}$ , it means that when c is equal to each of

positive integers plus 0, there always are  $\frac{4}{121+120c} = \frac{1}{X} + \frac{1}{Y} + \frac{1}{Z}$ .

After c is given any value,  $\frac{4}{121+120c}$  can be substituted by each of infinite more a sum of two unit fractions plus a proper fraction, and that these fractions are different from one another, as listed below.

$$\begin{aligned} & \frac{4}{121+120c} \\ &= \frac{1}{31+30c} + \frac{1}{(31+30c)(121+120c)} + \frac{2}{(31+30c)(121+120c)} \\ &= \frac{1}{32+30c} + \frac{1}{(32+30c)(121+120c)} + \frac{6}{(32+30c)(121+120c)} \\ &= \frac{1}{33+30c} + \frac{1}{(33+30c)(121+120c)} + \frac{10}{(33+30c)(121+120c)} \\ & \dots \\ &= \frac{1}{31+\alpha+30c} + \frac{1}{(31+\alpha+30c)(121+120c)} + \frac{4\alpha+2}{(31+\alpha+30c)(121+120c)}, \text{ where } \alpha \geq 0 \end{aligned}$$

and  $c \geq 0$ .

...

As listed above, we can first let  $\frac{1}{31+\alpha+30c} = \frac{1}{X}$  and

$$\frac{1}{(31+\alpha+30c)(121+120c)} = \frac{1}{Y}, \text{ then go to prove } \frac{4\alpha+2}{(31+\alpha+30c)(121+120c)} = \frac{1}{Z}$$

where  $\alpha \geq 0$  and  $c \geq 0$ , *ut infra*.

**Proof** First, let us compare the values of the numerator  $4\alpha+2$  and the denominator  $(31+\alpha+30c)(121+120c)$ .

Since there is  $c \geq 0$ , so  $31+\alpha+30c$  can always be greater than  $4\alpha+2$ . And then, let us just take  $31+\alpha+30c$  in the denominator, while reserve  $121+120c$  for later.

In the fraction  $\frac{4\alpha+2}{31+\alpha+30c}$ , since the numerator  $4\alpha+2$  is an even number, such that the denominator  $31+\alpha+30c$  must be an even numbers. Only in this case, it can reduce the fraction to become possibly an unit fraction, so  $\alpha$  in the denominator  $31+\alpha+30c$  can only be each of positive odd numbers.

After  $\alpha$  is assigned to odd numbers 1, 3, 5 and otherwise, and the numerator and the denominator of the fraction  $\frac{4\alpha+2}{31+\alpha+30c}$  after the values

assignment divided by 2, then the fraction  $\frac{4\alpha+2}{31+\alpha+30c}$  is turned into the

fraction  $\frac{3+4t}{m+15c}$  identically, where  $c \geq 0$ ,  $t \geq 0$  and  $m \geq 16$ .

The noteworthy point above is that  $3+4t$  and  $m+15c$  after the values assignment make up a fraction, they are on the same order of taking values of  $t$  and  $m$ , according to the order from small to large, i.e.

$$\frac{3+4t}{m+15c} = \frac{3}{16+15c}, \frac{7}{17+15c}, \frac{11}{18+15c}, \dots$$

Such being the case, letting the numerator and the denominator of the

fraction  $\frac{3+4t}{m+15c}$  divided by  $3+4t$ , then we get a temporary indeterminate

unit fraction, and its denominator is  $\frac{m+15c}{3+4t}$ , and its numerator is 1.

Thus, we are necessary to prove that the denominator  $\frac{m+15c}{3+4t}$  is able to become a positive integer in the case where  $t \geq 0$ ,  $m \geq 16$  and  $c \geq 0$ .

In the fraction  $\frac{m+15c}{3+4t}$ , due to  $m \geq 16$ , the numerator  $m+15c$  after the values assignment are infinite more consecutive positive integers, while the denominator  $3+4t=3, 7, 11$  and otherwise positive odd numbers.

The key above is that each value of  $3+4t$  after the values assignment can seek its integral multiples within infinite more consecutive positive integers of  $m+15c$  in the case where  $t \geq 0$ ,  $m \geq 16$  and  $c \geq 0$ .

As is known to all, there is a positive integer that contains the odd factor  $2n+1$  within  $2n+1$  consecutive positive integers, where  $n=1, 2, 3, \dots$

Like that, there is a positive integer that contains the odd factor  $3+4t$  within  $3+4t$  consecutive positive integers of  $m+15c$ , no matter which odd number that  $3+4t$  is equal to, where  $t \geq 0$ ,  $m \geq 16$  and  $c \geq 0$ . It is obvious that a fraction that consists of such a positive integer as the numerator and  $3+4t$  as the denominator is an improper fraction.

Undoubtedly, every such improper fraction that is found in this way, via the reduction, it is surely a positive integer.

That is to say,  $\frac{m+15c}{3+4t}$  as the denominator of the aforesaid temporary indeterminate unit fraction can become a positive integer, and the positive

integer is represented by  $\lambda$ . So, the fraction  $\frac{3+4t}{m+15c}$  is exactly  $\frac{1}{\lambda}$ .

For the unit fraction  $\frac{1}{\lambda}$ , multiply its denominator by  $121+120c$  reserved,

then we get the unit fraction  $\frac{1}{\lambda(121+120c)}$ , and let  $\frac{1}{\lambda(121+120c)} = \frac{1}{Z}$ .

Since  $\frac{1}{(31+\alpha+30c)(121+120c)} + \frac{4\alpha+2}{(31+\alpha+30c)(121+120c)}$  are equal to

$\frac{4\alpha+3}{(31+\alpha+30c)(121+120c)}$ , so if  $4\alpha+3$  serve as one numerator such that

$\frac{4\alpha+3}{(31+\alpha+30c)(121+120c)}$  is an unit fraction, then we can multiply the

denominator of the unit fraction by 2 to make a sum of two identical unit fractions, then again, convert them into the sum of two each other's -

distinct unit fractions by the formula  $\frac{1}{2r} + \frac{1}{2r} = \frac{1}{r+1} + \frac{1}{r(r+1)}$ .

Thus it can be seen, the fraction  $\frac{4\alpha+3}{(31+\alpha+30c)(121+120c)}$  is able to be expressed into a sum of two each other's -distinct unit fractions in the case where  $\alpha \geq 0$  and  $c \geq 0$ .

Let us synthesize the above results inferred, then get the equation

$$\frac{4}{121+120c} = \frac{1}{31+\alpha+30c} + \frac{1}{(31+\alpha+30c)(121+120c)} + \frac{1}{\lambda(121+120c)},$$

where  $\lambda$  is

an integer and  $\lambda = \frac{m+15c}{3+4t}$ ,  $t \geq 0$ ,  $m \geq 16$ , and  $c \geq 0$ .

In other words, we have proved  $\frac{4}{121+120c} = \frac{1}{X} + \frac{1}{Y} + \frac{1}{Z}$ .

The proof was thus brought to a close. As a consequence, the Erdős-Straus conjecture is tenable.

## References

[1] J. W. Sander, *On  $\frac{4}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}$  and Iwaniec's Half Dimensional Sieve*;

Journal of Number Theory, Volume 46, Issue 2, February 1994, Pages 123-136;

<https://www.sciencedirect.com/science/article/pii/S0022314X84710080?via%3Dihub>

[2] Wolfram Math world, the web's most extensive mathematics resource;

<https://mathworld.wolfram.com/Erdos-StrausConjecture.html>

[3] Konstantine Zelator, *An ancient Egyptian problem: the diophantine*

*equation  $\frac{4}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}$ ,  $n > or = 2$* ; arXiv: 0912.2458.