

# An Information Based Theory of Stationary Action

David Trost  
david\_trost@icloud.com

9-September-2023\*

*The quantization of energy proposed by Planck to account for the observed spectrum of black body radiation has associated with it a quantization of entropy. This in turn implies a quantization of observable information, directly implying observational uncertainty on the order of Planck's constant. The effect of that uncertainty is analyzed. In order to adhere strictly to the use of observable quantities, a probability measure is employed based on the distinguishability of statistical samples. This leads directly to the description of probability in terms of the absolute square of a complex amplitude. The Feynman rules may then be applied naturally for indistinguishable events without contradiction to the conventional rules for distinguishable events. This enables the straightforward calculation of the probability that a particle moves from one arbitrary point to another. The Feynman formulation of quantum phenomena and the principle of stationary action results when it is assumed that the classical action represents the measure of distinguishability. Parallel analysis on a Lorentz manifold yields the geodesic principle.*

## Introduction

Early efforts to understand black body radiation within the confines of classical physics focused on the entropy of electromagnetic radiation. In 1884 Boltzmann studied black body radiation in a perfectly reflecting enclosure. Treating the radiation pressure as that of a continuum gas he was able to define its entropy [1]. From that followed a theoretical basis for Stefan's empirically determined dependence of total radiated power on the fourth power of temperature.

Several years earlier Boltzmann had shown a correspondence between classical entropy and the discrete quantity that was later called the "statistical multiplicity"[2] of a molecular gas, though he made no use of this in his black body work. It was not until the mid-twentieth century introduction of information theory [3] that entropy could be understood as information lost due to the statistical treatment of trajectories in lieu of a more complete microscopic model of molecular states [4].

By 1900, Planck had developed a classical model of the entropy of a black body at temperature  $T$  in which electromagnetic dipole resonators operated in equilibrium with the radiant energy in Boltzmann's reflective enclosure. Comparing the latest empirical data to his model, he found it necessary to introduce the constant  $h$  limiting radiation energy to discrete multiples of  $h\nu$  [5]. This quantization of energy has the effect of limiting the entropy to increments of  $h\nu/T$  when expressed in the prevailing units, those of Boltzmann's constant  $k$ .

The appearance of entropy in discrete increments is consistent with the situation in statistical mechanics. In that case discrete values replace the continuum model since entropy is now based on the discrete statistical multiplicity of macro-states. As in the case of the continuum gas model, the entropy in the black body model will be interpreted as a paucity of accurate information, in this case due to some inherent natural limit.

Consider an ideal gas consisting of a single molecule. The Planck entropy implies an inability of the classical model to describe its trajectory in phase space with uncertainty less than  $h$ . It may be that a more accurate description of the trajectory is not possible. Alternately, the trajectory may be fully deterministic but some inherent limit in observational accuracy produces the uncertainty, even with perfectly accurate measuring equipment.

---

\*Initial text posted 25-March-2023

In either case the description of observable quantities in the classical model, like that of a continuum gas is only approximate, with the action of observed natural phenomena differing from the classical description. The result is an observed stochastic component of order  $h$  in the molecular trajectory. In this situation, Planck's constant  $h$  is a more convenient unit of entropy.

Let  $\eta$  be a system dependent parameter, then let

$$I_1 = \eta h \quad (1)$$

represent the information in a hypothetical, more accurate and possibly fully deterministic model that supersedes the inaccurate part of the information in the classical description. Let  $H_1$  represent the entropy of the more accurate model and  $H_0$  the entropy of the classical model. Then [6]

$$H_1 = H_0 - I_1 \quad (2)$$

If the more accurate model is both fully deterministic and completely accurate

$$H_1 = 0 \quad (3)$$

Then

$$H_0 = I_1 = \eta h \quad (4)$$

The analysis that follows explicitly acknowledges statistical uncertainties in observations of physical systems. Following classical practice, we assume no explicit limit on the accuracy of measuring instruments. Also, as in the classical model, the explicit effect of a measurement is not assumed in advance to significantly affect its own result, nor the results of future measurements.

The existence of uncertainty requires that the inherently stochastic nature of the result of observation be incorporated in the analysis. The choice of probability measure can be of profound importance [7]. Following modern practice, careful attention is paid to ensuring our analysis is based strictly on what can be observed. To that end we employ a probability measure based on stochastic outcomes that are equal in statistical distinguishability from one another.

Distinguishing one experimental outcome from another is necessarily a matter of distinguishing between their probability distributions. The distinguishability of probability distributions has been studied by Wootters [8]. It is measured by the quantity *statistical distance* on a probability space.

Consider two  $N$  sided loaded die with different loadings, where the differences in the probabilities of corresponding faces are  $\delta p_1 \cdots \delta p_N$ . The die are said to be distinguishable in  $n$  trials if

$$\frac{\sqrt{n}}{2} \left[ \sum_{i=1}^N \frac{(\delta p_i)^2}{p_i} \right] > 1 \quad (5)$$

Then the statistical distance  $\mathcal{S}$  between these dice on the appropriate probability space is defined by

$$\mathcal{S} = \lim_{n \rightarrow \infty} \frac{1}{\sqrt{n}} \times \left[ \begin{array}{l} \text{the maximum number of intermediate outcomes each of} \\ \text{which is distinguishable (in } n \text{ trials) from its neighbors} \end{array} \right] \quad (6)$$

Along with the stochastic analysis of what can be observed, the question remains: are these stochastic processes consistent with more deterministic, or even fully deterministic underlying natural processes, even if some of the variables necessary to make use of a more deterministic model are inherently hidden due to some natural limitation on their observability.

### Analysis

Let  $x = (x_0, x_1, x_2, x_3)$  represent the ordinary space and time of classical physics where  $x_0$  represents time and  $x_1, x_2$  and  $x_3$  represent three-dimensional Euclidean space. Let us consider a particle that moves from start point  $A$  to end point  $B$  by an unknown trajectory through this space and time. Let  $x(t)$  be an arbitrary trajectory with the same end points, where  $t$  is a time like parameter. We stipulate, in view of uncertainty, that for any value of the parameter  $t$  assigning a definite time and location on  $x(t)$  there may be a nonzero probability that the particle can be observed at any time and location  $x$ . This defines the probability distribution  $p(x, t)$ . The set of all possible probability distributions constitutes a probability space.

Let  $p(x, t)$  be piecewise differentiable with respect to  $t$ . The statistical distance between points on  $x(t)$  in the physical space may then be expressed as the statistical distance  $\mathcal{S}$  between corresponding points in the associated probability space [8].

$$d\mathcal{S}(t) = \frac{1}{2} \left\{ \int_{-\infty}^{\infty} dx_0 \iiint_{-\infty}^{\infty} dx_1 dx_2 dx_3 \frac{1}{p(x, t)} \left[ \frac{dp(x, t)}{dt} \right]^2 \right\}^{1/2} dt \quad (7)$$

This expression may be simplified by the substitution  $\zeta(x, t) = p^{1/2}(x, t)$ . Then

$$d\mathcal{S}(t) = \left\{ \int_{-\infty}^{\infty} dx_0 \iiint_{-\infty}^{\infty} dx_1 dx_2 dx_3 \left[ \frac{d\zeta(x, t)}{dt} \right]^2 \right\}^{1/2} dt \quad (8)$$

This new expression defines an element of length  $d\mathcal{S}$  in an infinite dimensional Euclidean  $\zeta$  space.

Since  $\zeta^2(x, t)$  is a probability, its integral over all of space and time is unity for any value of the parameter  $t$ . Thus  $d\mathcal{S}$  lies on the surface of an infinite dimensional unit hypersphere. Then statistical distance  $\mathcal{S}$  between points on  $x(t)$  is measured by the length of the arc traced on the surface of the hypersphere as the progress of parameter  $t$  traces out the trajectory in space and time between them [8].

Now let us divide the integral on the right-hand-side of (8) in two. Let  $[da(t)/dt]^2$  represent the portion of the integral for which  $x_0 < t$  and  $[db(t)/dt]^2$  the portion for which  $x_0 > t$ .

$$\left[ \frac{da(t)}{dt} \right]^2 = \int_{-\infty}^t dx_0 \iiint_{-\infty}^{\infty} dx_1 dx_2 dx_3 \left[ \frac{d\zeta(x, t)}{dt} \right]^2 \quad (9)$$

$$\left[ \frac{db(t)}{dt} \right]^2 = \int_t^{\infty} dx_0 \iiint_{-\infty}^{\infty} dx_1 dx_2 dx_3 \left[ \frac{d\zeta(x, t)}{dt} \right]^2 \quad (10)$$

Then  $a$  represents events nominally in the past and  $b$ , the future. Let us form the complex quantity  $a + ib = e^{i\theta}$  where  $\theta = \tan^{-1} b/a$ . The unit hypersphere is collapsed into the unit circle on the Argand plane, while the angle  $\theta(t)$  measures statistical distance.

$$d\mathcal{S}(t) = d\theta(t) \quad (11)$$

Now in order to express the probability  $p[x(t)]$  of the test particle being found at a point on the trajectory  $x(t)$  as an explicit function of the statistical distance  $\theta(t)$  we may define the probability amplitude

$$\varphi[x(t)] \equiv \zeta[x(t)] e^{i\theta(t)} \quad (12)$$

Then

$$p[x(t)] = |\varphi[x(t)]|^2 \quad (13)$$

Representation of the probability in terms of a Hilbert space has the effect of including the measure  $\theta$  in the statement of the probability. Sykora has studied the properties of this representation of probability [9]. He notes that evaluating experimental data in support of a theoretical model involves an ad hoc choice of statistical methodologies. This casts doubt on the confirmation. By specifying the measure along with the probability, the Hilbert space representation of probability avoids this ambiguity.

### The Feynman Rules

The identification of a probability with the absolute square of a complex quantity constitutes the Feynman amplitude-probability rule [10, 11].

While the Feynman rules are explicitly for the purpose of quantum mechanical calculation, the conditions assumed here in developing the probability amplitude consist of no more than a random variable with probability described by a function piecewise differentiable with respect to some parameter. This allows the Feynman rules to be treated as a feature of probability theory under these circumstances, when a measure

based on statistical distinguishability is employed. Indeed, the peculiarities of quantum theory, depending upon what can and cannot be measured, may be regarded as depending upon distinguishability.

The conventional Laplace rule for the probability  $p_L$  of an event that may occur by any of  $m$  alternative means is

$$p_L = \sum_{i=1}^m p_i = \sum_{i=1}^m |\varphi_i|^2 \quad (14)$$

where  $p_i$  represents the probability of each of the alternatives. The Laplace rule is empirical in nature, verified by counting occurrences of the various alternatives [12]. We know from a century of experience with quantum phenomena that when the alternatives are *indistinguishable* the probability  $p_F$  is [10].

$$p_F = |\varphi_F|^2 = \left| \sum_{i=1}^m \varphi_i \right|^2 \quad (15)$$

This constitutes the *Feynman amplitude sum rule* [11, 13].

The conventional Laplace rule for the probability  $p_L$  that  $m$  events all occur is

$$p_L = \prod_{i=1}^m p_i = \prod_{i=1}^m |\varphi_i|^2 \quad (16)$$

where the values of  $p_i$  represent the probability of each event. Our experience with quantum phenomena informs us that when the  $m$  events are indistinguishable the probability  $p_F$  is [10].

$$p_F = |\varphi_F|^2 = \left| \prod_{i=1}^m \varphi_i \right|^2 \quad (17)$$

This constitutes the *Feynman amplitude product rule* [11, 13]. While the two formulae yield identical probabilities, the latter establishes the phase  $\theta_F = \sum \theta_i$  of the probability amplitude for the combined event.

Remarkably, Goyal and Knuth have shown that the Feynman rules can coexist free of conflict with conventional probability [11]. Earlier Sykora noted that while probabilities are often described in terms of a single number, a measure is also necessary. Though it is frequently not explicitly stated, the need for this second real number is never-the-less implied [7]. Indeed, the complex representation of probability developed here and exhibited in quantum formulations has been shown to be a necessary result of any probability represented by a pair of real numbers [14]. In this case amplitude  $|\varphi|$  and phase  $\theta$  respectively describe the probability and the distinguishability of alternatives. The complex probability amplitude allows the former to be conveniently formulated in terms of the latter.

Feynman [13,15] citing von Neumann [16] has shown how the introduction of a means of observation produces an arbitrary unknown phase shift in the probability amplitudes of previously unobservable events. Let  $p_F = |\varphi_1 + \varphi_2|^2$  be the probability of an event with two indistinguishable alternative ways of occurring, signified by the two subscripts. Suppose now that a means of observing the alternatives is provided. The presence of the measuring equipment perturbs the phase of the probability amplitudes by arbitrary unknown amounts  $\theta_1$  and  $\theta_2$ . Now  $p_F = |\varphi_1 e^{i\theta_1} + \varphi_2 e^{i\theta_2}|^2$ . The multiple observations required to observe the statistical frequency of these alternatives require that their phases be averaged over all angles. This results in reversion to the Laplace rule  $p_L = |\varphi_1|^2 + |\varphi_2|^2$  as the sample size approaches infinity.

In view of this one may argue that (15) be treated as the more fundamental empirical rule of probability applying to any indistinguishable alternatives lying on a piecewise differentiable space and time continuum, while the conventional Laplace rule (14) becomes derivative of it. Until the possibility of an observation is present, only an amplitude exists without a corresponding probability. Where observation is possible, a frequency may be observed and a probability emerges. The real valued probability is

independent of phase, and the phase of the probability amplitude is lost with the emergence of a probability.

Von Neumann's arbitrary phase shift was determined based on the properties of the Dirac-von Neumann model of quantum phenomena. Given the undeniable long-term success of that model, the von Neumann phase shift has been treated here as an empirical result. The origin of the multiplication and summation rules as well as this phase shift behavior remain questions of interest.

To make proper use of the Feynman rules it becomes necessary to define all cases when events are inherently indistinguishable. Clearly that is the case when no means of measurement is present. In addition, in the more general case measurement results are indistinguishable when the statistical distance between them is less than unity. Kok [17] has shown this to be a consequence of the Kramer-Rao bound.

For uncertainty to be inherent there must be some mechanism, discussed below, whereby any variables describing underlying physical processes are either inherently random, or are inherently hidden from direct observation.

### The Path Integral

Let  $\Delta\theta = \int_{x(t)} [d\theta(t)/dt] dt$ . This is the statistical distance traced by a particle as it traverses the path  $x(t)$ . Let  $m$  be an arbitrary integer and let  $\delta\theta = \Delta\theta/m$  so that  $x[\theta(t)]$  is divided into  $m$  equally distinguishable segments. Then the probability amplitude for the  $j^{\text{th}}$  segment is  $\varphi_j = \mathcal{A}_j^{(m)} e^{i\delta\theta}$  where  $\mathcal{A}_j^{(m)} = [p_j^{(m)}]^{1/2}$  while  $p_j^{(m)}$ , dependent on the value of  $m$ , is the probability for the  $j^{\text{th}}$  segment were a measurement possible.

The probability amplitude  $\varphi_{\mathcal{P}}[x(t)]$  for a test particle following an arbitrary path  $\mathcal{P} = x(t)$ , when the individual points cannot be observed, is the product of the probability amplitudes that the particle is found at each of the  $m$  intervals on  $x[\theta(t)]$ .

$$\varphi_{\mathcal{P}}[x(t)] = \lim_{m \rightarrow \infty} \prod_{j=1}^m [\mathcal{A}_j^{(m)} e^{i\delta\theta}] = \mathcal{A} e^{i\Delta\theta} \quad (18)$$

where  $\mathcal{A}^2 = \left[ \lim_{m \rightarrow \infty} \prod_{j=1}^m \mathcal{A}_j^{(m)} \right]^2$  is the probability that the path  $x(t)$  is followed when observation of the path is possible.

We, know based on many decades of empirical experience verifying the Feynman formulation of quantum mechanics, that the proper expression for  $\varphi_{\mathcal{P}}[x(t)]$  is [10]

$$\varphi_{\mathcal{P}}[x(t)] = \text{Const} e^{iS/\hbar} \quad (19)$$

where  $S$  is the classical action. Then the substitution

$$S/\hbar = \Delta\theta \quad (20)$$

yields the Feynman result. The classical action  $S$  corresponds to the statistical distance  $S$  traced by a particle as it traverses  $x(t)$  while the classical Lagrangian corresponds to the rate of change of statistical distance with time.

The probability that a test particle goes from start point  $A$  to end point  $B$  by any path is then computed according to the Feynman amplitude sum rule with the path integral replacing the summation in (15).

$$p(BA) = \left| \int_A^B \varphi(BA) \mathcal{D}x(t) \right|^2 \quad (21)$$

### Quantum Gravity

The derivation of the Feynman formalism for ordinary quantum mechanics developed here does not rely on the Euclidean nature of the physical space to which those rules are applied. A parallel derivation can be applied to a particle trajectory on a Lorentz manifold provided an uncertainty is postulated in the observed *four-space location* of a particle on the manifold. This leads directly to the geodesic principle of general relativity in lieu of Hamilton's principle. The equivalence of these two principles in the flat space limit [18] establishes a correspondence between the two forms of uncertainty.

Let  $x^* = (x_0^*, x_1^*, x_2^*, x_3^*)$  represent the four-dimensional Lorentzian manifold of general relativity, where  $x_0^*$  represents time and  $x_1^*, x_2^*$  and  $x_3^*$  represent the three spatial dimensions. Substituting  $x^*$  for  $x$  in (7) through (11) we find the equivalent definition of statistical distance. Continuing in the same vein through (18), we find the equivalent expression for the probability amplitude of an arbitrary path in terms of the statistical length of the path  $\Delta\theta$ .

We omit (19) and (20) which identify statistical distance with the classical action, and substitute  $s/s_c$  for  $\Delta\theta$  where  $s = \int_{x^*(t)} [dx^*(t)/dt] dt$  is the length of  $x^*(t)$  and  $s_c$  is a small increment of length on the Lorentzian manifold characterizing uncertainty in spacetime location. Then the probability amplitude  $\varphi_p^*[x^*(t)]$  for a particle to follow an arbitrary path  $x^*(t)$  from  $A$  to  $B$  is

$$\varphi_p^*[x^*(t)] = \mathcal{A}^* e^{is/s_c} \quad (22)$$

As the path integral (21) yields Hamilton's principle, the new path integral

$$p^*(BA) = \left| \int_A^B \varphi_p^*(BA) \mathcal{D}x^*(t) \right|^2 \quad (23)$$

yield the geodesic principle. The dynamic of general relativity is recovered directly while the principle of stationary action follows in the flat space limit [18]. This yields the relativistic equivalent to the Feynman space time formulation of non-relativistic quantum mechanics [15] providing a plausible and equally general model of quantum gravity subject to empirical validation.

The association of uncertainty with the geometric quantity spacetime-location provides some additional clarity. The presence of some yet to be defined source of uncertainty in the geometry of spacetime provides a manifest source of the limitations of classical physics. It remains unclear whether randomness in spacetime location is inherent to spacetime, or is due to some inherent limit on observational accuracy. Both cases are discussed below.

In either case inherent uncertainty in the location of measuring devices makes all location dependent measurements inherently uncertain. It also provides a plausible explanation for the von Neumann phase shift.

### The Black Body Spectrum Revisited

When viewed in the laboratory frame, the uncertainty of location in spacetime must appear as a small apparently random motion. This in turn results in a small zero-point energy. As early as 1913 employing purely classical analysis, Einstein and Stern showed that the assumption of zero-point energy  $h\nu$  in Planck's dipole oscillators led to the Planck spectrum *without the independent assumption of energy quantization* [19].

Milonni has extended that analysis [15] noting each Planck oscillator is in equilibrium with an associated field mode of Planck's cavity. Employing the same zero-point energy, the equipartition theorem requires the oscillator and field each have energy  $h\nu/2$ .

Let us now momentarily assume that deterministic physical laws resembling to the laws of classical physics continue to govern even at the microscopic level, subject to some undefined source of uncertainty in observed spacetime location. Since the apparently, or actually random motion of the electron and the field at the same location are identical, we would expect no coupling between the zero-point motion of the dipole oscillator and the zero-point component of the field under this assumption.

Returning to Milonni, still employing purely classical analysis he demonstrates that when there is no interaction between the random components of the oscillator and the field, black body spectral density is

$$\rho(\nu) = \frac{8\pi h\nu^3/c^3}{e^{h\nu/kT} - 1} + 4\pi h\nu^3/c^3 \quad (24)$$

in agreement with quantum electrodynamic theory. Though short of proof, this hints that even at the unobservable level the concept of spacetime location remains valid and fully deterministic laws resembling the classical ones may prevail. These restrictions further appear to limit the observed randomness to variations in the spacetime metric.

### Discussion

The present analysis leads naturally to the Feynman formulation under the assumption that our ability to know the state of physical phenomena is inherently imperfect. It relies on a supplemented form of conventional probability theory that employs the Feynman rules as the empirically determined rules of probability for indistinguishable states.

A similar point of view can be applied to the Dirac-von Neumann formalism. Consistent with the " $\psi$ -epistemic" view [20], the probability amplitude represents the state of information available about the system. When the amplitude is defined this way, its collapse does not represent a change in the physical system. Instead, it indicates the state of available information about the system has changed as the result of a measurement. A probability has emerged, while simultaneously the phase of a probability amplitude has vanished.

The probability amplitude may be regarded as representing the available information about a conditional probability [11] based on the state of information prior to measurement. The amplitude after measurement represents a new conditional probability based on the new state of information generated by the measurement.

Goyal's analysis has shown [21] that the logic of the Feynman formalism is equivalent to that of Dirac-von Neumann when it is supplemented with a *no-disturbance postulate*. This posits that there exists a class of *trivial measurements* which have no effect on the probability amplitude. Trivial measurements are defined by the property that they yield no new information about the system being measured. This principle is a natural consequence of the epistemic interpretation of the probability amplitude. With no change in information the conditional probabilities of subsequent outcomes are unchanged.

The no-disturbance postulate resides uncomfortably alongside the notion that uncertainty is caused by the process of measurement. The apparent conflict is eliminated with the adoption of the  $\psi$ -epistemic viewpoint coupled with the proposed entropic origin of uncertainty. The fact that the no disturbance principle requires a change in information to change the probability amplitude is a strong indicator of the latter's epistemic nature.

The entropy and information considerations that motivate the present analysis allow for fully deterministic underlying spacetime locations of both particles and waves that are beyond our ability to accurately observe, hence appearing stochastic in nature. Alternatively, they also allow for a spacetime location inherently stochastic in nature, not just appearance. Similarly, nothing in the  $\psi$ -epistemic viewpoint prevents the existence of fully deterministic, nor of inherently stochastic spacetime [20].

### Wave Particle Duality

The empirically derived Laplace rules of probability among distinguishable alternatives have been shown here to lead naturally to the description of particle probabilities in terms of a complex probability amplitude with wavelike characteristics. With the adoption of the more general Feynman rules, also empirically derived, these probability amplitudes add as do the amplitudes of physical wave phenomena when observation of intermediate events is not possible. This imparts wavelike properties to classical particles resulting strictly from the novel rules of probability.

Planck's derivation of the black body spectrum was based almost entirely on a combination of classical mechanics and electromagnetic theory. It deviated only with an unexplained quantization of electromagnetic radiation. Planck demonstrated that the observed black body spectrum was consistent with the effect of energy quantization on the entropy of radiation.

Particle like behavior of classical electromagnetic waves with entropy empirically imposed by the observed black body spectrum were famously explored as early as 1905 [22, 23]. Einstein's classical analysis of the empirically determined high frequency portion of the black body spectrum found its entropy matched that of an ideal molecular gas with particle energy concentrated in a narrow band around the value  $h\nu$ . Particle like behavior in classical wave phenomena thus explained the photoelectric effect twenty years before the advent of modern quantum theory.

With the benefit of modern information theoretic insights not available in 1905, the present analysis allows us to associate the entropy of the black body spectrum with our inability to account for the apparently, or actually random spacetime locations of both resonator and field. As a result, information about the energy of the radiation field as well as the corresponding entropy at the location of the particles are quantized, rather than the field energy itself. The quantization of field entropy due to the resulting uncertainty mimics that of a particulate gas. This accounts for the appearance of photons.

While the present analysis takes no exception to the notion that light consists of photons in all observable phenomena, it also allows for the existence of unobservable fully deterministic underlying behavior in which electromagnetic effects are not quantized, as in the Einstein-Stern-Milonni black body analysis. Though this is an isolated result, it suggests that deterministic physical processes may continue to operate in the quantum regime even though the location of events in spacetime may suffer inherent limitations in observability. Alternately, there may be a random component in spacetime itself. Further investigation is called for before generalizations be made with confidence, as was the case with Planck's isolated black body result. Let us examine both cases.

#### Fully Deterministic Spacetime

The general relativistic model contains within itself a mechanism whereby an observer with only local information will observe a small zero-point energy. Correspondence between Newtonian mechanics and general relativity occurs when energies of objects under observation are suitably small, and there are no variations in the spacetime metric due to events outside the range of observation [24]. The latter of these conditions precludes from consideration a background level of broadband gravitational radiation.

If such background exists then, it can be expected to impose on the classical picture a small apparently random source of zero-point energy. Even in the full general relativistic model, background gravitational radiation that appears stochastic to an observer with only local knowledge must add an apparently random component to the predictable trajectories of ponderable masses.

The full nature of such a stochastic background of gravitational radiation is an open question [25]. Neither a spectrum nor a characteristic time for the gravitational background is known. That this is the source of uncertainty is of course speculation. If this is the case, the general relativistic model is the not-fully-predictable but fully deterministic model. The lack of predictability stems from our inability to know the background gravitational radiation in anything but stochastic terms.

In the fully deterministic spacetime model, the preexisting laws of general relativity, including classical mechanics, are assumed to be in place while quantum phenomena are caused by an additional small stochastic component in the knowledge of the observer. This model is characterized by fully deterministic location in fully deterministic spacetime with fully deterministic laws of physics, impaired by an inherent limit on observability of deterministic spacetime locations.

Aside from the gravitational background, it is also conceivable that some unknown deterministic model more accurate than general relativity prevails. In this case the difference between the two models would appear as stochastic behavior providing an alternate or additional source of uncertainty.

Regardless of the origin of the spacetime perturbations contemplated as a source of uncertainty, it must be borne in mind that the spacetime metric is a logical construct that cannot be directly measured, but only inferred based on the behavior of ponderable masses and the belief that the Einstein equation holds sway. This is doubly true of the background gravitational radiation under discussion in the present theory. For this component of the metric, the observer may only connect the generating motions of the ponderable masses to the resulting disturbances in a statistical sense. Thus, unlike special relativity, neither general relativity nor the present theory is formulated strictly in terms of observable quantities.

While we cannot be certain, it seems likely Einstein had this aspect of his general theory in mind when, late in life, he made this comment on Ernst Mach's considerable contribution to his thinking [26]:

*I see Mach's greatness in his incorruptible skepticism and independence; in my younger years, however, Mach's epistemological position also influenced me very greatly, a position which today appears to me to be essentially untenable. For he did not place in the correct light the essentially constructive and speculative nature of thought and more especially scientific thought; in consequence of which he condemned theory on precisely those points where its constructive-speculative character unconcealably comes to light, as for example in the kinetic atomic theory.*

### Spacetime With a Random Component

In the alternative to the deterministic case, the preexisting laws of mechanics are not assumed. Let us call this the bootstrap model in recognition of how these laws may come into being. In this model uncertainty is assumed due to fully random variations in the spacetime metric not tied to any underlying deterministic process. This possibility introduces an intriguing problem in our understanding of the laws of physics.

The macroscopically deterministic laws of mechanics are generally described by differential equations. It is in the nature of these equations that macroscopic behavior follows directly from behavior at the microscopic level, yet we are assuming randomness at the microscopic level. How then could it come to be that differential equations provide a near perfect description of nature at the macroscopic level?

The analysis presented here is entirely classical in nature up to and including the Feynman amplitude probability rule. The appearance of the product and sum rules is unexplained. In the face of randomness at the microscopic level, these rules provide a mechanism for the laws of mechanics to emerge from randomness. The path integral assures that the observability of particles that just happen to follow paths of near stationary length in spacetime will be coherently amplified, while the observability of those that do not will be coherently suppressed. Thus, both the principle of stationary action, and the geodesic principle are bootstrapped into existence on the strength of the Feynman rules.

Despite randomness at the microscopic level, a system at least macroscopically described by deterministic differential equations may be observed. In this way the general relativistic model, indicative of what can be observed in a universe highly random at the microscopic level, may arise out of that randomness. In this model then, the Feynman multiplication and addition rules are elevated to fundamental laws of nature, giving rise to the laws of mechanics.

A central feature of the present discussion is the concept of inherent uncertainty. It posits that there exists a class of observational results that are inherently unpredictable by any theory. This holds true even though there may be deterministic hidden variables describing some or all the underlying phenomena. In either case observational results appear random. In the fully deterministic proposal, the randomness is only apparent, resulting from the observer's inability to distinguish deterministic undulation of the spacetime metric from motion with respect to the metric. In the alternative the spacetime metric has a truly random component. As a result of these alternatives, the precise definition of inherent uncertainty remains for now uncertain itself.

---

[1] P. W. Milonni, *The Quantum Vacuum an Introduction to Quantum Electrodynamics*, Academic Press (1994) p. 2.

[2] G. H. Wannier, *Statistical Physics*, Dover (1966) p. 85.

- 
- [3] C. E. Shannon and W. Weaver, *The Mathematical Theory of Communication*, Univ. of Illinois Press (1949)
- [4] A. C. Melissinos, F. Lobkowicz, *Physics for Scientists and Engineers*, Saunders (1975) v.1, p.509
- [5] M. J. Klein, Thermodynamics and Quanta in Planck's Work, *Physics Today*, (Nov. 1966) p. 23
- [6] L. Brillouin, The Negentropy Principle of Information, *J. Appl. Phys.* **24**, 1152 (1953)
- [7] S. Sykora, Quantum Theory and the Bayesian Inference Problems, *J. Stat. Phys.*, **11**, p. 17 (1974)
- [8] W. K. Wootters, Statistical Distance and Hilbert Space, *Phys. Rev. D*, **23**, 357 (1981).
- [9] S. Sykora, Quantum Theory and the Bayesian Inference Problems, *J. Stat. Phys.*, **11**, p. 17 (1974)
- [10] R. P. Feynman, A. R. Hibbs, *Quantum Mechanics and Path Integrals*, McGraw Hill (1965), pp. 5-6.
- [11] P. Goyal, K. H. Knuth, Quantum Theory and Probability Theory: Their Relationship and Origin in Symmetry, *Symmetry*, **3**, pp. 171-206 (2011)
- [12] R. B. Ash, *Basic Probability Theory*, Wiley (1970) pp. 13-14
- [13] R. P. Feynman, The concept of Probability in Quantum Mechanics, *Second Berkeley Symposium on Mathematical Statistics and Probability*, U. Cal. Press (1951)
- [14] P. Goyal, Origin of Complex Amplitudes and Feynman's Rules, *Phys. Rev. A*, **81**, (2010)
- [15] R. P. Feynman, Space-Time Approach to Non-Relativistic Quantum Mechanics, *Review of Modern Physics*, **20**, 2, pp. 367-387 (1948)
- [16] J. von Neumann, *Mathematical Foundations of Quantum Mechanics*, Princeton Univ. Press (1955), translation of *Mathematische Grundlegeren der Quantenmechanik*, Julius Springer (1932)
- [17] P. Kok, Tutorial: Statistical Distance and Fisher information  
[https://www.pieter-kok.staff.shef.ac.uk/docs/geometrical\\_Cramer-Rao.pdf](https://www.pieter-kok.staff.shef.ac.uk/docs/geometrical_Cramer-Rao.pdf)
- [18] A. Einstein, Hamilton's Principle and the General Theory of Relativity, *The Principle of Relativity*, Dover, p. 167, reprint of Methuen (1923)
- [19] P. W. Milonni, *The Quantum Vacuum an Introduction to Quantum Electrodynamics*, pp. 14-17.
- [20] N. Harrigan, R. W. Spekkens, Einstein, Incompleteness, and the Epistemic View of Quantum States, *Foundations of Physics*, **40**(2), pp. 125-157, (2010)
- [21] P. Goyal, Derivation of Quantum Theory from Feynman's Rules, *Phys. Rev. A*, **89**(3), p.032120 (2014)
- [22] A. Einstein, On a Heuristic Point of View Concerning the Production and Transformation of Light, translated and reprinted in: J. Stachel, *Einstein's Miraculous Year*, Princeton pp. 177-198 (1998)
- [23] T. S. Kuhn, *Black-Body Theory and the Quantum Discontinuity 1894-1912*, Univ. of Chicago Press, pp. 180-183, (1978)
- [24] A. Einstein, The Foundation of the General Theory of relativity, *The Principle of Relativity*, Dover, p. 157, reprint of Methuen (1923)
- [25] B. Allen, The Stochastic Gravity-Wave Background: Sources and Detection, *Relativistic Gravitation and Gravitational Radiation Proceedings*, Cambridge Univ. Press, conference C95-09-26, (1997)
- [26] A. Einstein, Autobiographical Notes, in *Albert Einstein Philosopher Scientist*, ed. P. A. Schlipp, MJF Books, p. 21, (1949)