

An introduction to the Smarandache Double factorial function

Felice Russo
Via A. Infante 7
 67051 Avezzano (Aq) Italy

In [1], [2] and [3] the Smarandache Double factorial function is defined as:

Sdf(n) is the smallest number such that Sdf(n)!! is divisible by n, where the double factorial is given by [4]:

$$m!! = 1 \times 3 \times 5 \times \dots \times m, \text{ if } m \text{ is odd};$$

$$m!! = 2 \times 4 \times 6 \times \dots \times m, \text{ if } m \text{ is even}.$$

In this paper we will study this function and several examples, theorems, conjectures and problems will be presented. The behaviour of this function is similar to the other Smarandache functions introduced in the chapter I.

In the table below the first 100 values of function Sdf(n) are given:

n	Sdf(n)	n	Sdf(n)	n	Sdf(n)	n	Sdf(n)	n	Sdf(n)
1	1	21	7	41	41	61	61	81	15
2	2	22	22	42	14	62	62	82	82
3	3	23	23	43	43	63	9	83	83
4	4	24	6	44	22	64	8	84	14
5	5	25	15	45	9	65	13	85	17
6	6	26	26	46	46	66	22	86	86
7	7	27	9	47	47	67	67	87	29
8	4	28	14	48	6	68	34	88	22
9	9	29	29	49	21	69	23	89	89
10	10	30	10	50	20	70	14	90	12
11	11	31	31	51	17	71	71	91	13
12	6	32	8	52	26	72	12	92	46
13	13	33	11	53	53	73	73	93	31
14	14	34	34	54	18	74	74	94	94
15	5	35	7	55	11	75	15	95	19
16	6	36	12	56	14	76	38	96	8
17	17	37	37	57	19	77	11	97	97
18	12	38	38	58	58	78	26	98	28
19	19	39	13	59	59	79	79	99	11
20	10	40	10	60	10	80	10	100	20

According to the experimental data the following two conjectures can be formulated:

Conjecture 4.1 The series $\sum_{n=1}^{\infty} Sdf(n)$ is asymptotically equal to $a \cdot n^b$ where a and b are close to 0.8834.. and 1.759.. respectively.

Conjecture 4.2 The series $\sum_{n=1}^{\infty} \frac{1}{Sdf(n)}$ is asymptotically equal to $a \cdot n^b$ where a and b are close to 0.9411.. and 0.49.. respectively.

Let's start now with the proof of some theorems.

Theorem 4.1. $Sdf(p)=p$ where p is any prime number.

Proof. For $p=2$, of course $Sdf(2)=2$. For p odd instead observes that only for $m=p$ the factorial of first m odd integers is a multiple of p , that is $1 \cdot 3 \cdot 5 \cdot 7 \dots p = (p-2)!! \cdot p$.

Theorem 4.2. For any squarefree even number n ,

$$Sdf(n) = 2 \cdot \max\{p_1, p_2, p_3, \dots, p_k\}$$
where $p_1, p_2, p_3, \dots, p_k$ are the prime factors of n .

Proof. Without loss of generality let's suppose that $n = p_1 \cdot p_2 \cdot p_3$ where $p_3 > p_2 > p_1$ and $p_1 = 2$. Given that the factorial of even integers must be a multiple of n of course the smallest integer m such that $2 \cdot 4 \cdot 6 \dots \cdot m$ is divisible by n is $2 \cdot p_3$. Infact for $m = 2 \cdot p_3$ we have :

$$2 \cdot 4 \cdot 6 \dots 2 \cdot p_2 \dots 2 \cdot p_3 = (2 \cdot p_2 \cdot p_3) \cdot (4 \cdot 6 \dots 2k \cdot 2) = k \cdot (2 \cdot p_2 \cdot p_3) \text{ where } k \in \mathbb{N}$$

Theorem 4.3. For any squarefree composite odd number n ,
 $Sdf(n) = \max\{p_1, p_2, \dots, p_k\}$ where p_1, p_2, \dots, p_k are the prime factors of n .

Proof. Without loss of generality let suppose that $n = p_1 \cdot p_2$ where p_1 and p_2 are two distinct primes and $p_2 > p_1$. Of course the factorial of odd integers up to p_2 is a multiple of n because being $p_1 < p_2$ the factorial will contain the product $p_1 \cdot p_2$ and therefore $n \mid 1 \cdot 3 \cdot 5 \cdot \dots \cdot p_1 \cdot p_2$.

Theorem 4.4. $\sum_{n=1}^{\infty} \frac{1}{Sdf(n)}$ diverges.

Proof. This theorem is a direct consequence of the divergence of sum $\sum_p \frac{1}{p}$ where p is any prime number.

In fact $\sum_{k=1}^{\infty} \frac{1}{Sdf(k)} > \sum_{p=2}^{\infty} \frac{1}{p}$ according to the theorem 4.1 and this proves the theorem.

Theorem 4.5 The $Sdf(n)$ function is not additive that is $Sdf(n+m) \neq Sdf(n) + Sdf(m)$ for $(n,m) = 1$.

Proof. In fact for example $Sdf(2+15) \neq Sdf(2) + Sdf(15)$.

Theorem 4.6 The $Sdf(n)$ function is not multiplicative, that is $Sdf(n \cdot m) \neq Sdf(n) \cdot Sdf(m)$ for $(n,m) = 1$.

Proof. In fact for example $Sdf(3 \cdot 4) \neq Sdf(3) \cdot Sdf(4)$.

Theorem 4.7 $Sdf(n) \leq n$

Proof. If n is a squarefree number then based on theorems 4.1, 4.2 and 4.3 $Sdf(n) \leq n$. Let's now consider the case when n is not a squarefree number. Of course the maximum value of the $Sdf(n)$ function cannot be larger than n because when we arrive in the factorial to n for sure it is a multiple of n .

Theorem 4.8 $\sum_{n=1}^{\infty} \frac{Sdf(n)}{n}$ diverges.

Proof. In fact $\sum_{k=1}^{\infty} \frac{Sdf(k)}{k} > \sum_{p=2}^{\infty} \frac{Sdf(p)}{p}$ where p is any prime number and of course $\sum_p \frac{Sdf(p)}{p}$ diverges because the number of primes is infinite [5] and $Sdf(p)=p$.

Theorem 4.9 $Sdf(n) \geq 1$ for $n \geq 1$

Proof. This theorem is a direct consequence of the $Sdf(n)$ function definition. In fact for $n=1$, the smallest m such that 1 divide $Sdf(1)$ is trivially 1. For $n \neq 1$, m must be greater than 1 because the factorial of 1 cannot be a multiple of n .

Theorem 4.10 $0 < \frac{Sdf(n)}{n} \leq 1$ for $n \geq 1$

Proof. The theorem is a direct consequence of theorem 4.7 and 4.9.

Theorem 4.11 $Sdf(p_k\#) = 2 \cdot p_k$ where $p_k\#$ is the product of first k primes (primorial) [4].

Proof. The theorem is a direct consequence of theorem 4.2.

Theorem 4.12 The equation $\frac{Sdf(n)}{n} = 1$ has an infinite number of solutions.

Proof. The theorem is a direct consequence of theorem 4.1 and the well-known fact that there is an infinite number of prime numbers [5].

Theorem 4.13 The even (odd respectively) numbers are invariant under the application of Sdf function, namely $Sdf(\text{even})=\text{even}$ and $Sdf(\text{odd})=\text{odd}$

Proof. Of course this theorem is a direct consequence of the Sdf(n) function definition.

Theorem 4.14 The diophantine equation $Sdf(n) = Sdf(n+1)$ doesn't admit solutions.

Proof. In fact according to the previous theorem if n is even (odd respectively) then Sdf(n) also is even (odd respectively). Therefore the equation $Sdf(n)=Sdf(n+1)$ can not be satisfied because Sdf(n) that is even should be equal to Sdf(n+1) that instead is odd.

Conjecture 4.3 The function $\frac{Sdf(n)}{n}$ is not distributed uniformly in the interval]0,1].

Conjecture 4.4 For any arbitrary real number $\varepsilon > 0$, there is some number $n \geq 1$ such that $\frac{Sdf(n)}{n} < \varepsilon$

Let's now start with some problems related to the Sdf(n) function.

Problem 1. Use the notation $FSdf(n)=m$ to denote, as already done for the $Zt(n)$ and $Zw(n)$ functions, that m is the number of different integers k such that $Zw(k)=n$.

Example $FSdf(1)=1$ since $Sdf(1)=1$ and there are no other numbers n such that $Sdf(n)=1$

Study the function $Fsd(n)$.

Evaluate $\lim_{m \rightarrow \infty} \frac{\sum_{k=1}^m \frac{FSdf(k)}{k}}{m}$

Problem 2. *Is the difference $|Sdf(n+1)-Sdf(n)|$ bounded or unbounded?*

Problem 3. *Find the solutions of the equations: $\frac{Sdf(n+1)}{Sdf(n)} = k = \frac{Sdf(n)}{Sdf(n+1)}$ where k is any positive integer and $n > 1$ for the first equation.*

Conjecture 4.5 *The previous equations don't admits solutions.*

Problem 4. *Analyze the iteration of $Sdf(n)$ for all values of n . For iteration we intend the repeated application of $Sdf(n)$. For example the k -th iteration of $Sdf(n)$ is:*

$$Sdf^k(n) = Sdf(Sdf(\dots(Sdf(n))\dots)) \quad \text{where } Sdf \text{ is repeated } k \text{ times.}$$

For all values of n , will each iteration of $Sdf(n)$ produces always a fixed point or a cycle?

Problem 5. *Find the smallest k such that between $Sdf(n)$ and $Sdf(k+n)$, for $n > 1$, there is at least a prime.*

Problem 6. *Is the number $0.1232567491011\dots$ where the sequence of digits is $Sdf(n)$ for $n \geq 1$ an irrational or transcendental number? (We call this number the Pseudo-Smarandache-Double Factorial constant).*

Problem 7. *Is the Smarandache Euler-Mascheroni sum (see chapter II for definition) convergent for $Sdf(n)$ numbers? If yes evaluate the convergence value.*

Problem 8. Evaluate $\sum_{k=1}^{\infty} (-1)^k \cdot Sdf(k)^{-1}$

Problem 9. Evaluate $\prod_{n=1}^{\infty} \frac{1}{Sdf(n)}$

Problem 10. Evaluate $\lim_{k \rightarrow \infty} \frac{Sdf(k)}{\theta(k)}$ where $\theta(k) = \sum_{n \leq k} \ln(Sdf(n))$

Problem 11. Are there m, n, k non-null positive integers for which $Sdf(n \cdot m) = m^k \cdot Sdf(n)$?

Problem 12. Are there integers $k > 1$ and $n > 1$ such that $(Sdf(n))^k = k \cdot Sdf(n \cdot k)$?

Problem 13. Solve the problems from 1 up to 6 already formulated for the $Zw(n)$ function also for the $Sdf(n)$ function.

Problem 14. Find all the solution of the equation $Sdf(n) \neq Sdf(n!)$

Problem 15. Find all the solutions of the equation $Sdf(n^k) = k \cdot Sdf(n)$ for $k > 1$ and $n > 1$.

Problem 16. Find all the solutions of the equation $Sdf(n^k) = n \cdot Sdf(k)$ for $k > 1$.

Problem 17. Find all the solutions of the equation $Sdf(n^k) = n^m \cdot Sdf(m)$ where $k > 1$ and $n, m > 0$.

Problem 18. For the first values of the $Sdf(n)$ function the following inequality is true:

$$\frac{n}{Sdf(n)} \leq \frac{1}{8} \cdot n + 2 \quad \text{for } 1 \leq n \leq 1000$$

Is this still true for $n > 1000$?

Problem 19. For the first values of the $Sdf(n)$ function the following inequality is true:

$$\frac{Sdf(n)}{n} \leq \frac{1}{n^{0.73}} \quad \text{for } 1 \leq n \leq 1000$$

Is this still true for all values of $n > 1000$?

Problem 20. For the first values of the $Sdf(n)$ function the following inequality hold:

$$\frac{1}{n} + \frac{1}{Sdf(n)} < n^{-\frac{1}{4}} \quad \text{for } 2 < n \leq 1000$$

Is this still true for $n > 1000$?

Problem 21. For the first values of the $Sdf(n)$ function the following inequality holds:

$$\frac{1}{n \cdot Sdf(n)} < n^{-\frac{5}{4}} \quad \text{for } 1 \leq n \leq 1000$$

Is this inequality still true for $n > 1000$?

Problem 22. Study the convergence of the Smarandache Double factorial harmonic series:

$$\sum_{n=1}^{\infty} \frac{1}{Sdf^a(n)} \quad \text{where } a > 0 \text{ and } a \in \mathbb{R}$$

Problem 23. Study the convergence of the series:

$$\sum_{n=1}^{\infty} \frac{x_{n+1} - x_n}{Sdf(x_n)}$$

where x_n is any increasing sequence such that $\lim_{n \rightarrow \infty} x_n = \infty$

Problem 24. Evaluate

$$\lim_{n \rightarrow \infty} \frac{\sum_{k=2}^n \frac{\ln(Sdf(k))}{\ln(k)}}{n}$$

Is this limit convergent to some known mathematical constant?

Problem 25. Solve the functional equation:

$$Sdf(n)^r + Sdf(n)^{r-1} + \Lambda \wedge Sdf(n) = n$$

where r is an integer ≥ 2 .

What about the functional equation:

$$Sdf(n)^r + Sdf(n)^{r-1} + \Lambda \wedge Sdf(n) = k \cdot n$$

where r and k are two integers ≥ 2 .

Problem 26. Is there any relationship between $Sdf\left(\prod_{k=1}^m m_k\right)$ and $\sum_{k=1}^m Sdf(m_k)$?

References.

- [1] F. Smarandache, *"Only Problems, not Solutions!"*, Xiquan, Publishing House, Phoenix-Chicago, 1990, 1991, 1993;
- [2] F. Smarandache, *"The Florentin Smarandache papers"* special collection, Arizona State University, Hayden Library, Tempe, Box 871006, AZ 85287-1006, USA;
- [3] C. Dumitrescu and V. Seleacu, *Some notions and questions in Number Theory*, Erhus Univ. Press, Glendale 1994
- [4] E. Weisstein, *CRC Concise Encyclopedia of Mathematics*, CRC Press, 1999
- [5] P. Ribenboim, *The book of prime numbers records*, Second edition, New York, Springer-Verlag, 1989