

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

# Maximizing deviation method for neutrosophic multiple attribute decision making with incomplete weight information

Rıdvan Şahin<sup>a</sup>, Peide Liu<sup>b</sup>

<sup>a</sup>Faculty of Education, Bayburt University, Bayburt, 69000, Turkey

<sup>b</sup>School of Management Science and Engineering, Shandong University of Finance and Economics, Jinan 250014, Shandong, China

Corresponding authors: [mat.ridone@gmail.com](mailto:mat.ridone@gmail.com) (Şahin R), and [peide.liu@gmail.com](mailto:peide.liu@gmail.com) (Liu P)

---

## Keywords

Neutrosophic sets  
Single valued neutrosophic sets  
Interval neutrosophic sets  
Aggregation operators  
Neutrosophic multiple attribute decision making  
Maximizing deviation method

## Abstract

This paper develops a method for solving the multiple attribute decision making problems with the single valued neutrosophic information or interval neutrosophic information. We first propose two discrimination functions referred to as score function and accuracy function for ranking the neutrosophic numbers. An optimization model to determine the attribute weights that are partly known is established based on the maximizing deviation method. For the special situations where the information about attribute weights is completely unknown, we propose another optimization model. A practical and useful formula which can be used to determine the attribute weights is obtained by solving a proposed non-linear optimization problem. To aggregate the neutrosophic information corresponding to each alternative, we utilize the neutrosophic weighted averaging (NWA) operators which are the single valued neutrosophic weighted averaging (SVNWA) operator and the interval neutrosophic weighted averaging (INWA) operator. Thus we can determine the order of alternatives and choose the most desirable one(s) based on the score function and accuracy function. Finally, some illustrative examples are presented to verify the proposed approach and to present its effectiveness and practicality.

---

## 1. Introduction

Zadeh introduced the degree of membership/truth ( $t$ ) in 1965 and proposed the concept of fuzzy set. Atanassov introduced the degree of nonmembership/falsehood ( $f$ ) in 1986 and defined the intuitionistic fuzzy set. (To date, the intuitionistic fuzzy sets have been widely applied in solving MCDM problems (Peng et al. 2015b; Wang et al. 2013a, 2013b, 2014)). Using the degree of indeterminacy/neutrality ( $i$ ) as independent component in 1995, Smarandache initiated the neutrosophic set theory. He has coined the words “neutrosophy” and neutrosophic. In 2013, he redefined the neutrosophic set to  $n$  components:  $t_1, t_2, \dots; i_1, i_2, \dots; f_1, f_2, \dots$ .

But, a neutrosophic set will be difficult to apply in real scientific and engineering fields. Therefore, Wang et al. (2005, 2010) proposed the concepts of a single valued neutrosophic set (SVNS) and an interval neutrosophic set (INS) which are an instance of a neutrosophic set, and provided set-theoretic operators and various properties of SVNSs and INSs. Recently, the theory of neutrosophic set has received more and more attentions (Liu et al. 2014; Liu and Wang 2014; Liu and Shi 2015; Peng et al. 2014, 2015a, 2015c; Sahin and Kucuk 2014; Ye 2013; Ye 2014a, 2014b, 2014c, 2014d, 2014e; Zhang et al. 2014, 2015; Broumi et al. 2015; Broumi and Smarandache 2014, 2015; Tian 2015). Zhang et al. (2014) proposed some neutrosophic aggregation operators, such as the interval neutrosophic weighted averaging (INWA) operator and the interval neutrosophic weighted geometric (INWG) operator, and applied the operators to solve the multiple attribute group decision making problems with interval neutrosophic information.

For the above researches on the multiple attribute decision making (MADM) problems with interval neutrosophic information, we can suppose the attribute weights are fully known. However, in real decision making, because of time pressure, lack of knowledge or data, and the expert's limited expertise about the problem domain, the information about attribute weights is incompletely known or completely unknown. So, the existing MADM under neutrosophic environment will be impractical for such situations. Therefore, it is necessary to study this issue. In this paper, our aim is to solve the MADM problems in which the attribute values take the form of neutrosophic information and attribute weights are incompletely known or completely unknown based on the maximizing deviation method. In Section 2, we summarize the some basic concepts related to a neutrosophic set and its instances, single valued neutrosophic set and interval neutrosophic set. A score function and an accuracy function is also proposed for ranking neutrosophic numbers in this section. Section 3 introduces the neutrosophic MADM (NMADM) method under neutrosophic environment, in which the information about attribute weights is partly known and the attribute values take the form of neutrosophic numbers. An optimization model based on the maximizing deviation method is established to determine the attribute weights. For the special situations where the information about attribute weights is completely unknown, we develop another optimization model which provides a simple and exact formula. To aggregate the neutrosophic information corresponding to each alternative, we utilize the neutrosophic weighted averaging (NWA) operators which are the single valued neutrosophic weighted averaging (SVNWA) operator and the interval neutrosophic weighted averaging (INWA) operator. Thus we can determine the order of alternatives and choose the most desirable one(s) based on the score function and accuracy function. In Section 4, some illustrative examples are presented to verify the developed approach and to demonstrate its practicality and effectiveness. Section 5 concludes the paper and presents some results.

## 2. Preliminaries

In the subsection, we give some concepts related to neutrosophic sets, single valued neutrosophic sets and interval neutrosophic sets.

### 2.1 Neutrosophic set

**Definition 1.** (Smarandache 1998) Let  $X$  be a universe of discourse, then a neutrosophic set is defined as:

$$A = \{ \langle x, F_A(x), T_A(x), I_A(x) \rangle : x \in X \},$$

which is characterized by a truth-membership function  $T_A: X \rightarrow ]0^-, 1^+[$ , an indeterminacy-membership function  $I_A: X \rightarrow ]0^-, 1^+[$  and a falsity-membership function  $F_A: X \rightarrow ]0^-, 1^+[$ .

There is no restriction on the sum of  $T_A(x)$ ,  $I_A(x)$  and  $F_A(x)$ , so  $0^- \leq \sup T_A(x) + \sup I_A(x) + \sup F_A(x) \leq 3^+$ .

In the following, we adopt the representations  $u_A(x)$ ,  $p_A(x)$  and  $v_A(x)$  instead of  $T_A(x)$ ,  $I_A(x)$  and  $F_A(x)$ , respectively.

Wang et al. (2010) defined the single valued neutrosophic set which is an instance of neutrosophic set as follows:

### 2.1 Single valued neutrosophic sets

**Definition 2.** (Wang et al. 2010) Let  $X$  be a universe of discourse, then a single valued neutrosophic set is defined as:

$$A = \{ \langle x, u_A(x), p_A(x), v_A(x) \rangle : x \in X \}$$

where,  $u_A: X \rightarrow [0,1]$ ,  $p_A: X \rightarrow [0,1]$  and  $v_A: X \rightarrow [0,1]$  with  $0 \leq u_A(x) + p_A(x) + v_A(x) \leq 3$  for all  $x \in X$ . The intervals  $u_A(x)$ ,  $p_A(x)$  and  $v_A(x)$  denote the truth-membership degree, the indeterminacy-membership degree and the falsity membership degree of  $x$  to  $A$ , respectively.

We will denote the set of all the SVNSs in  $X$  by  $\mathcal{Q}$ . A single valued neutrosophic number (SVNN) is denoted by  $\tilde{a} = \langle u, p, v \rangle$  for convenience.

We give a score function and an accuracy function for ranking SVNNs as follows;

**Definition 3.** Let  $\tilde{a} = \langle u, p, v \rangle$  be a single valued neutrosophic number. Then a score function  $S$  of the single valued neutrosophic number can be defined by

$$S(\tilde{a}) = \frac{1 + u - 2p - v}{2} \quad (1)$$

where  $S(\tilde{a}) \in [-1,1]$ .

The score function  $S$  is reduced the score function proposed by Li (2005) if  $p = 0$  and  $u + v \leq 1$ .

**Example 1.** Let  $\tilde{a}_1 = (0.5,0.2,0.6)$  and  $\tilde{a}_2 = (0.6,0.4,0.2)$  be two single valued neutrosophic numbers for two alternatives. Then, by applying *Definition 3*, we can obtain

$$S(\tilde{a}_1) = \frac{1 + 0.5 - 2 \times 0.2 - 0.6}{2} = 0.25$$

$$S(\tilde{a}_2) = \frac{1 + 0.6 - 2 \times 0.4 - 0.2}{2} = 0.30.$$

In this case, we can say that alternative  $\tilde{a}_2$  is better than  $\tilde{a}_1$ .

**Definition 4.** Let  $a = \langle u, p, v \rangle$  be a single valued neutrosophic number, an accuracy function  $H$  of the single valued neutrosophic number can be defined by

$$H(a) = u - p(1 - u) - v(1 - p) \quad (2)$$

where  $H(a) \in [-1,1]$ .

When the value of  $H(\tilde{a})$  increases, we say that the degree of accuracy of the single valued neutrosophic number  $\tilde{a}$  increases.

**Example 2.** Let  $\tilde{a}_1 = (0.3,0.1,0.4)$  and  $\tilde{a}_2 = (0.5,0.1,0.3)$  be two single valued neutrosophic numbers for two alternatives. Then, by applying *Definition 4*, we can obtain  $H(\tilde{a}_1) = -0.13$  and  $H(\tilde{a}_2) = 0.18$ .

In this case, we can say that alternative  $\tilde{a}_2$  is better than  $\tilde{a}_1$ .

With respect to the score function  $S$  and the accuracy function  $H$ , a method for comparing SVNNs can be defined as follows;

**Definition 5.** Let  $\tilde{a}_1 = \langle u_1, p_1, v_1 \rangle$  and  $\tilde{a}_2 = \langle u_2, p_2, v_2 \rangle$  be two single valued neutrosophic values. Then we have

- (1) if  $S(\tilde{a}_1) > S(\tilde{a}_2)$ , then  $\tilde{a}_1$  is greater than  $\tilde{a}_2$ , denoted by  $\tilde{a}_1 \succ \tilde{a}_2$ ,
- (2) if  $S(\tilde{a}_1) = S(\tilde{a}_2)$  and  $H(\tilde{a}_1) > H(\tilde{a}_2)$  then  $\tilde{a}_1$  is greater than  $\tilde{a}_2$ , denoted by  $\tilde{a}_1 \succ \tilde{a}_2$ .

**Example 3.** Let  $\tilde{a}_1 = (0.6,0.2,0.2)$  and  $\tilde{a}_2 = (0.5,0.1,0.3)$  be two single valued neutrosophic numbers for two alternatives. Then, by applying *Definition 5*, we can obtain  $S(\tilde{a}_1) = S(\tilde{a}_2) = 0.5$  and  $H(\tilde{a}_1) = 0.36$ ,  $H(\tilde{a}_2) = 0.18$ . Then it implies that  $\tilde{a}_1 \succ \tilde{a}_2$ .

Based on the study given in Zhang et al. (2014), we define two weighted aggregation operators related to SVNSs as follows;

**Definition 6.** Let  $\tilde{a}_j = \langle u_j, p_j, v_j \rangle (j = 1, 2, \dots, n)$  be a collection of single valued neutrosophic numbers, and SVNWA:  $\mathcal{Q}_n \rightarrow \mathcal{Q}$ , if

$$\text{SVNWA}_\omega(\tilde{a}_1, \tilde{a}_2, \dots, \tilde{a}_n) = \sum_{j=1}^n \omega_j \tilde{a}_j = \left( 1 - \prod_{j=1}^n (1 - u_j)^{\omega_j}, \prod_{j=1}^n p_j^{\omega_j}, \prod_{j=1}^n v_j^{\omega_j} \right) \quad (3)$$

where  $\omega_j$  is the weight of  $\tilde{a}_j (j = 1, 2, \dots, n)$ ,  $\omega_j \in [0, 1]$  and  $\sum_{j=1}^n \omega_j = 1$ , then SVNWA is called single valued neutrosophic weighted average operator. Especially, when  $\omega_j = 1/n (j = 1, 2, \dots, n)$ , then the SVNWA is called an arithmetic average operator for SVNNs.

Similarly, we can define the single valued neutrosophic weighted geometric average (SVNWG) operator.

**Definition 7.** Let  $\tilde{a}_j = \langle u_j, p_j, v_j \rangle (j = 1, 2, \dots, n)$  be a collection of single valued neutrosophic numbers, and SVNWG:  $\mathcal{Q}_n \rightarrow \mathcal{Q}$ , if

$$\text{SVNWG}_\omega(\tilde{a}_1, \tilde{a}_2, \dots, \tilde{a}_n) = \prod_{j=1}^n \tilde{a}_j^{\omega_j} = \left( \prod_{j=1}^n u_j^{\omega_j}, 1 - \prod_{j=1}^n (1 - p_j)^{\omega_j}, 1 - \prod_{j=1}^n (1 - v_j)^{\omega_j} \right) \quad (4)$$

where  $\omega_j$  is the weight of  $\tilde{a}_j (j = 1, 2, \dots, n)$ ,  $\omega_j \in [0, 1]$  and  $\sum_{j=1}^n \omega_j = 1$ , then SVNWG is called single valued neutrosophic weighted geometric average operator. Especially, when  $\omega_j = 1/n (j = 1, 2, \dots, n)$ , then SVNWG is called a geometric average operator for SVNNs.

The aggregation results of the SVNWA and SVNWG operators are still SVNSs.

**Definition 8.** (Majumdar and Samanta 2014) Let  $\tilde{a}_1 = \langle u_1, p_1, v_1 \rangle$  and  $\tilde{a}_2 = \langle u_2, p_2, v_2 \rangle$  be two single valued neutrosophic numbers. Then the normalized Hamming distance measure between  $\tilde{a}_1$  and  $\tilde{a}_2$  is defined as:

$$d(\tilde{a}_1, \tilde{a}_2) = \frac{1}{3} (|u_1 - u_2| + |p_1 - p_2| + |v_1 - v_2|). \quad (5)$$

Wang et al. (2005) extended the concept of single valued neutrosophic set to interval neutrosophic set (INS) which is a further instance of the NSs. The fundamental characteristic of the INS is that the values of its truth-membership function, indeterminacy-membership function and falsity membership function are intervals rather than exact numbers.

## 2.2 Interval neutrosophic sets

**Definition 9.** Wang et al. (2005) Let  $X$  be a universe of discourse and  $\text{Int}[0, 1]$  be the set of all closed subsets of  $[0, 1]$ . Then an interval neutrosophic set is defined as:

$$A = \{ \langle x, u_A(x), p_A(x), v_A(x) \rangle : x \in X \}$$

where  $u_A: X \rightarrow \text{Int}[0, 1]$ ,  $p_A: X \rightarrow \text{Int}[0, 1]$  and  $v_A: X \rightarrow \text{Int}[0, 1]$  with  $0 \leq \sup u_A(x) + \sup p_A(x) + \sup v_A(x) \leq 3$  for all  $x \in X$ . The intervals  $u_A(x)$ ,  $p_A(x)$  and  $v_A(x)$  denote the truth-membership degree, the indeterminacy-membership degree and the falsity membership degree of  $x$  to  $A$ , respectively.

For convenience, if let  $u_A(x) = [u_A^L(x), u_A^U(x)]$ ,  $p_A(x) = [p_A^L(x), p_A^U(x)]$  and  $v_A(x) = [v_A^L(x), v_A^U(x)]$ , then

$$A = \{ \langle x, [u_A^L(x), u_A^U(x)], [p_A^L(x), p_A^U(x)], [v_A^L(x), v_A^U(x)] \rangle : x \in X \}$$

with the condition,  $0 \leq \sup u_A^U(x) + \sup p_A^U(x) + \sup v_A^U(x) \leq 3$  for all  $x \in X$ . Here, we only consider the sub-unitary interval of  $[0, 1]$ . Therefore, an interval neutrosophic set is clearly a neutrosophic set.

We will denote the set of all the INSs in  $X$  by  $\mathcal{F}$ . An interval neutrosophic number (INN) is denoted by  $\tilde{b} = \langle [u^-, u^+], [p^-, p^+], [v^-, v^+] \rangle$  for convenience.

We give the score function and accuracy function of an INN as follows.

**Definition 10.** Let  $\tilde{b} = \langle [u^-, u^+], [p^-, p^+], [v^-, v^+] \rangle$  be an interval neutrosophic number, a score function  $S$  of the single valued neutrosophic number can be defined by

$$S(\tilde{b}) = \frac{2 + u^- + u^+ - 2p^- - 2p^+ - v^- - v^+}{4} \quad (6)$$

where  $S(\tilde{b}) \in [-1, 1]$ .

**Example 4.** Let  $\tilde{b}_1 = ([0.6, 0.4], [0.3, 0.1], [0.1, 0.3])$  and  $\tilde{b}_2 = ([0.1, 0.6], [0.2, 0.3], [0.1, 0.4])$  be two interval neutrosophic numbers for two alternatives. Then, by *Definition 10*, we can obtain  $S(\tilde{b}_1) = 0.65$  and  $S(\tilde{b}_2) = 0.30$ .

In this case we can say that alternative  $\tilde{b}_1$  is better than  $\tilde{b}_2$ .

**Definition 11.** Let  $\tilde{b} = \langle [u^-, u^+], [p^-, p^+], [v^-, v^+] \rangle$  be an interval neutrosophic number, an accuracy function  $H$  of the single valued neutrosophic number can be defined by

$$H(\tilde{b}) = \frac{1}{2}(u^- + u^+ - p^+(1 - u^+) - p^-(1 - u^-) - v^+(1 - p^-) - v^-(1 - p^+)) \quad (7)$$

where  $H(\tilde{b}) \in [-1, 1]$ .

The larger the value of  $H(\tilde{b})$  is, the more the degree of accuracy of the single valued neutrosophic value  $\tilde{b}$  is.

The accuracy function  $H$  is reduced the accuracy function proposed by Nayagam et al. (2011) if  $p^-, p^+ = 0$  and  $u^+ + v^+ \leq 1$ .

With respect to the score function  $S$  and the accuracy function  $H$ , we define a method for comparing INNs as follows;

**Definition 12.** Let  $\tilde{b}_1 = \langle [u_1^-, u_1^+], [p_1^-, p_1^+], [v_1^-, v_1^+] \rangle$  and  $\tilde{b}_2 = \langle [u_2^-, u_2^+], [p_2^-, p_2^+], [v_2^-, v_2^+] \rangle$  be two interval neutrosophic numbers. Then we have

- (1) if  $S(\tilde{b}_1) > S(\tilde{b}_2)$ , then  $\tilde{b}_1$  is greater than  $\tilde{b}_2$ , denoted by  $\tilde{b}_1 > \tilde{b}_2$ ,
- (2) if  $S(\tilde{b}_1) = S(\tilde{b}_2)$  and  $H(\tilde{b}_1) > H(\tilde{b}_2)$  then  $\tilde{b}_1$  is greater than  $\tilde{b}_2$ , denoted by  $\tilde{b}_1 > \tilde{b}_2$ .

Next, we give two weighted aggregation operators related to INNs.

**Definition 13.** (Zhang 2014) Let  $\tilde{b}_j = \langle [u_j^-, u_j^+], [p_j^-, p_j^+], [v_j^-, v_j^+] \rangle$  ( $j = 1, 2, \dots, n$ ) be a collection of interval neutrosophic values, and INWA:  $\mathcal{F}_n \rightarrow \mathcal{F}$ , if

$$\text{INWA}_\omega(\tilde{b}_1, \tilde{b}_2, \dots, \tilde{b}_n) = \sum_{j=1}^n \omega_j \tilde{b}_j = \left( \left[ 1 - \prod_{j=1}^n (1 - u_j^-)^{\omega_j}, 1 - \prod_{j=1}^n (1 - u_j^+)^{\omega_j} \right], \left[ \prod_{j=1}^n (p_j^-)^{\omega_j}, \prod_{j=1}^n (p_j^+)^{\omega_j} \right], \left[ \prod_{j=1}^n (v_j^-)^{\omega_j}, \prod_{j=1}^n (v_j^+)^{\omega_j} \right] \right), \quad (8)$$

where  $\omega_j$  is the weight of  $\tilde{b}_j$  ( $j = 1, 2, \dots, n$ ),  $\omega_j \in [0, 1]$  and  $\sum_{j=1}^n \omega_j = 1$ , then INWA is called interval neutrosophic weighted average operator. Especially, when  $\omega_j = 1/n$  ( $j = 1, 2, \dots, n$ ), then the INWA is called an arithmetic average operator for INNs.

**Definition 14.** (Zhang 2014) Let  $\tilde{b}_j = \langle [u_j^-, u_j^+], [p_j^-, p_j^+], [v_j^-, v_j^+] \rangle$  ( $j = 1, 2, \dots, n$ ) be a collection of interval neutrosophic numbers, and INWG:  $\mathcal{F}_n \rightarrow \mathcal{F}$ , if

$$\begin{aligned}
\text{INWG}_\omega(\tilde{b}_1, \tilde{b}_2, \dots, \tilde{b}_n) &= \prod_{j=1}^n \tilde{b}_j^{\omega_j} \\
&= \left( \left[ \prod_{j=1}^n (u_j^-)^{\omega_j}, \prod_{j=1}^n (u_j^+)^{\omega_j} \right], \left[ 1 - \prod_{j=1}^n (1 - p_j^-)^{\omega_j}, 1 - \prod_{j=1}^n (1 - p_j^+)^{\omega_j} \right], \right. \\
&\quad \left. \left[ 1 - \prod_{j=1}^n (1 - v_j^-)^{\omega_j}, 1 - \prod_{j=1}^n (1 - v_j^+)^{\omega_j} \right] \right), \tag{9}
\end{aligned}$$

where  $\omega_j$  is the weight of  $\tilde{b}_j$  ( $j = 1, 2, \dots, n$ ),  $\omega_j \in [0, 1]$  and  $\sum_{j=1}^n \omega_j = 1$ , then INWG is called interval neutrosophic weighted geometric average operator. Especially, when  $\omega_j = 1/n$  ( $j = 1, 2, \dots, n$ ), then the INWG is called a geometric average operator for INNs.

The aggregation results of the INWA and INWG operators are still INNs.

**Definition 15.** (Ye 2014a) Let  $\tilde{b}_1 = \langle [u_1^-, u_1^+], [p_1^-, p_1^+], [v_1^-, v_1^+] \rangle$  and  $\tilde{b}_2 = \langle [u_2^-, u_2^+], [p_2^-, p_2^+], [v_2^-, v_2^+] \rangle$  be two interval neutrosophic numbers. Then the normalized Hamming distance measure between  $\tilde{b}_1$  and  $\tilde{b}_2$  is defined as:

$$d(\tilde{b}_1, \tilde{b}_2) = \frac{1}{6} (|u_1^- - u_2^-| + |p_1^- - p_2^-| + |v_1^- - v_2^-| + |u_1^+ - u_2^+| + |p_1^+ - p_2^+| + |v_1^+ - v_2^+|).$$

From the above analysis, we develop a method based on the maximizing deviation for the neutrosophic multiple attribute decision making problems in which attribute values for alternatives are the single valued neutrosophic value and the interval neutrosophic value.

### 3. Maximizing deviation method for neutrosophic information

Suppose that  $A = \{A_1, A_2, \dots, A_m\}$  is the set of alternatives and  $C = \{C_1, C_2, \dots, C_n\}$  is a set of criteria or attributes. The attribute weights are partly known or completely unknown. Let  $\omega = (\omega_1, \omega_2, \dots, \omega_n)^T$  be the weight vector of attributes, such that  $\sum_{j=1}^n \omega_j = 1$ ,  $\omega_j \geq 0$  ( $j = 1, 2, \dots, n$ ) and  $\omega_j$  refers to the weight of attribute  $C_j$ .  $D = \{d_1, d_2, \dots, d_t\}$  denotes the set of decision makers (DMs), and  $\lambda = \{\lambda_1, \lambda_2, \dots, \lambda_t\}$  denotes the weight vector of DMs,  $\lambda_t \in [0, 1]$ ,  $k = 1, 2, \dots, t$ ,  $\sum_{k=1}^t \lambda_k = 1$ . Assume that  $A^{(k)} = (a_{ij}^{(k)})_{m \times n}$  is the decision matrix provided by the DM  $d_k \in D$ ,  $a_{ij}^{(k)}$  is a neutrosophic value for alternative  $A_i$  associated with the attribute  $C_j$ . If  $A^{(k)} = (a_{ij}^{(k)})_{m \times n} = (u_{ij}^{(k)}, p_{ij}^{(k)}, v_{ij}^{(k)})_{m \times n}$ , it is a single valued neutrosophic decision matrix, where  $u_{ij}^{(k)}$  indicates the degree that the alternative  $A_i$  satisfies the attribute  $C_j$  and  $p_{ij}^{(k)}$  indicates the degree that the alternative  $A_i$  is indeterminacy on the attribute  $C_j$ , whereas  $v_{ij}^{(k)}$  indicates the degree that the attribute  $A_i$  does not satisfy the attribute  $C_j$  given by the decision-maker  $d_k$ . We have the conditions  $u_{ij}^{(k)} \in [0, 1]$ ,  $p_{ij}^{(k)} \in [0, 1]$ , and  $v_{ij}^{(k)} \in [0, 1]$ ,  $0 \leq u_{ij}^{(k)} + p_{ij}^{(k)} + v_{ij}^{(k)} \leq 3$  for  $i = 1, 2, \dots, m$  and  $j = 1, 2, \dots, n$ . Similarly, if  $A^{(k)} = (a_{ij}^{(k)})_{m \times n} = ([u_{ij}^{(k)-}, (u_{ij}^{(k)+})], [(p_{ij}^{(k)-}, (p_{ij}^{(k)+})], [(v_{ij}^{(k)-}, (v_{ij}^{(k)+})])_{m \times n}$ , it is an interval neutrosophic decision matrix, where  $[(u_{ij}^{(k)-}, (u_{ij}^{(k)+})]$  indicates the degree that the alternative  $A_i$  satisfies the attribute  $C_j$  and  $[(p_{ij}^{(k)-}, (p_{ij}^{(k)+})]$  indicates the degree that the alternative  $A_i$  is indeterminacy on the attribute  $C_j$ , whereas  $[(v_{ij}^{(k)-}, (v_{ij}^{(k)+})]$  indicates the degree that the attribute  $A_i$  does not satisfy the attribute  $C_j$  given by the decision-maker  $d_k$ . Here, we have the condition  $0 \leq \sup(u_{ij}^{(k)+}) + \sup(p_{ij}^{(k)+}) + \sup(v_{ij}^{(k)+}) \leq 3$  for  $i = 1, 2, \dots, m$  and  $j = 1, 2, \dots, n$ .

Obtaining an overall preference value by synthesizing the performance values of all alternatives of each expert is an important step in decision process.

In this paper, we will utilize the **SVNWA** and **INWA** operators as the main aggregation operators for two different **methods**, respectively.

**Definition 16.** Suppose that  $A^{(k)} = (a_{ij}^{(k)})_{m \times n} = (u_{ij}^{(k)}, p_{ij}^{(k)}, v_{ij}^{(k)})_{m \times n}$  is a single valued neutrosophic decision matrix and  $\tilde{r}_i = (\tilde{r}_{i1}, \tilde{r}_{i2}, \dots, \tilde{r}_{in})$  is the vector of attribute values corresponding to the alternative  $A_i$ ,  $i = 1, 2, \dots, m$ . Then the overall preference value of alternative  $A_i$  for DM  $d_k$  can be expressed as,  $i = 1, 2, \dots, m$ ,

$$\tilde{r}_i = (u_{ij}^{(k)}, p_{ij}^{(k)}, v_{ij}^{(k)}) = \text{SVNWA}_\omega(\tilde{r}_{i1}, \tilde{r}_{i2}, \dots, \tilde{r}_{in}) = \left( 1 - \prod_{j=1}^n (1 - u_j^{(k)})^{\omega_j}, \prod_{j=1}^n p_j^{(k)\omega_j}, \prod_{j=1}^n v_j^{(k)\omega_j} \right) \quad (10)$$

where  $\omega = (\omega_1, \omega_2, \dots, \omega_n)^T$  denotes the weight vector of attributes.

**Definition 17.** Suppose that  $A^{(k)} = (a_{ij}^{(k)})_{m \times n} = ([u_{ij}^{-(k)}, u_{ij}^{+(k)}], [p_{ij}^{-(k)}, p_{ij}^{+(k)}], [v_{ij}^{-(k)}, v_{ij}^{+(k)}])_{m \times n}$  is an interval **neutrosophic** decision matrix and  $\tilde{z}_i = (\tilde{z}_{i1}, \tilde{z}_{i2}, \dots, \tilde{z}_{in})$  is the vector of attribute values corresponding to the alternative  $A_i$ ,  $i = 1, 2, \dots, m$ . Then the overall preference value of alternative  $A_i$  for DM  $d_k$  can be expressed as,  $i = 1, 2, \dots, m$ ,

$$\begin{aligned} \tilde{z}_i &= \left( [u_{ij}^{-(k)}, u_{ij}^{+(k)}], [p_{ij}^{-(k)}, p_{ij}^{+(k)}], [v_{ij}^{-(k)}, v_{ij}^{+(k)}] \right) = \text{INWA}_\omega(\tilde{z}_{i1}, \tilde{z}_{i2}, \dots, \tilde{z}_{in}) \\ &= \left( \left[ 1 - \prod_{j=1}^n (1 - u_{ij}^{-(k)})^{\omega_j}, 1 - \prod_{j=1}^n (1 - u_{ij}^{+(k)})^{\omega_j} \right], \right. \\ &\quad \left. \left[ \prod_{j=1}^n (p_{ij}^{-(k)})^{\omega_j}, \prod_{j=1}^n (p_{ij}^{+(k)})^{\omega_j} \right], \left[ \prod_{j=1}^n (v_{ij}^{-(k)})^{\omega_j}, \prod_{j=1}^n (v_{ij}^{+(k)})^{\omega_j} \right] \right), \end{aligned} \quad (11)$$

where  $\omega = (\omega_1, \omega_2, \dots, \omega_n)^T$  be the weight vector of attributes.

Because many practical group decision making problems are complex and uncertain, and human thinking is inherently subjective, the information about attribute weights is usually incomplete. Generally speaking, the incomplete attribute weight information can be expressed as the following relationships among the weights, for  $i \neq j$ :

*Form 1. A weak ranking:*  $\omega_i \geq \omega_j$ ;

*Form 2. A strict ranking:*  $\omega_i - \omega_j \geq \alpha_i$  ( $\alpha_i > 0$ );

*Form 3. A ranking of differences:*  $\omega_i - \omega_j \geq \omega_k - \omega_l$  ( $j \neq k \neq l$ )

*Form 4. A ranking with multiples:*  $\omega_i \geq \alpha_i \omega_j$  ( $0 \leq \alpha_i \leq 1$ )

*Form 5. An interval form:*  $\alpha_i \leq \omega_i \leq \alpha_i + \varepsilon_i$ , ( $0 \leq \alpha_i \leq \alpha_i + \varepsilon_i \leq 1$ ).

Wang (1998) developed the maximizing deviation method for handling the multiple attribute decision making problems characterized by numerical information. In decision making problem, it is essential to rank them by comparing alternatives. The larger the ranking value  $\tilde{r}_i$  (or  $\tilde{z}_i$ ) is, the better corresponding alternative  $A_i$  is. If **an attribute is creating little differences on all alternatives**, it implies that such an attribute has a small important in decision process. Contrary, if an attribute has very clear differences in terms of the performance values of each alternative, we say that such an attribute should be in the foreground in selecting the best alternative. That is, if one attribute has a similar effect among alternatives, it should be **assigned** with a small weight; otherwise the attribute which makes larger deviations should be **assigned** a bigger weight. Especially, if all alternatives have a very similar performance value in term of a given attribute, then such an attribute will not have much effect **on ranking the alternatives**. In other word, such an attribute should be

assigned with a very small weight. Also, Wang (1998) put forward that zero should be assigned with the corresponding to attribute.

To determine the differences among the performance values of all alternatives, we adopt the deviation method. For the DM  $d_k$  and the attribute  $C_j$ , the deviation of alternative  $A_i$  to all the other alternatives can be expressed as follows:

$$H_{ij}^{(k)}(\omega) = \sum_{s=1}^m d(a_{ij}^{(k)}, a_{sj}^{(k)}) \omega_j, \quad \begin{matrix} i = 1, 2, \dots, m \\ j = 1, 2, \dots, n \end{matrix}$$

Let

$$H_j^{(k)}(\omega) = \sum_{i=1}^m H_{ij}^{(k)}(\omega) = \sum_{i=1}^m \sum_{s=1}^m d(a_{ij}^{(k)}, a_{sj}^{(k)}) \omega_j, \quad j = 1, 2, \dots, n.$$

Then  $H_j^{(k)}(\omega)$  gives the deviation value of all alternatives to other alternatives for the attribute  $A_i$  and the DM  $d_k$ .

Using the single valued neutrosophic sets and the interval neutrosophic sets, we can select a weight vector  $\omega$  for maximize operator of all deviation values with respect to all the attributes and all the DMs.

### 3.1 Maximizing deviation method for single valued neutrosophic sets

In the subsection, we construct a non-linear programming model with single valued neutrosophic information, as follows:

$$(M-1) \begin{cases} \max H(\omega) = \sum_{k=1}^t \lambda_k \sum_{j=1}^n \sum_{i=1}^m \sum_{s=1}^m d(a_{ij}^{(k)}, a_{sj}^{(k)}) \omega_j \\ \text{subject to } \omega_j \geq 0, \sum_{j=1}^n \omega_j = 1, \quad j = 1, 2, \dots, n \end{cases} \quad (12)$$

where  $\lambda_k$  is the weight of DM  $d_k$ , and

$$d(a_{ij}^{(k)}, a_{sj}^{(k)}) = \frac{1}{3} (|u_{ij} - u_{sj}| + |p_{ij} - p_{sj}| + |v_{ij} - v_{sj}|).$$

By solving the model (M-1), we get the optimal solution  $\omega = (\omega_1, \omega_2, \dots, \omega_n)^T$ , which can be used as the weight vector of attributes.

If the attribute weights is completely unknown, we can establish another programming model:

$$(M-2) \begin{cases} \max H(\omega) = \sum_{k=1}^t \lambda_k \frac{1}{3} \sum_{j=1}^n \sum_{i=1}^m \sum_{s=1}^m \omega_j (|u_{ij} - u_{sj}| + |p_{ij} - p_{sj}| + |v_{ij} - v_{sj}|) \\ \text{subject to } \omega_j \geq 0, \sum_{j=1}^n \omega_j^2 = 1, \quad j = 1, 2, \dots, n \end{cases}$$

To solve this model, we construct the Lagrange function:

$$L(\omega, \pi) = \sum_{k=1}^t \lambda_k \frac{1}{3} \sum_{j=1}^n \sum_{i=1}^m \sum_{s=1}^m \omega_j (|u_{ij} - u_{sj}| + |p_{ij} - p_{sj}| + |v_{ij} - v_{sj}|) + \frac{\pi}{6} \left( \sum_{j=1}^n \omega_j^2 - 1 \right) \quad (13)$$

where  $\pi$  is the Lagrange multiplier.

Then we compute the partial derivatives of L as follows:



$$\begin{cases} \frac{\partial L}{\partial \omega_j} = \sum_{k=1}^t \lambda_k \sum_{i=1}^m \sum_{s=1}^m (|u_{ij} - u_{sj}| + |p_{ij} - p_{sj}| + |v_{ij} - v_{sj}| + \pi \omega_j = 0) \\ \frac{\partial L}{\partial \pi} = \sum_{j=1}^n \omega_j^2 - 1 = 0 \end{cases}$$

From Eq.(13), we get a simple and exact formula for determining the attribute weights as follows:

$$\omega_j^* = \frac{\sum_{k=1}^t \lambda_k \sum_{i=1}^m \sum_{s=1}^m (|u_{ij} - u_{sj}| + |p_{ij} - p_{sj}| + |v_{ij} - v_{sj}|)}{\sqrt{\sum_{j=1}^n (\sum_{k=1}^t \lambda_k \sum_{i=1}^m \sum_{s=1}^m (|u_{ij} - u_{sj}| + |p_{ij} - p_{sj}| + |v_{ij} - v_{sj}|))^2}}. \quad (14)$$

By normalizing  $\omega_j^*$  ( $j = 1, 2, \dots, n$ ) be a unit, we have

$$\omega_j = \frac{\sum_{k=1}^t \lambda_k \sum_{i=1}^m \sum_{s=1}^m (|u_{ij} - u_{sj}| + |p_{ij} - p_{sj}| + |v_{ij} - v_{sj}|)}{\sum_{j=1}^n \sum_{k=1}^t \lambda_k \sum_{i=1}^m \sum_{s=1}^m (|u_{ij} - u_{sj}| + |p_{ij} - p_{sj}| + |v_{ij} - v_{sj}|)}. \quad (15)$$

### 3.2. Maximizing deviation method for interval neutrosophic sets

Similar to the previous method, we also construct a non-linear programming model with interval neutrosophic information, as follows;

$$(M-3) \begin{cases} \max H(\omega) = \sum_{k=1}^t \lambda_k \sum_{j=1}^n \sum_{i=1}^m \sum_{s=1}^m d(a_{ij}^{(k)}, a_{sj}^{(k)}) \omega_j \\ \text{subject to } \omega_j \geq 0, \sum_{j=1}^n \omega_j = 1, \quad j = 1, 2, \dots, n \end{cases}$$

where  $\lambda_k$  is the weight of DM  $d_k$ , and

$$d(a_{ij}^{(k)}, a_{sj}^{(k)}) = \frac{1}{6} (|u_{ij}^- - u_{sj}^-| + |p_{ij}^- - p_{sj}^-| + |v_{ij}^- - v_{sj}^-| + |u_{ij}^+ - u_{sj}^+| + |p_{ij}^+ - p_{sj}^+| + |v_{ij}^+ - v_{sj}^+|).$$

Solving the model (M-3), we get the optimal solution  $\omega = (\omega_1, \omega_2, \dots, \omega_n)^T$ , which can be used as the weight vector of attributes.

If the attribute weights is completely unknown, we can establish another programming model:

$$(M-4) \begin{cases} \max H(\omega) = \sum_{k=1}^t \lambda_k \frac{1}{6} \sum_{j=1}^n \sum_{i=1}^m \sum_{s=1}^m \omega_j (\Delta_{upv}) \\ \text{subject to } \omega_j \geq 0, \sum_{j=1}^n \omega_j^2 = 1, \quad j = 1, 2, \dots, n \end{cases}$$

where  $\lambda_k$  is the weight of DM  $d_k$  and  $\Delta_{upv} = |u_{ij}^- - u_{sj}^-| + |p_{ij}^- - p_{sj}^-| + |v_{ij}^- - v_{sj}^-| + |u_{ij}^+ - u_{sj}^+| + |p_{ij}^+ - p_{sj}^+| + |v_{ij}^+ - v_{sj}^+|$ .

To solve this model, we construct the Lagrange function:

$$L(\omega, \pi) = \sum_{k=1}^t \lambda_k \frac{1}{6} \sum_{j=1}^n \sum_{i=1}^m \sum_{s=1}^m \omega_j (\Delta_{upv}) + \frac{\pi}{12} \left( \sum_{j=1}^n \omega_j^2 - 1 \right), \quad (16)$$

where  $\pi$  is the Lagrange multiplier and  $\Delta_{upv} = |u_{ij}^- - u_{sj}^-| + |p_{ij}^- - p_{sj}^-| + |v_{ij}^- - v_{sj}^-| + |u_{ij}^+ - u_{sj}^+| + |p_{ij}^+ - p_{sj}^+| + |v_{ij}^+ - v_{sj}^+|$ .

Then we compute the partial derivatives of L as follows:

$$\begin{cases} \frac{\partial L}{\partial \omega_j} = \sum_{k=1}^t \lambda_k \sum_{i=1}^m \sum_{s=1}^m ((\Delta_{upv}) + \pi \omega_j = 0) \\ \frac{\partial L}{\partial \pi} = \sum_{j=1}^n \omega_j^2 - 1 = 0 \end{cases} \quad (17)$$

where,  $\Delta_{upv} = |u_{ij}^- - u_{sj}^-| + |p_{ij}^- - p_{sj}^-| + |v_{ij}^- - v_{sj}^-| + |u_{ij}^+ - u_{sj}^+| + |p_{ij}^+ - p_{sj}^+| + |v_{ij}^+ - v_{sj}^+|$ .

From Eq. (17), we get a simple and exact formula for determining the attribute weights as follows:

$$\omega_j^* = \frac{\sum_{k=1}^t \lambda_k \sum_{i=1}^m \sum_{s=1}^m (\Delta_{upv})}{\sqrt{\sum_{j=1}^n (\sum_{k=1}^t \lambda_k \sum_{i=1}^m \sum_{s=1}^m (\Delta_{upv}))^2}} \quad (18)$$

By normalizing  $\omega_j^*$  ( $j = 1, 2, \dots, n$ ) be a unit, we have

$$\omega_j = \frac{\sum_{k=1}^t \lambda_k \sum_{i=1}^m \sum_{s=1}^m (\Delta_{upv})}{\sum_{j=1}^n \sum_{k=1}^t \lambda_k \sum_{i=1}^m \sum_{s=1}^m (\Delta_{upv})}, \quad (19)$$

where  $\Delta_{upv} = |u_{ij}^- - u_{sj}^-| + |p_{ij}^- - p_{sj}^-| + |v_{ij}^- - v_{sj}^-| + |u_{ij}^+ - u_{sj}^+| + |p_{ij}^+ - p_{sj}^+| + |v_{ij}^+ - v_{sj}^+|$ .

Using by MatLab software with optimization toolbox or Lindo/Lingo software package, the solution of aforementioned maximization problem could be easily solved by a few simple calculations.

With respect to the aforementioned models, we establish a practical and suitable methods for solving the NMADM problems. In our methods, the attribute weights is partly known or completely unknown, and the attribute values are the single valued neutrosophic information or interval neutrosophic information. The methods are described by the following steps:

*Method (1): Maximizing deviation method for single valued neutrosophic sets*

**Step1.** Let  $A^{(k)} = (a_{ij}^{(k)})_{m \times n}$  be a single valued neutrosophic decision matrix, where  $a_{ij}^{(k)} = (u_{ij}^{(k)}, p_{ij}^{(k)}, v_{ij}^{(k)})$  is an attribute value, given by the decision maker  $d_k$ , for the alternative  $A_i$  with respect to the attribute  $C_j$  and  $\tilde{r}_i = (\tilde{r}_{i1}, \tilde{r}_{i2}, \dots, \tilde{r}_{in})$  be the vector of attribute values corresponding to the alternative  $A_i$ .

**Step2.** If the attribute weights are partly known, then we solve the model (M-1) to obtain the attribute weights. If the information about the attribute weights is completely unknown, then we use the model (M-2).

**Step3.** (i) Utilize the weight vector  $\omega = (\omega_1, \omega_2, \dots, \omega_n)^T$  of attributes and by Eq. (10), and obtain the matrix of overall single valued neutrosophic preference values  $\tilde{r}_i$  corresponding to the alternative  $A_i (i = 1, 2, \dots, m)$ .  
(ii) By using the SVNWA operator again and weights of decision makers, compute the collective overall single valued neutrosophic preference values  $\tilde{r}_i$  of alternative  $A_i (i = 1, 2, \dots, m)$ .

**Step4.** Calculate the scores  $S(\tilde{r}_i)$  of the collective overall single valued neutrosophic preference values  $\tilde{r}_i (i = 1, 2, \dots, m)$  to rank all the alternatives  $A_i (i = 1, 2, \dots, m)$  and then to select the best one(s).

If there is no difference between two scores  $S(\tilde{r}_i)$  and  $S(\tilde{r}_j)$ , then we need to calculate the accuracy degrees  $H(\tilde{r}_i)$  and  $H(\tilde{r}_j)$  of the collective overall single valued neutrosophic preference values  $\tilde{r}_i$  and  $\tilde{r}_j$ , respectively, and then rank the alternatives  $A_i$  and  $A_j$  corresponding to the accuracy degrees  $H(\tilde{r}_i)$  and  $H(\tilde{r}_j) (i, j = 1, 2, \dots, m)$ .

**Step 5.** Rank all the alternatives  $A_i (i = 1, 2, \dots, m)$  and select the best one(s) in accordance with  $S(\tilde{r}_i)$  and  $H(\tilde{r}_i)$

**Step 6.** End.

*Method (2): Maximizing deviation method for interval neutrosophic sets*

**Step1.** Let  $A^{(k)} = (a_{ij}^{(k)})_{m \times n}$  be an interval neutrosophic decision matrix, where  $a_{ij}^{(k)} = ([u_{ij}^{-(k)}, u_{ij}^{+(k)}], [p_{ij}^{-(k)}, p_{ij}^{+(k)}], [v_{ij}^{-(k)}, v_{ij}^{+(k)}])$  is an attribute value, given by the decision maker  $d_k$ , for the alternative  $A_i$  with respect to the attribute  $C_j$  and  $\tilde{z}_i = (\tilde{z}_{i1}, \tilde{z}_{i2}, \dots, \tilde{z}_{in})$  be the vector of attribute values corresponding to the alternative  $A_i$ .

**Step2.** If the attribute weights are partly known, then we solve the model (M-3) to obtain the attribute weights. If the information about the attribute weights is completely unknown, then we solve the model (M-4).

**Step3.** (i) Utilize the weight vector  $\omega = (\omega_1, \omega_2, \dots, \omega_n)^T$  of attributes and by Eq. (11), and obtain the matrix of overall interval neutrosophic preference values  $\tilde{z}_i$  corresponding to the alternative  $A_i (i = 1, 2, \dots, m)$ .  
(ii) By using the INWA operator again and weights of decision makers, compute the collective overall interval neutrosophic preference values  $\tilde{z}_i$  of alternative  $A_i (i = 1, 2, \dots, m)$ .

**Step4.** Calculate the scores  $S(\tilde{z}_i)$  of the overall interval neutrosophic preference value  $\tilde{z}_i (i = 1, 2, \dots, m)$  to rank all the alternatives  $A_i (i = 1, 2, \dots, m)$  and then to select the best one(s).

If there is no difference between two scores  $S(\tilde{z}_i)$  and  $S(\tilde{z}_j)$ , then we need to calculate the accuracy degrees  $H(\tilde{z}_i)$  and  $H(\tilde{z}_j)$  of the collective overall interval neutrosophic preference values  $\tilde{z}_i$  and  $\tilde{z}_j$ , respectively, and then rank the alternatives  $A_i$  and  $A_j$  in accordance with the accuracy degrees  $H(\tilde{z}_i)$  and  $H(\tilde{z}_j) (i, j = 1, 2, \dots, m)$ .

**Step 5.** Rank all the alternatives  $A_i (i = 1, 2, \dots, m)$  and select the best one(s) in accordance with  $S(\tilde{z}_i)$  and  $H(\tilde{z}_i)$ .

**Step 6.** End.

#### 4. Numerical examples

**Example 5.** Let us consider decision making problem adapted from Xu and Xia (2012). An automotive company is desired to select the most appropriate supplier for one of the key elements in its manufacturing process. After pre-evaluation, five suppliers have remained as alternatives for further evaluation. In order to evaluate alternative suppliers, a committee composed of three decision makers has been formed. The committee selects four attributes to evaluate the alternatives; (1) product quality  $C_1$ , (2) relationship closeness  $C_2$ , (3) delivery performance  $C_3$ , (4) price  $C_4$ . Decision makers (without loss of generality), whose weight vector is  $\lambda = (\lambda_1, \lambda_2, \lambda_3, \lambda_4) = (\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4})$ , use the single valued neutrosophic values to evaluate the four possible alternatives  $A_i (i = 1, 2, 3, 4)$  under the above four attributes and construct the single valued neutrosophic decision matrices  $A^{(k)} = (a_{ij}^{(k)})_{m \times n} \quad k = (1, 2, 3, 4)$ , as listed in Table 1-4.

**Table 1:** Decision matrices  $A^{(1)}$  given by DM-1

	$C_1$	$C_2$	$C_3$	$C_4$
$A_1$	(0.4,0.2,0.3)	(0.4,0.2,0.3)	(0.2,0.2,0.5)	(0.7,0.2,0.3)
$A_2$	(0.6,0.1,0.2)	(0.6,0.1,0.2)	(0.5,0.2,0.3)	(0.5,0.1,0.2)
$A_3$	(0.3,0.2,0.3)	(0.5,0.2,0.3)	(0.1,0.5,0.2)	(0.1,0.4,0.5)
$A_4$	(0.7,0.2,0.1)	(0.6,0.1,0.2)	(0.4,0.3,0.2)	(0.4,0.5,0.1)

**Table 2:** Decision matrices  $A^{(2)}$  given by DM-2

	$C_1$	$C_2$	$C_3$	$C_4$
$A_1$	(0.1,0.3,0.5)	(0.5,0.1,0.5)	(0.3,0.1,0.6)	(0.4,0.1,0.4)
$A_2$	(0.2,0.5,0.4)	(0.3,0.4,0.3)	(0.2,0.3,0.1)	(0.2,0.3,0.5)
$A_3$	(0.5,0.2,0.6)	(0.2,0.4,0.3)	(0.5,0.2,0.5)	(0.1,0.5,0.3)
$A_4$	(0.2,0.4,0.2)	(0.1,0.1,0.3)	(0.1,0.5,0.4)	(0.5,0.3,0.1)

**Table 3:** Decision matrices  $A^{(3)}$  given by DM-3

	$C_1$	$C_2$	$C_3$	$C_4$
$A_1$	(0.3,0.2,0.1)	(0.3,0.1,0.3)	(0.1,0.4,0.5)	(0.2,0.3,0.5)

$A_2$	(0.6,0.1,0.4)	(0.6,0.4,0.2)	(0.5,0.4,0.1)	(0.5,0.2,0.4)
$A_3$	(0.3,0.3,0.6)	(0.4,0.2,0.4)	(0.2,0.3,0.2)	(0.3,0.5,0.1)
$A_4$	(0.3,0.6,0.1)	(0.5,0.3,0.2)	(0.3,0.3,0.6)	(0.4,0.3,0.2)

**Table 4:** Decision matrices  $A^{(4)}$  given by DM-4

	$C_1$	$C_2$	$C_3$	$C_4$
$A_1$	(0.2,0.2,0.3)	(0.3,0.2,0.3)	(0.2,0.3,0.5)	(0.4,0.2,0.5)
$A_2$	(0.4,0.1,0.2)	(0.6,0.3,0.5)	(0.1,0.2,0.2)	(0.5,0.1,0.2)
$A_3$	(0.3,0.5,0.1)	(0.2,0.2,0.3)	(0.5,0.4,0.3)	(0.5,0.3,0.2)
$A_4$	(0.3,0.1,0.1)	(0.2,0.1,0.4)	(0.2,0.3,0.2)	(0.3,0.1,0.6)

Then, we use the approach developed to obtain the most desirable alternative(s).

**Case1:** Assume that the attribute weights are partly known and the weight information is given as follows:

$$\left\{ \begin{array}{l} 0.18 \leq \omega_1 \leq 0.20, 0.15 \leq \omega_2 \leq 0.25, 0.30 \leq \omega_3 \leq 0.35, 0.30 \leq \omega_4 \leq 0.40, \\ \omega_j \geq 0, \sum_{j=1}^4 \omega_j = 1, \quad j = 1, 2, 3, 4 \end{array} \right\}$$

**Step (1)** Obtain the decision matrix  $A^{(k)} = (a_{ij}^{(k)})_{m \times n}$  given by the DM  $d_k$  and all the components  $a_{ij}^{(k)}$  are single valued neutrosophic values (See Table 1-4).

**Step (2)** Utilize the model (M-1) to establish the following **non-linear** programming model:

$$\left\{ \begin{array}{l} \max H(\omega) = 1.06\omega_1 + 0.83\omega_2 + 1.63\omega_3 + 1.23\omega_4 \\ \text{subject to } \omega_j \geq 0, \sum_{j=1}^n \omega_j = 1, \quad j = 1, 2, \dots, n \end{array} \right.$$

Solving this model, we obtain the weight vector of attributes:  $\omega = (0.18, 0.15, 0.35, 0.32)$ .

**Step (3)** By the weight vector  $\omega = (0.18, 0.15, 0.35, 0.32)$  and by Eq. (10), we obtain the overall single valued neutrosophic preference values  $\tilde{r}_i$  of the alternatives  $A_i (i = 1, 2, 3, 4)$ , as shown in Table 5.

**Table 5:** Matrix of the overall preference values with respect to party known attributes weights

	$d_1$	$d_2$	$d_3$	$d_4$
$\tilde{r}_1$	(0.4684,0.2000,0.3587)	(0.3371,0.1218,0.4962)	(0.2023,0.2616,0.3466)	(0.4142,0.2305,0.4224)
$\tilde{r}_2$	(0.5355,0.1274,0.2305)	(0.2158,0.3434,0.2532)	(0.5355,0.2497,0.2219)	(0.4634,0.1502,0.2294)
$\tilde{r}_3$	(0.2124,0.3440,0.3065)	(0.3524,0.2975,0.4064)	(0.2833,0.3324,0.2166)	(0.3171,0.3422,0.2162)
$\tilde{r}_4$	(0.5016,0.2785,0.1414)	(0.2699,0.3204,0.2170)	(0.3665,0.3399,0.2593)	(0.3866,0.1468,0.2784)

By using the SVNWA operator again (here, take  $\lambda = (\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4})$  as the DM's weight vector), thus we get the collective overall single valued neutrosophic preference values  $\tilde{r}_i$  of alternatives  $A_i$ ,

$$\begin{aligned} \tilde{r}_1 &= (0.3630, 0.1957, 0.4018), \tilde{r}_2 = (0.4511, 0.2013, 0.2335), \\ \tilde{r}_3 &= (0.2931, 0.3285, 0.2763), \tilde{r}_4 = (0.3868, 0.2583, 0.2169). \end{aligned}$$

**Step (4)** Calculate the scores  $S(\tilde{r}_i)$  the collective overall single valued neutrosophic preference values  $\tilde{r}_i (i = 1, 2, 3, 4)$ .

$$S(\tilde{r}_1) = 0.2848, \quad S(\tilde{r}_2) = 0.4074, \quad S(\tilde{r}_3) = 0.1798, \quad S(\tilde{r}_4) = 0.3265.$$

**Step (5)** Rank all the alternatives  $A_i (i = 1, 2, 3, 4)$  in accordance with the scores  $S(\tilde{r}_i) (i = 1, 2, 3, 4)$  of the collective overall single valued neutrosophic preference values  $\tilde{r}_i (i = 1, 2, 3, 4)$ :  $A_2 > A_4 > A_1 > A_3$  and thus  $A_2$  is the most desirable alternative.

**Case2:** If the attribute weights are completely unknown, we propose another approach to determine the most desirable alternative(s).

**Step (1)** See (Step1).

**Step (2)** Utilize the Eq. (15) to obtain the weight vector of attributes:  $\omega = (0.2238, 0.1748, 0.3427, 0.2587)$ .

**Step (3)** Utilize the weight vector  $\omega = (0.2238, 0.1748, 0.3427, 0.2587)$  and by Eq. (10), we obtain the overall single valued neutrosophic preference values  $\tilde{r}_i$  of the alternatives  $A_i (i = 1, 2, 3, 4)$ , as shown in Table 6.

**Table 6:** Matrix of the overall preference values with respect to completely unknown attributes weights

	$d_1$	$d_2$	$d_3$	$d_4$
$\tilde{r}_1$	(0.4465, 0.2000, 0.3573)	(0.3291, 0.1278, 0.5023)	(0.2201, 0.2495, 0.3189)	(0.4329, 0.2298, 0.4078)
$\tilde{r}_2$	(0.5425, 0.1268, 0.2298)	(0.2184, 0.3536, 0.506)	(0.5425, 0.2451, 0.2203)	(0.4571, 0.1536, 0.2347)
$\tilde{r}_3$	(0.2323, 0.3275, 0.2979)	(0.3680, 0.2861, 0.4173)	(0.2867, 0.3189, 0.2412)	(0.2907, 0.3457, 0.2112)
$\tilde{r}_4$	(0.5213, 0.2580, 0.1431)	(0.2470, 0.3145, 0.2275)	(0.3658, 0.3503, 0.2495)	(0.3951, 0.1457, 0.2568)

By using the SVNWA operator again (here, take  $\lambda = (\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4})$  as the DM's weight vector), thus we get the collective overall single valued neutrosophic preference values  $\tilde{r}_i$  of alternatives  $A_i$ ,

$$\tilde{r}_1 = (0.3614, 0.1956, 0.3909), \quad \tilde{r}_2 = (0.4541, 0.2027, 0.2336),$$

$$\tilde{r}_3 = (0.2961, 0.3188, 0.2821), \quad \tilde{r}_4 = (0.3902, 0.2537, 0.2137).$$

**Step (4)** Calculate the scores  $S(\tilde{r}_i)$  of the collective overall single valued neutrosophic preference values  $\tilde{r}_i (i = 1, 2, 3, 4)$ .

$$S(\tilde{r}_1) = 0.2895, \quad S(\tilde{r}_2) = 0.4075, \quad S(\tilde{r}_3) = 0.1881, \quad S(\tilde{r}_4) = 0.3345$$

**Step (5)** Rank all the alternatives  $A_i (i = 1, 2, 3, 4)$  in accordance with the scores  $S(\tilde{r}_i) (i = 1, 2, 3, 4)$  the collective overall single valued neutrosophic preference values  $\tilde{r}_i (i = 1, 2, 3, 4)$ :  $A_2 > A_4 > A_1 > A_3$  and thus the most desirable alternative is  $A_2$ .

**Example 6.** Let us consider decision making problem adapted from Wei et al. (2013). Suppose an organization plans to implement ERP system. The first step is to form a project team that consists of CIO and two senior representatives from user departments. By collecting all possible information about ERP vendors and systems, project team chooses four potential ERP systems  $A_i (i = 1, 2, 3, 4)$  as candidates. The company employs some external professional organizations (or experts) to aid this decision making. The project team selects four attributes to evaluate the alternatives: (1) function and technology  $C_1$ , (2) strategic fitness  $C_2$ , (3) vendor's ability  $C_3$ , and (4) vendor's reputation  $C_4$ . Decision makers (without loss of generality), take weight vector  $\lambda = (\lambda_1, \lambda_2, \lambda_3) = (\frac{1}{3}, \frac{1}{3}, \frac{1}{3})$  and use the interval neutrosophic values to evaluate the four possible alternatives  $A_i (i = 1, 2, 3, 4)$  under the above four attributes and construct the interval neutrosophic decision matrices  $A^{(k)} = (a_{ij}^{(k)})_{m \times n} \quad k = (1, 2, 3)$ , as listed in Table 7-9.

**Table 7:** Decision matrices  $A^{(1)}$  given by DM-1

	$C_1$	$C_2$	$C_3$	$C_4$
$A_1$	([0.4, 0.5], [0.2, 0.3], [0.3, 0.5])	([0.3, 0.4], [0.3, 0.6], [0.2, 0.4])	([0.2, 0.5], [0.2, 0.6], [0.3, 0.5])	([0.5, 0.6], [0.3, 0.5], [0.2, 0.5])
$A_2$	([0.6, 0.7], [0.1, 0.2], [0.2, 0.3])	([0.1, 0.3], [0.1, 0.4], [0.2, 0.5])	([0.4, 0.5], [0.2, 0.5], [0.3, 0.7])	([0.2, 0.4], [0.1, 0.4], [0.3, 0.3])
$A_3$	([0.3, 0.4], [0.2, 0.3], [0.3, 0.4])	([0.3, 0.6], [0.2, 0.3], [0.2, 0.5])	([0.2, 0.7], [0.2, 0.4], [0.3, 0.6])	([0.2, 0.6], [0.4, 0.7], [0.2, 0.7])
$A_4$	([0.2, 0.6], [0.1, 0.1], [0.1, 0.2])	([0.2, 0.5], [0.4, 0.5], [0.1, 0.6])	([0.3, 0.5], [0.1, 0.3], [0.2, 0.2])	([0.4, 0.4], [0.1, 0.6], [0.1, 0.5])

**Table 8:** Decision matrices  $A^{(2)}$  given by DM-2

	$C_1$	$C_2$	$C_3$	$C_4$
$A_1$	([0.4, 0.6], [0.1, 0.3], [0.2, 0.4])	([0.3, 0.5], [0.1, 0.4], [0.3, 0.4])	([0.4, 0.5], [0.2, 0.4], [0.1, 0.3])	([0.3, 0.6], [0.3, 0.6], [0.3, 0.6])

$A_2$	([0.3,0.5], [0.1,0.2], [0.2,0.3])	([0.3,0.4], [0.2,0.2], [0.1,0.3])	([0.2,0.7], [0.3,0.5], [0.3,0.6])	([0.2,0.5], [0.2,0.7], [0.1,0.2])
$A_3$	([0.5,0.6], [0.2,0.3], [0.3,0.4])	([0.1,0.4], [0.1,0.3], [0.3,0.5])	([0.5,0.5], [0.4,0.6], [0.3,0.4])	([0.1,0.2], [0.1,0.4], [0.5,0.6])
$A_4$	([0.3,0.4], [0.1,0.2], [0.1,0.3])	([0.3,0.3], [0.1,0.5], [0.2,0.4])	([0.2,0.3], [0.4,0.5], [0.5,0.6])	([0.3,0.3], [0.2,0.3], [0.1,0.4])

**Table 9:** Decision matrices  $A^{(3)}$  given by DM-3

	$C_1$	$C_2$	$C_3$	$C_4$
$A_1$	([0.1,0.3], [0.2,0.3], [0.4,0.5])	([0.3,0.3], [0.1,0.3], [0.3,0.4])	([0.2,0.6], [0.3,0.5], [0.3,0.5])	([0.4,0.6], [0.3,0.4], [0.2,0.3])
$A_2$	([0.3,0.6], [0.3,0.5], [0.3,0.5])	([0.3,0.4], [0.3,0.4], [0.3,0.5])	([0.3,0.5], [0.2,0.4], [0.1,0.5])	([0.1,0.2], [0.3,0.5], [0.3,0.4])
$A_3$	([0.4,0.5], [0.2,0.4], [0.2,0.4])	([0.2,0.3], [0.1,0.1], [0.3,0.4])	([0.1,0.4], [0.2,0.6], [0.3,0.6])	([0.4,0.5], [0.2,0.6], [0.1,0.3])
$A_4$	([0.2,0.4], [0.3,0.4], [0.1,0.3])	([0.1,0.4], [0.2,0.5], [0.1,0.5])	([0.3,0.6], [0.2,0.4], [0.2,0.2])	([0.2,0.4], [0.3,0.3], [0.2,0.6])

Then, we use the approach developed to obtain the most desirable alternative(s).

**Case1:** Assume that the attribute weights are partly known and the weight information is given as follows:

$$\left\{ \begin{array}{l} 0.16 \leq \omega_1 \leq 0.18, 0.20 \leq \omega_2 \leq 0.25, 0.20 \leq \omega_3 \leq 0.30, 0.35 \leq \omega_4 \leq 0.40 \\ \omega_j \geq 0, \sum_{j=1}^4 \omega_j = 1, \quad j = 1, 2, 3, 4 \end{array} \right\}$$

**Step (1)** Obtain the decision matrix  $A^{(k)} = (a_{ij}^{(k)})_{m \times n}$  given by the DM  $d_k$  and all the components  $a_{ij}^{(k)}$  are interval neutrosophic values (See Table 7-9).

**Step (2)** Utilize the model (M-3) to establish the following non-linear programming model:

$$\left\{ \begin{array}{l} \max H(\omega) = 0.71\omega_1 + 0.69\omega_2 + 0.81\omega_3 + 1.01\omega_4 \\ \text{subject to } \omega_j \geq 0, \sum_{j=1}^n \omega_j = 1, \quad j = 1, 2, \dots, n \end{array} \right.$$

Solving this model, we obtain the weight vector of attributes:  $\omega = (0.16, 0.20, 0.24, 0.40)$ .

**Step (3)** By the weight vector  $\omega = (0.16, 0.20, 0.24, 0.40)$  and by Eq. (11), we obtain the overall interval neutrosophic preference values  $\tilde{z}_i$  of the alternatives  $A_i (i = 1, 2, 3, 4)$ , as shown in Table 10.

**Table 10:** Matrix of the overall preference values with respect to party known attributes weights

	$d_1$	$d_2$	$d_3$
$\tilde{z}_1$	([0.383,0.525], [0.255,0.499], [0.235,0.487])	([0.341,0.558], [0.183,0.449], [0.215,0.439])	([0.292,0.510], [0.225,0.380], [0.267,0.389])
$\tilde{z}_2$	([0.315,0.469], [0.118,0.377], [0.259,0.407])	([0.237,0.541], [0.197,0.411], [0.145,0.301])	([0.225,0.296], [0.272,0.478], [0.230,0.457])
$\tilde{z}_3$	([0.237,0.601], [0.263,0.451], [0.235,0.576])	([0.288,0.396], [0.155,0.397], [0.368,0.491])	([0.299,0.441], [0.174,0.392], [0.181,0.392])
$\tilde{z}_4$	([0.309,0.481], [0.131,0.367], [0.118,0.359])	([0.277,0.317], [0.184,0.352], [0.169,0.421])	([0.209,0.455], [0.250,0.372], [0.155,0.397])

By using the INWA operator again (here, take  $\lambda = (\frac{1}{3}, \frac{1}{3}, \frac{1}{3})$  as the DM's weight vector), thus we get the collective overall interval neutrosophic preference values  $\tilde{z}_i$  of alternatives  $A_i$ ,

$$\begin{aligned} \tilde{z}_1 &= ([0.3401, 0.5318], [0.2196, 0.4406], [0.2388, 0.4345]), \\ \tilde{z}_2 &= ([0.2606, 0.4720], [0.1854, 0.4207], [0.2059, 0.3831]), \\ \tilde{z}_3 &= ([0.2754, 0.4874], [0.1930, 0.4134], [0.2506, 0.4816]), \\ \tilde{z}_4 &= ([0.2654, 0.4218], [0.1829, 0.4469], [0.1462, 0.3922]). \end{aligned}$$

**Step (4)** Compute the scores  $S(\tilde{z}_i)$  of the collective overall interval neutrosophic preference values  $\tilde{z}_i (i = 1, 2, 3, 4)$ .

$$S(\tilde{z}_1) = 0.2194, \quad S(\tilde{z}_2) = 0.2328, \quad S(\tilde{z}_3) = 0.2044, \quad S(\tilde{z}_4) = 0.2222.$$

**Step (5)** Rank all the alternatives  $A_i (i = 1, 2, 3, 4)$  in accordance with the scores  $S(\tilde{z}_i) (i = 1, 2, 3, 4)$  of the collective overall interval valued preference values  $\tilde{z}_i (i = 1, 2, 3, 4)$ :  $A_2 \succ A_4 \succ A_1 \succ A_3$  and thus  $A_2$  is the most desirable alternative.

**Case2:** If the attribute weights are completely unknown, we proposed another approach to determine the most desirable alternative(s).

**Step (1)** See Step (1).

**Step (2)** Utilize the Eq. (19) to obtain the weight vector of attributes:  $\omega = 0.2224, 0.2155, 0.2518, 0.3103$

**Step (3)** Utilize the weight vector  $\omega = 0.2224, 0.2155, 0.2518, 0.3103$  and by Eq. (11), we obtain the overall interval neutrosophic preference values  $\tilde{z}_i$  of the alternatives  $A_i (i = 1, 2, 3, 4)$  (see Table 11).

**Table 11:** Matrices of the overall preference values with respect to completely unknown attributes weights

	$d_1$	$d_2$	$d_3$
$\tilde{z}_1$	([0.369, 0.514], [0.247, 0.485], [0.242, 0.476])	([0.349, 0.556], [0.167, 0.425], [0.207, 0.421])	([0.270, 0.488], [0.216, 0.373], [0.282, 0.406])
$\tilde{z}_2$	([0.345, 0.492], [0.119, 0.362], [0.251, 0.414])	([0.245, 0.542], [0.189, 0.371], [0.153, 0.314])	([0.243, 0.427], [0.270, 0.476], [0.227, 0.466])
$\tilde{z}_3$	([0.245, 0.592], [0.248, 0.419], [0.242, 0.552])	([0.318, 0.427], [0.165, 0.390], [0.351, 0.475])	([0.293, 0.437], [0.172, 0.372], [0.194, 0.405])
$\tilde{z}_4$	([0.292, 0.496], [0.134, 0.325], [0.119, 0.336])	([0.276, 0.323], [0.175, 0.348], [0.174, 0.415])	([0.206, 0.458], [0.248, 0.383], [0.147, 0.375])

By using the INWA operator again (here, take  $\lambda = (\frac{1}{3}, \frac{1}{3}, \frac{1}{3})$  as the DM's weight vector), thus we get the collective overall interval neutrosophic preference values  $\tilde{z}_i$  of alternatives  $A_i$ ,

$$\begin{aligned}\tilde{z}_1 &= ([0.3308, 0.5203], [0.2080, 0.4260], [0.2425, 0.4344]), \\ \tilde{z}_2 &= ([0.2795, 0.4892], [0.1832, 0.4008], [0.2067, 0.3938]), \\ \tilde{z}_3 &= ([0.2861, 0.4915], [0.1922, 0.3941], [0.2555, 0.4745]), \\ \tilde{z}_4 &= ([0.2590, 0.4303], [0.1808, 0.4389], [0.1454, 0.3747]).\end{aligned}$$

**Step (4)** Compute the scores  $S(\tilde{z}_i)$  of the overall interval neutrosophic preference values  $\tilde{z}_i (i = 1, 2, 3, 4)$ .

$$S(\tilde{z}_1) = 0.2265, \quad S(\tilde{z}_2) = 0.2503, \quad S(\tilde{z}_3) = 0.2187, \quad S(\tilde{z}_4) = 0.2324.$$

**Step (5)** Rank all the alternatives  $A_i (i = 1, 2, 3, 4)$  in accordance with the scores  $S(\tilde{z}_i) (i = 1, 2, 3, 4)$  of the overall interval neutrosophic preference values  $\tilde{z}_i (i = 1, 2, 3, 4)$ :  $A_2 > A_4 > A_1 > A_3$  and thus the most desirable alternative is  $A_2$

From the examples, we can see that the proposed neutrosophic decision-making methods are more suitable for real scientific and engineering applications because they can handle not only incomplete information but also the indeterminate information and inconsistent information existing in real situations. Therefore, the technique proposed in this paper extends the existing decision making methods and provides a new way for decision makers.

By a comparative study with existing methods, we can represent the useable and feasibility of the developed group decision-making method. Here, we discuss some methods used to determine the final ranking order of all the alternatives with the single valued neutrosophic information, which are based on the cosine similarity measure and the correlation coefficient (Ye 2013), the weighted cross-entropy (Ye 2014b), the aggregation operators (Ye 2014e) and the outranking approach (Peng et al. 2014, Zhang et al. 2015). In these methods, the weights of decision makers and attribute weights are completely known and the decision process is carried out in the opinion of only a decision maker. In fact, in many MAGDM with neutrosophic information, because of time pressure, lack of knowledge or data, and the decision makers' limited expertise about the problem domain, the information about the weights of decision makers and attributes are incompletely known or completely unknown. Our method has a group decision making approach and utilizes the maximizing deviation method to determine the weight values that are incompletely known or completely unknown of decision makers and attributes, respectively, which is more flexible and reasonable; while the Ye (2013, 2014b, 2014e)'s method, Peng et al. (2014) and Zhang et al. (2015)'s methods ask the decision makers to provide the weight values of decision makers and attributes in advance, which is subjective and sometime cannot yield the persuasive results.

1 With respect to above analyses, a single valued neutrosophic set and an interval neutrosophic set is a special  
2 case of a neutrosophic set, and a neutrosophic set is a set where each element of the universe has the degrees  
3 of truth, indeterminacy and falsity, which lie within  $]0^-, 1^+]$ , the non-standard unit interval. In particular, the  
4 uncertainty presented here, i.e. the indeterminacy factor, is independent of truth and falsity values, whereas  
5 the incorporated uncertainty is dependent on the degree of belongingness and non-belongingness of  
6 intuitionistic fuzzy sets. Therefore, this leads to the theory that intuitionistic fuzzy sets are a special case of  
7 single valued neutrosophic sets. Moreover, SNSs can solve some problems that are beyond the scope of  
8 fuzzy sets and intuitionistic fuzzy sets. Therefore, the proposed MAGDM approach under single valued  
9 neutrosophic environment can be used also to solve MADM problems with fuzzy information and  
10 intuitionistic information. Thus, the comparison shows that our method has its great superiority in handling  
11 the ambiguity and uncertainty inherent in MAGDM problems with neutrosophic information.  
12  
13

## 14 5. Conclusions

15  
16 DMs have a major role to provide the information about alternatives in decision making process. Because of  
17 time pressure, lack of knowledge or data, and the expert's limited expertise about the problem domain, the  
18 information about attribute weights given by DMs is partly known or completely unknown. Recently, some  
19 authors proposed many of methods to overcome the limitations. In this paper, we first defined two  
20 discrimination functions such that score function and accuracy function used to rank the neutrosophic  
21 numbers. Considering by the idea that the attribute with a larger deviation value among alternatives should  
22 be assigned with a larger weight, we then established a method called the maximizing deviation method to  
23 compute the optimal weights of attributes under neutrosophic environment, in which the attribute values are  
24 characterized in terms of neutrosophic values. When aggregating the neutrosophic information  
25 corresponding to each alternative, we utilize the neutrosophic weighted averaging (NWA) operators, the  
26 single valued neutrosophic weighted averaging (SVNWA) operator and the interval neutrosophic weighted  
27 averaging (INWA) operator. Thus one can easily determines the order of alternatives and can chooses the  
28 most desirable one(s) based on the proposed score function and accuracy function. Finally, an application of  
29 developed approach is given to explain its effectiveness and practicality. Our method is straightforward and  
30 has no loss of information. In the future, we shall continue working in application of the neutrosophic  
31 multiple attribute decision-making.  
32  
33  
34  
35  
36  
37

## 38 Acknowledgment

39 This paper is supported by the National Natural Science Foundation of China (Nos. 71471172 and  
40 71271124), and the authors also would like to express appreciation to the anonymous reviewers and Editors  
41 for their very helpful comments that improved the paper.  
42  
43

## 44 References

- 45  
46 Atanassov K (1986) Intuitionistic fuzzy sets. *Fuzzy Sets and Systems*, (20) 87–96.  
47 Broumi S, Ye J and Smarandache F (2015) An Extended TOPSIS Method for Multiple Attribute Decision Making based on Interval  
48 Neutrosophic Uncertain Linguistic Variables, *Neutrosophic Sets and Systems*, 8 23-32.  
49 Broumi S and Smarandache F (2014) Single valued neutrosophic trapezoid linguistic aggregation operators based multi-attribute decision  
50 making, *Bulletin of Pure & Applied Sciences- Mathematics and Statistics*, 33 (2) 135-155.  
51 Broumi S and Smarandache F (2015) New Operations on Interval Neutrosophic Sets, *Journal of new theory*, (1) 24-37.  
52 Li DF (2005) Multiattribute decision making models and methods using intuitionistic fuzzy sets. *Journal of Computer and System Sciences* (70)  
53 73–85.  
54 Lui P, Chu Y, Li Y and Chen Y.(2014) Some generalized neutrosophic number Hamacher aggregation operators and their application to Group  
55 Decision Making, *International Journal of Fuzzy Systems* 16(2)242-255  
56 Liu P and Wang Y (2014) Multiple attribute decision-making method based on single-valued neutrosophic normalized weighted Bonferroni  
57 mean, *Neural Computing and Applications*, 25 (7-8), 2001-2010.  
58 Liu P Shi L (2015) The generalized hybrid weighted average operator based on interval neutrosophic hesitant set and its application to multiple  
59 attribute decision making, *Neural Computing and Applications*, 26 (2) 457-471.  
60 Majumdar P and Samanta SK (2014) On similarity and entropy of neutrosophic sets, *J. Intell. Fuzzy Syst.* 26 (3) 1245–1252.  
61 Nayagam V LG, Muralikrishnan S and Sivaraman G (2011) Multi-criteria decision-making method based on interval-valued intuitionistic fuzzy  
62 sets. *Expert Systems with Applications* 38(3) 1464–1467.  
63  
64  
65



- 1 Peng JJ, Wang JQ, Wang J, Zhang HY and Chen XH (2015a) Simplified neutrosophic sets and their applications in multi-criteria group decision-  
2 making problems, *International Journal of Systems Science*, DOI:10.1080/00207721.2014.994050, 201.
- 3 Peng JJ, Wang JQ, Wang J, Wu, XH and Chen XH (2015b) The Fuzzy Cross-entropy for Intuitionistic Hesitant Fuzzy Sets and its Application in  
4 Multi-criteria Decision -making, *International Journal of Systems Science*, DOI:10.1080/00207721.2014.993744.
- 5 Peng JJ, Wang JQ, Wang J, Wu, XH and Chen XH (2015c) Multi-valued Neutrosophic Sets and Power Aggregation Operators with Their  
6 Applications in Multi-criteria Group Decision- making Problems, *International Journal of Computational Intelligence Systems*, 8(4):345-  
7 363.
- 8 Peng JJ Wang J Zhang HY Chen XH (2014) An outranking approach for multi-criteria decision-making problems with simplified neutrosophic  
9 sets, *Applied Soft Computing*, 25 336-346
- 10 Sahin R and Kucuk A (2014) Subsethood measure for single valued neutrosophic sets, *Journal of Intelligent and Fuzzy Systems*, DOI:  
11 10.3233/IFS-141304.
- 12 Smarandache, F (1998) *Neutrosophy. Neutrosophic Probability, Set, and Logic*, Amer. Res. Press, Rehoboth, USA, 105 p.
- 13 Xu, Z. S., and Xia, M. M. (2012). Hesitant fuzzy entropy measures and their use in multi-attribute decision making. *International Journal of*  
14 *Intelligent Systems* (27) 799–822.
- 15 Tian ZP Wang J Zhang HY Chen XH, Wang JQ (2015) Simplified neutrosophic linguistic normalized weighted Bonferroni mean operator and its  
16 application to multi-criteria decision-making problems, *FILOMAT*.
- 17 Wang H Smarandache F Zhang YQ and Sunderraman R (2010) Single valued neutrosophic sets, *Multispace and Multistructure* (4) 410-413.
- 18 Wang H Smarandache F Zhang YQ Sunderraman R (2005) Interval neutrosophic sets and logic: theory and applications in computing. Hexis,  
19 Arizona.
- 20 Wang YM (1998) Using the method of maximizing deviations to make decision for multi-indices, *System Engineering and Electronics* (7) 24–26
- 21 Wang J Zhou P Li KJ Zhang HY Chen XH (2014) Multi-criteria decision-making method based on normal intuitionistic fuzzy-induced  
22 generalized aggregation operator, *TOP*, 22 1103-1122.
- 23 Wang J Nie R Zhang H Chen X (2013a) Intuitionistic fuzzy multi-criteria decision-making method based on evidential reasoning, *Applied Soft*  
24 *Computing*, 13(4), 1823-1831.
- 25 Wang J Nie R Zhang H Chen X (2013b) New operators on triangular intuitionistic fuzzy numbers and their applications in system fault analysis,  
26 *Information Sciences*, 251, 79-95
- 27 Wei G, Zhao X, Lin R and Wang H (2013) Uncertain linguistic Bonferroni mean operators and their application to multiple attribute decision  
28 making, *Applied Mathematical Modelling* 37(7) 5277–5285.
- 29 Ye J (2013) Multicriteria decision-making method using the correlation coefficient under single-valued neutrosophic environment, *International*  
30 *Journal of General Systems* 42(4) 386-394.
- 31 Ye J (2014a) Similarity measures between interval neutrosophic sets and their applications in Multi-criteria decision-making. *Journal of*  
32 *Intelligent and Fuzzy Systems*, (26) 165-172.
- 33 Ye J (2014b) Single valued neutrosophic cross-entropy for multi-criteria decision making problems, *Appl. Math. Model.* 38 (3) 1170–1175.
- 34 Ye J (2014c) Trapezoidal neutrosophic set and its application to multiple attribute decision making, *Neural Computing and Applications*, DOI:  
35 10.1007/s00521-014-1787-6
- 36 Ye J (2014d) Some aggregation operators of interval neutrosophic linguistic numbers for multiple attribute decision making, *Journal of Intelligent*  
37 *& Fuzzy Systems* (27) 2231-2241.
- 38 Ye J (2014e) A multicriteria decision-making method using aggregation operators for simplified neutrosophic sets, *J. Intell. Fuzzy Syst.* 26 (5)  
39 2459–2466.
- 40 Zadeh LA (1965) Fuzzy sets, *Inf. Control* 8 338–353.
- 41 Zhang HY, Wang JQ, Chen XH (2014) Interval neutrosophic sets and their application in multicriteria decision making problems. *The Scientific*  
42 *World Journal*. DOI: 10.1155/2014/645953.
- 43 Zhang HY Wang J Chen XH (2015) An Outranking Approach for Multi-criteria Decision-making Problems with Interval-valued  
44 Neutrosophic Sets, *Neural Computing and Applications*, DOI: 10.1007/s00521-015-1882-3.
- 45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65