

-(Kushik) Kaushal Timilsina

Riemann's Hypothesis and Manifolds


(-not an attempt to prove, just a new perspective)

"An infinite dimensional manifold is a ^{compact} surface by itself"

Part 1: The identification of the origin

(that encloses nothing)

1.1) Real line




1.2) Fold the real line into a coil as:

a) start with a point and fix the origin to it.

b) Rotate clockwise on a spiral about a center (not yet clear), and map $(0, \infty)$ on the clockwise rotation; moving inwards on the spiral.

c) Rotate counterclockwise starting at 0 and map $(-\infty, 0)$

("run to $-\infty$ from 0"); moving outwards on the spiral.



1.3) Define a map from the part of the spiral $[0, \infty)$ to the complex plane, identifying each point with a complex number

$S = a + ib$. ("take the spiral to be a disc")

This is pretty sloppy, as the center is not well defined; which should be done but

I am unable to do it.

Question: Can you identify the center to the origin (0); this would mean to convert the real line into a closed curve, given that you can fix ~~the~~ a finite radius (0 and center relation) to define the center of the spiral from the origin (0).

- To accommodate the whole real line, the torsion ("not exactly") of the curve is defined accordingly.

Hypothesis (1) Whether you can form a closed curve by identification will depend on where you start (0, center relation). Only special maps can define the identification.

I'm just being verbose here, but this could probably require non-linear mapping or some kind of non-Cauchy convergence on partitions of the domain $[0, \infty)$.

Part 2: Mysterious: homeomorphism? diffeomorphism? or homeomorphism?

2.1) consider $V \subset \mathbb{R}^n$ (n dimensional Euclidean space).

2.2) Can you construct a vector field, V' at every point in V such that an orthonormal basis (linearly independent could be enough) $\{e_1, e_2, \dots, e_n\}$ at a point p , can span V' at every point? ("You can construct the basis at other points by linear combination of $\{e_1, e_2, \dots, e_n\}$ ")

2.3) Define $\varphi: V \rightarrow \mathbb{M}^n$; which is already constructed as $f_\alpha: U_\alpha \rightarrow \mathbb{M}^n, \alpha: 1, \dots, n$ where U_α are open sets (domains). (" φ is a map from V to an already well defined manifold M ").


2.4) Assign norm 1 vectors $\{f_1, f_2, \dots, f_n\}$ to U_α 's such that for the set of points $\{q_1, q_2, q_3, \dots, q_n \mid q_i \in U_i\}$, the map $\phi_i: q_i \rightarrow q_i \cdot f_i$ (where dot is the scalar product) is defined such that the set of vectors $\{q_i \cdot f_i \mid i=1, \dots, n\}$ form an n -dimensional vector space W , such that $q_i \cdot f_i$ are linearly independent.

2.5) ~~what~~

2.5) A map from $V' \rightarrow W$ can be defined by composition.
 $\{e_i\}$ on V' were defined orthonormal by brute force. $\{q_i h_i\}$ on W were defined orthonormal by brute force.
 linearly independent.

2.6) Define a homomorphism $\omega: V' \rightarrow W$ if for any operation defined on $\{e_1, e_2, \dots, e_n\}$ at p with between a vector in V' at a point p in V and a vector in W , and between a vector in V' at another point q in V and the same vector in W give the same result.
 $\omega(\sum_i d_i e_{ip}, \sum_i q_i h_i) = \text{operation}(\sum_i d_i e_{iq}, \sum_i q_i h_i)$
 $\forall p, q$ in V . ("back to the question in 2.2")

Part 3: Riemann's hypothesis zeta function.

3.1) n linearly dependent vectors may add upto 0 as do the n th roots of unity. 

3.2) The Riemann zeta function:

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}, \text{Re}(s) > 1.$$

3.3) Definition: ~~Consider~~
 Consider the sum, $1^{-s} + 2^{-s} + 3^{-s} + \dots + n^{-s}$
 define: $Z_1 = (|g_{11}|)^{-s} + (|g_{12}| + |g_{22}|)^{-s} + \dots$
 $+ (|g_{1n}| + |g_{2n}| + \dots + |g_{nn}|)^{-s}$
 where g_{ij} is the i th orthonormal basis of

V^j (j dimensional vector space) such that
 $V^j \subset \mathbb{F}^n, \forall j$ and V^j is a subspace of
 \mathbb{F}^n , where \mathbb{F}^n is defined as $\underbrace{\mathbb{F} \times \mathbb{F} \times \dots \times \mathbb{F}}_n$,
 \mathbb{F} is a field.

3.4) Definition: $Z_2 = (|g_{11}| + (g_{12} + g_{22}) + (g_{13} + g_{23} + g_{33}) + \dots + (g_{1n} + g_{2n} + \dots + g_{nn})|^{-s})$ with g_{ij} 's as defined in 3.3.

3.5) If $Z_2 \neq 0$, \mathbb{F}^n direct sum of V^j 's over j
 $= \mathbb{F}^n$ and V^j 's are subspaces that are
disjoint in some sense, $\cup V^j$'s is ~~the~~ a
partition of \mathbb{F}^n which is equal to \mathbb{F}^n .
If $Z_2 = 0$, the V^j 's are not disjoint.
This establishes some form of
compactness.

Part 4: The new perspective.

4.1) Define the construction in part 2 using U_α 's (2.3) as different refinements of the closed curve in part 1 (whose origin is identified with the center).

4.2) Define an operation (2.6)

$$\left(\sum_i^n \alpha_i e_{ip}, \sum_j^n q_j h_j \right) = \left(\sum_i^n \alpha_i e_{ip} \right) \wedge \left(\sum_j^n q_j h_j \right)$$

where \wedge is some sort of exterior derivative product (by definition).

(Because it gives out a vector and would be 0 if $\sum_i^n e_{ip} \alpha_i$ was in some sense parallel to $\sum_i^n q_i h_i$.)

~~with the bilinearity of \wedge on the component~~

$$\Rightarrow \text{operation } \left(\sum_i^n \alpha_i e_{ip}, \sum_i^n q_i h_i \right) = \alpha_i q_i \sum_i^n e_i h_i$$

~~(and now we drop p in e_i using 2.6 definition of homomorphism.)~~

4.3) Plug $\alpha_i q_i \sum_i^n e_i h_i$

$$4.3) \text{ plug } \left(\sum_i^n \alpha_i e_i \right) \wedge \left(\sum_j^n q_j h_j \right)$$

(dropped the p on e_i by 2.6 definition of homomorphism)

into Z_1 and Z_3 such that

g_{ij} is the i 'th component of

$$\left(\sum_i^n \alpha_i e_i \right) \wedge \left(\sum_j^n q_j h_j \right)$$

("without defining the i^{th} unit vector, this is a pretty big formula").

4.4) Hypothesis: with this setup, if $Z_1 = Z_2$ for a finite integer n , (some kind of saturation of the triangle inequality) $|\vec{a} + \vec{b}| \leq |\vec{a}| + |\vec{b}|$, in some sense

on a finite dimensional manifold M^n you can use U_α from the definition of M^n you can form a V (as in part 2) that can ~~cover~~ be defined at every point in M .

(In part 2, ~~things~~ ~~were~~ the relation between U_α 's and V was defined by construction. But here we get V from U_α 's.)

Hypothesis (2): $Z_1 = Z_2 = 0 \Leftrightarrow M^n$ is closed
(M^n is a compact surface.)

4.5) Define $Z_3 = (|g_{11}|)^s + (|g_{12}| + |g_{22}|)^s + \dots$
 \dots infinitely many such that

$$Z_3 = \lim_{n \rightarrow \infty} Z_n$$

$$Z_4 = (|g_{11}| + |g_{12}| + |g_{22}| + |g_{13}| + |g_{23}| + |g_{33}|) \dots \quad (s)$$

$$Z_4 = \lim_{n \rightarrow \infty} Z_n$$

(the definition of limit is sloppy here as indicated at the end of part 1, it is not clearly defined here based on convergence.)

Hypothesis (3): Riemann's hypothesis $\Leftrightarrow \zeta_3 = \zeta_4 = 0$
zeroes the existence of

which is to say that \sim Riemann's hypothesis zeroes
implies (and vice versa) that an infinite zeroes
dimensional manifold is a compact surface
(closed). Where? (what is the ambient space?)
Well, by itself is the answer (possibly).

Rephrase: An infinite dimensional manifold is
a compact surface by itself. (but there is nothing
that it encloses.)

Summary about this perspective

Riemann zeta function.

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}$$

- Many other perspectives on the zeta function try to explore the parameter, s .
- This perspective explore the possibilities on two other parameters of the zeta function - (i) n being natural numbers (ii) the sum being infinite.


- Hypothesis (3) is based on Hypothesis (2) and Hypothesis (2) is based on Hypothesis (1).

which is to say that if there exists and can be defined an identification as defined in part 1, Hypothesis (2) and Hypothesis (3) are implied.

- Indeed the zeta function depends on the parameter s as well, which could be describing the uniqueness of Riemann's zeroes.

Part 5: Motivations:

(5.1) If ^{one of the principal} curvature of a plane (S^2) blows up \rightarrow it forms a 1 dimensional line, locally, which is a 1 dimensional singularity in S^2 .




(5.2) The singularity of a black hole is where the 4d spacetime (Lorentzian manifold) forms (collapses into) a singularity (intuitively 0 dimensional, but I don't really know) 1d singularity.

(5.3) Speculations on what lies on the "other side" of the singularity include wide range of notions.

(5.4) But if you collapse an infinite dimensional manifold into a singularity, perhaps you might not come out some where but ~~some~~ singularity is another singularity on the same manifold.

"sloppy figure" to demonstrate, two points ^{both} on the "outer surface" of the sphere identified by irregularity.



Even though you cannot ^(possibly) go there, what lies beyond a ~~is~~ singularity in an infinite dimensional manifold is the same manifold.

"Intuitively its just a surface, there is no inside"

(5.5) Question: Can two manifolds of different dimension be connected by a singularity? ~~no~~

("Is there a 100 dimensional universe beyond that black hole?", well how about the other one?")

(5.6) It can be difficult to start with this question for two finite dimensional surfaces. ^{manifolds} So we start with infinite two dimensional manifolds because Hypothesis (3) [the new perspective on the Riemann hypothesis], makes it easier.

Hypothesis (4) Two infinite dimensional manifolds M^m, N^n can be connected by a singularity. | Hypothesis (4) is based on Hypothesis (3).

I have not yet understood how infinite dimensional manifolds are generally treated, but this hypothesis would be constructed based on Cantor's theorem.

(5.7) Hypothesis (5) An infinite dimensional "sphere" (sloppy) turned inside out or, An infinite dimensional "It has the same information on the outer surface as in the inner surface and indeed ~~the~~ the surfaces can be identified and hence there is no inside the sphere, it's just the surface!"
Hypothesis (5) is based on hypothesis (4).

(5.7). Hypothesis (6): A globally orientable ~~compact~~ manifold can store global information iff it is globally compact. A locally orientable manifold can store local information if it is locally compact.

Question: Can infinite dimensional manifolds store global information?

- it seems like the answer is No, but local information yes.

- this question has major implications for defining conservation laws on manifolds.

(5.8) I do not yet understand clearly how loops in spacetime are realized. But this setting, can define loops in spacetime as intersection between spacetime manifold and itself. It doesn't always have to be ~~inf~~ 0-dimensional singularity, and neither does it even have to be a singularity. If we can work on infinite dimensional manifolds with singularities, we can soon develop intersections of finite dimensional manifolds (which includes self intersection)

This will help us explore notions of higher dimensions, and ideas of creation, annihilation and conservation.

(5.9) Aim: The aim of this discussion is to build an understanding towards "A Theory of intersection of manifolds"

(5.10) In this discussion, $[0, \infty)$ was collapsed into a disc in complex coordinates which would later be used to build U_α 's. Its position $s = at + ib$ also plays an important part in the Riemann zeta function,

- Is it possible to collapse the whole complex plane into a similar spiral (or perhaps a stack of discs)?

In general, can you collapse any topological space into an analogous structure ("make it closed" is the heart of the problem) and use it to build a more abstract version of a manifold?

- For example the Kähler manifold is defined with a 2-form structure. (Can you ~~define~~ work with constructions from n -form structures?)

(5.11) Corollary of Hypothesis (6):

Black holes can not store local information but they store global information. So perhaps one may not traverse beyond the singularity, but two people jumping into "entangled" black holes can meet at the center $\Leftrightarrow (ER = EPR)$.

- In the formulation of ER = EPR, "a theory of intersection of manifolds" will help us better understand what entanglement geometrically means.

(5.12) Integrating geometric flows in a theory of intersection of manifolds could describe the foundations of the dynamics of interactions between manifolds; setting a new path for theories of quantum gravity.

(5.13). The fact that the Riemann zeta function describes the distribution of prime numbers, could hint at the spiral used in part 1 to be described by a map from the real line that preserves the distribution of primes. This could mean that this map could be a function like for example, von Mangoldt function, which behaves differently for primes and non-primes. This could describe the relationship between s in the zeta function and the different orders of infinities that we have used in defining infinite dimensional manifolds. The fact that $\zeta(s) \neq 0$ could tell us something like "there is no set whose cardinality is strictly between the cardinality of natural numbers and the cardinality of the set of real numbers"; helping extend this perspective to the Riemann hypothesis.