

On the L-functions from Generalized Riemann hypothesis, Birch and Swinnerton-Dyer conjecture, and the Prime numbers from Polignac's and Twin prime conjectures

By John Y. C. Ting, the corresponding author who contributes to all work in this research paper.

Abstract

Dirichlet eta function (proxy function for Riemann zeta function as generating function for all nontrivial zeros) and Sieve of Eratosthenes (generating algorithm for all prime numbers) are essentially infinite series. We apply infinitesimals to their outputs and analyze L-functions of elliptic curves for Birch and Swinnerton-Dyer conjecture. Riemann hypothesis asserts the complete set of all nontrivial zeros from Riemann zeta function is located on its critical line. It is proven to be true when usefully regarded as an Incompletely Predictable Problem. We ignore even prime number 2. The complete set with derived subsets of Odd Primes contain arbitrarily large number of elements and satisfy Prime number theorem for Arithmetic Progressions, Generic Squeeze theorem and Theorem of Divergent-to-Convergent series conversion for Prime numbers. Having these theorems satisfied by all Odd Primes, Polignac's and Twin prime conjectures are separately proven to be true when usefully regarded as Incompletely Predictable Problems.

Contents

1. Generic Introduction to Generic Numbers, Generic Terms and Generic Equations	2
1.1. Linking Generalized Riemann hypothesis to Birch and Swinnerton-Dyer conjecture	8
2. General notations including Prime number theorem for Arithmetic Progressions and creating <i>de novo</i> Infinite Series	24
3. Generic Squeeze theorem as a novel mathematical tool in Number theory	32
4. Applying the Theorem of Divergent-to-Convergent series conversion for Prime numbers to Polignac's and Twin prime conjectures	36
5. Subtypes of Countably Infinite Sets with Incompletely Predictable entities from Riemann zeta function and Sieve of Eratosthenes	39

Keywords: Birch and Swinnerton-Dyer conjecture, Centroid of n -dimensional geometric object, Hodge conjecture, Polignac's and Twin prime conjectures, Generalized Riemann hypothesis, Theory of Symmetry from Langlands program

AMS Classification: Primary: 11M26, 11A41; Secondary: 81Q10.

©John Yuk Ching Ting.

6. Conclusions including applying infinitesimals to outputs from Sieve of Eratosthenes and Riemann zeta function	41
Acknowledgements and Declarations	45
References	46
Appendix A. Predictability properties of Dirichlet L -series from Dirichlet L -functions that are relevant to Riemann hypothesis, and Birch and Swinnerton-Dyer conjecture	46

1. Generic Introduction to Generic Numbers, Generic Terms and Generic Equations

Generic Summary of this research paper: Polignac's and Twin prime conjectures are proposals on the property [known as Cardinality] of a particular **Generic Number** [known as Prime numbers]. Here, Twin prime conjecture (on even Prime gap = 2) is a subset of Polignac's conjecture (on all even Prime gaps 2, 4, 6, 8, 10...). Generic L-functions, that must obey three Axioms [stated in Remark 1.2], are an important type of **Generic Equations** [with infinitely-many **Generic Terms**] having major properties of Algebraic rank, Analytic rank, trivial zeros and nontrivial zeros. Generalized Riemann hypothesis and "subset" Riemann hypothesis are conjectures on, respectively, all (Generic) L-functions and two specific "prototypical" L-functions (known as Riemann zeta function and its *proxy* [via Analytic continuation] Dirichlet eta function, which are often [*incorrectly*] interchanged with each other). Together, they conjectured entire sets of unique nontrivial zeros of any L-function must always be located on the so-called $\sigma = \frac{1}{2}$ -Critical Line. An important caveat: Riemann zeta function DO NOT have any nontrivial zeros and DO NOT have its first trivial zero given by $s = 0$ *versus* Dirichlet eta function DO have all nontrivial zeros [albeit its first nontrivial zero is NOT given by $t = 0$] and DO have its first trivial zero given by $s = 0$. Birch and Swinnerton-Dyer (BSD) conjecture concern two nominated properties of a particular type of "more complex" L-functions present in all Elliptic curves [classified as degree 2; based on degree of these L-functions being the number $J + 2K$ of Gamma factors occurring in their functional equations]. More specifically, it (mainly) proposes these particular L-functions that are "individualized" for each and every Elliptic curves have Algebraic rank = Analytic rank [\equiv Mordell-Weil rank]. Mordell-Weil theorem states that the set of rational points on an abelian variety over a number field forms a finitely generated abelian group, hence isomorphic to a group of the form $T \oplus \mathbb{Z}^r$, where T is a finite torsion group. The integer $r \geq 0$ is the Mordell-Weil rank of abelian variety [whereby Elliptic functions with their Elliptic curves are abelian variety]. Mordell-Weil rank of ever higher r values; viz, representing ever higher Analytic rank values for Elliptic curves, can technically be extremely difficult to compute.

L-functions for elliptic curves, as function of complex variable $s = \sigma \pm it$, can have Analytic rank of zero or non-zero integer values. Analytic rank = 0 \implies associated L-functions NEVER have first nontrivial zero given by real variable $t = 0$. Analytic rank ≥ 1 [viz, 1, 2, 3, 4, 5... up to an arbitrarily large number value] \implies associated L-functions ALWAYS have first nontrivial zero given by real variable

$t = 0$. We immediately recognize some overlapping aspects of the above major conjectures detailed as follows: The two L-functions from Riemann hypothesis also possess Analytic rank [= 0 in both cases]. The L-functions of every Elliptic curves also have unique nontrivial zeros (spectrum) that are all located on [Analytically normalized] $\sigma = \frac{1}{2}$ -Critical Line. L-functions of various Elliptic curves can have Analytic rank of 0, 1, 2, 3, 4, 5... whereby we expect greater "frequency" and "complexity" and/or altered "internal symmetry" in nontrivial zeros (spectrums) associated with increasing values of Analytic rank.

Symmetric power of an L-function: Let $L(s)$ be an L-function given by an Euler product $L(s) = \prod_{p \notin S} \prod_{j=1}^r \left(1 - \frac{\alpha_j}{p^s}\right)^{-1} \times \prod_{p \in S} L_p(s)$, where S is a finite set of primes. The symmetric n^{th} power of $L(s)$ is an

L-function given by an Euler product $L(s, \text{sym}^n) = \prod_{p \notin S} \prod_{\substack{\text{degree-}n \\ \text{monomials } m}} \left(1 - \frac{m(\alpha_1, \dots, \alpha_r)}{p^s}\right)^{-1} \times \prod_{p \in S} L_p(s, \text{sym}^n)$.

The Euler factors at the primes $p \in S$ (the "bad" primes) are computed via a more complicated recipe which involves a non-trivial amount of information about the underlying object. The degree of an Euler factor at one of the bad primes will be smaller than the degree of the Euler factors outside the set S .

Remark 1.1. Formal statements on Birch and Swinnerton-Dyer conjecture: The central value of an L-function, which could be computed using the formula mentioned below, is its value at central point of Critical Strip. The central point of an L-function is the point on real axis of Critical Line. Equivalently, it is the fixed point of functional equation. In Analytic normalization, the central point is $s = \frac{1}{2}$. In Arithmetic normalization, it is $s = \frac{w+1}{2}$, where w is the weight of L-function. Rodriguez-Villegas and Zagier[5] have proven a formula, conjectured by Gross and Zagier[1], for central value of $L(s, \chi^{2n-1})$, namely $L(1/2, \chi^{2n-1}) = 2 \frac{(2\pi \sqrt{7})^n \Omega^{2n-1} A(n)}{(n-1)!}$ where $\Omega = \frac{\Gamma(1/7)\Gamma(2/7)\Gamma(4/7)}{4\pi^2}$. Using their notation, they variously prove that $A(n) = B(n)^2$ where $B(1) = 1/2$ and $B(n)$ is an integer for $n > 1$; and that $A(n)$ is given by a remarkable recursion formula [not stated here in this paper]. Then the accompanying incredible [derived] result, in one fell swoop, proves the non-vanishing of $L(1/2, \chi^{2n-1})$ for all odd n .

BSD conjecture relates the order of vanishing (or analytic rank) and the leading coefficient of the L-function associated to an elliptic curve E defined over a number field K at central point $s = 1$ to certain arithmetic data, the BSD invariants of E .

The *weak* form of the BSD conjecture states just that the analytic rank r_{an} (that is, the order of vanishing of vanishing of $L(E, s)$ at $s = 1$), is equal to the rank r of E/K .

The *strong* form of the BSD conjecture states also that the leading coefficient of the L-function is given by the formula $\frac{1}{r!} L^{(r)}(E, 1) = |d_K|^{1/2} \cdot \frac{\#\text{III}(E/K) \cdot \Omega(E/K) \cdot \text{Reg}(E/K) \cdot \prod_{\mathfrak{p}} c_{\mathfrak{p}}}{\#E(K)_{\text{tor}}^2}$.

The quantities appearing in this formula are as follows: d_K is the discriminant of K ; r is the rank of $E(K)$; $\text{III}(E/K)$ is the Tate-Shafarevich group of E/K ; $\text{Reg}(E/K)$ is the regulator of E/K ; $\Omega(E/K)$ is the global period of E/K ; $c_{\mathfrak{p}}$ is the Tamagawa number of E at each prime \mathfrak{p} of K ; $E(K)_{\text{tor}}$ is the torsion order of $E(K)$.

Implicit in the strong form of BSD conjecture is that the Tate-Sharafevich group $\text{III}(E/K)$ is finite. There is a similar conjecture for abelian varieties over number fields.

We assign the defining polynomial [see Remark 1.3 below] to be univariate Polynomial $[x]$ for Riemann zeta function when based on linear equation " P " $x = 0$. We [arbitrarily] assign the defining polynomial to be univariate Polynomial $[\pm x]$ for its *proxy* function Dirichlet eta function when based on linear equation " D " $\pm x = 0$. An elliptic curve " E " over a field K is a smooth projective curve of genus 1 together with a distinguished point O [viz, the unique point at infinity serving as the identity element]; whereby E over rational numbers \mathbb{Q} has a Weierstrass equation of the form $E: y^2 = x^3 + ax + b$ with $a, b \in \mathbb{Z}$ such that its discriminant $\Delta := -16(4a^3 + 27b^2) \neq 0$ [viz, being square-free in x with the curve being non-singular]. This bivariate equation E can also be written as $y^2 - x^3 - ax - b = 0$. Elliptic curves are abelian variety; viz, they have an algebraically defined group law with respect to being an abelian group. The algebraic condition discriminant $\Delta \neq 0$ geometrically imply the graph of E has no cusps, self-intersections, or isolated points. In addition, the real graph of a non-singular curve has two components if its discriminant is positive, and one component if it is negative. One can [analogously] assign the defining polynomial for an elliptic curve to be a bivariate Polynomial $[y^2 - x^3 - ax - b]$. This is a "mixed" Polynomial $P(x, y) = \text{Monomial } P(x) \text{ of degree } 3 + \text{Monomial } P(y) \text{ of degree } 2$.

The four basic arithmetic operations are addition, subtraction, multiplication and division. Product (viz, multiplication) of a sequence, denoted by \prod , can be finite $\prod_{i=1}^N$ or infinite $\prod_{i=1}^{\infty}$. This extended operation in multiplication should be differentiated from dot product, matrix multiplication, scalar multiplication, multiplication of vectors, etc. In our elaborate scheme of defining the "Generic Equation", infinite product of a sequence such as Euler product is NOT an Equation by itself, although it could mathematically form a Term, or part of a Term, in an Equation.

Euler product is defined as the expansion of a Dirichlet series into an infinite product indexed by prime numbers. In general, if a is a bounded multiplicative function, then the Dirichlet series $\sum_n \frac{a(n)}{n^s}$ is equal to $\prod_p P(p, s)$ for $\text{Re}(s) > 1$ where the product is taken over prime numbers p , and $P(p, s)$ is the sum $\sum_{k=0}^{\infty} \frac{a(p^k)}{p^{ks}} = 1 + \frac{a(p)}{p^s} + \frac{a(p^2)}{p^{2s}} + \frac{a(p^3)}{p^{3s}} + \dots$. In fact, if we consider these as formal generating functions, the existence of such a formal Euler product expansion is a necessary and sufficient condition that $a(n)$ be multiplicative: this exactly imply $a(n)$ is the product of the $a(p^k)$ whenever n factors as the product of the powers p^k of distinct primes p . An important special case is when $a(n)$ is totally multiplicative, so that $P(p, s)$ is a geometric series. Then $P(p, s) = \frac{1}{1 - \frac{a(p)}{p^s}}$ as is the case for Riemann zeta function, where $a(n) = 1$, and more generally for Dirichlet characters.

A Dirichlet series is a formal series of the form $F(s) = \sum_{n=1}^{\infty} \frac{a_n}{n^s}$ where $a_n \in \mathbb{C}$, $s = \sigma \pm it$ with $\sigma, t \in \mathbb{R}$, and $i = \sqrt{-1}$ is the imaginary unit. A Dirichlet L-function is an L-function defined by a Dirichlet series

of the form $L(s, \chi) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s}$, where χ is a Dirichlet character. Here, a Dirichlet character is a function $\chi : \mathbb{Z} \rightarrow \mathbb{C}$ together with a positive integer q called the modulus such that χ is completely multiplicative, i.e. $\chi(mn) = \chi(m)\chi(n)$ for all integers m and n , and χ is periodic modulo q , i.e. $\chi(n+q) = \chi(n)$ for all n . If $(n, q) > 1$ then $\chi(n) = 0$, whereas if $(n, q) = 1$, then $\chi(n)$ is a root of unity. The character χ is primitive if its conductor is equal to its modulus.

Remark 1.2. Definition and Axioms of all L-functions. An (analytic) L-function is a Dirichlet series that has an Euler product and satisfies a certain type of functional equation, and allows analytic continuation. Then this L-function is also called Dirichlet L-function, associated with its Dirichlet L-series, which can be meromorphically continued to the complex plane, have an Euler product $L(s, \chi) = \prod_p (1 - \chi(p)p^{-s})^{-1}$, and satisfy a functional equation of the form $\Lambda(s, \chi) = q^{\frac{s}{2}} \Gamma_{\mathbb{R}}(s) L(s, \chi) = \varepsilon_{\chi} \overline{\Lambda}(1-s)$, where q is the conductor of χ . Thus the three main defining features of L-functions are Axiom I: Analytic continuation [viz, technique to extend the domain of a given analytic function]. Axiom II: Euler product [over prime p as previously defined above]. Axiom III: Functional equation [created using gamma factors, see below].

The complex functions $\Gamma_{\mathbb{R}}(s) := \pi^{-\frac{s}{2}} \Gamma(\frac{s}{2})$ and $\Gamma_{\mathbb{C}}(s) := 2(2\pi)^{-s} \Gamma(s)$ that appear in functional equation of an L-function are known as gamma factors. Here $\Gamma(s) := \int_0^{\infty} e^{-t} t^{s-1} dt$ is Euler's gamma function. The gamma factors satisfy $\Gamma_{\mathbb{C}}(s) = \Gamma_{\mathbb{R}}(s) \Gamma_{\mathbb{R}}(s+1)$ and can also be viewed as "missing" factors of the Euler product of an L-function corresponding to (real or complex) archimedean places.

All known analytic L-functions have functional equations that can be written in the form

$\Lambda(s) := N^{\frac{s}{2}} \prod_{j=1}^J \Gamma_{\mathbb{R}}(s + \mu_j) \prod_{k=1}^K \Gamma_{\mathbb{C}}(s + \nu_k) \cdot L(s) = \varepsilon \overline{\Lambda}(1-s)$ where N is an integer, $\Gamma_{\mathbb{R}}$ and $\Gamma_{\mathbb{C}}$ are defined in terms of the Γ -function, $\text{Re}(\mu_j) = 0$ or 1 (assuming Selberg's eigenvalue conjecture), and $\text{Re}(\nu_k)$ is a positive integer or half-integer, $\sum \mu_j + 2 \sum \nu_k$ is real, and ε is the sign of the functional equation. With those restrictions on the spectral parameters, the data in the functional equation is specified uniquely. The integer $d = J + 2K$ is the degree of the L-function. The integer N is the conductor (or level) of the L-function. The pair $[J, K]$ is the signature of the L-function.

The functional equation for $\zeta(s)$ is $\zeta(s) = 2^s \pi^{s-1} \sin\left(\frac{\pi s}{2}\right) \Gamma(1-s) \zeta(1-s)$.

The functional equation for $\eta(s)$ is $\eta(-s) = 2 \frac{1-2^{-s-1}}{1-2^{-s}} \pi^{-s-1} s \sin\left(\frac{\pi s}{2}\right) \Gamma(s) \eta(s+1)$.

The Riemann zeta function $\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = \frac{1}{1^s} + \frac{1}{2^s} + \frac{1}{3^s} + \dots$, having convergence when $\Re(s) > 1$, is prototypical "non-alternating zeta function (harmonic series)" L-function. Riemann zeta function is the only L-function of degree 1 and conductor 1, and (conjecturally) it is the only primitive L-function with a unique pole [located at $s = 1$]; and is analytically continued to entire complex plane as Dirichlet eta function $\eta(s) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^s} = \frac{1}{1^s} - \frac{1}{2^s} + \frac{1}{3^s} - \frac{1}{4^s} + \dots$. The Dirichlet eta function, having convergence

when $\Re(s) > 0$, is prototypical "alternating zeta function (harmonic series)" L-function. This continuation is defined by relationship $\eta(s) = \gamma \cdot \zeta(s) = (1 - 2^{1-s}) \zeta(s)$ where $\gamma = 1 - 2^{1-s}$ is the proportionality factor.

Euler product expressions are possible for Riemann zeta function $\zeta(s)$ and its L-function $L_\zeta(s)$, Mobius function $\mu(n)$, Leibniz formula (for $\frac{\pi}{4}$), Sum-of-Divisors function $\sigma(n)$, Euler's totient function $\Phi(n)$, Elliptic curves $\zeta_E(s)$ with their associated L-functions $L_E(s)$ and modular forms, etc.

The Euler product expression for Riemann zeta function $\zeta(s)$ having convergence when $\Re(s) > 1$ is

$$\sum_{n=1}^{\infty} \frac{1}{n^s} = \prod_{p \text{ prime}} \frac{1}{1 - p^{-s}} = \frac{1}{1 - 2^{-s}} \cdot \frac{1}{1 - 3^{-s}} \cdot \frac{1}{1 - 5^{-s}} \cdot \frac{1}{1 - 7^{-s}} \cdot \frac{1}{1 - 11^{-s}} \cdots \frac{1}{1 - p^{-s}} \cdots$$

"Alternating" Euler product expression for Dirichlet eta function $\eta(s)$ having convergence when $\Re(s) > 0$

$$\text{is } \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^s} = \prod_{p \text{ prime}} \frac{1}{1 \pm p^{-s}} = \frac{1}{1 + 2^{-s}} \cdot \frac{1}{1 - 3^{-s}} \cdot \frac{1}{1 + 5^{-s}} \cdot \frac{1}{1 - 7^{-s}} \cdot \frac{1}{1 + 11^{-s}} \cdots \frac{1}{1 \pm p^{-s}} \cdots$$

Modular forms and the Modularity theorem. Having special spectacular properties resulting from surprising array of internal symmetries, modular forms describe several types of complex functions which have a certain type of functional equation and growth condition. The q -expansion of a modular form $f(z)$ is its Fourier expansion at the cusp $z = i\infty$, expressed as a power series $\sum_{n=0}^{\infty} a_n q^n$ in the variable $q = e^{2\pi iz}$.

As exemplified by Dedekind eta function $\eta(z) = q^{1/24} \prod_{n \geq 1} (1 - q^n)$ with $\prod_{n \geq 1} (1 - q^n)$ as basic construction block of infinite products, this implies that many types of infinite products with this construction block are modular [but, of course, not similar-looking "micmicker" products of the type $\prod (1 - q^{n^2})$ or $\prod (1 - q^n)^n$].

L-function associated with a modular form can often be expressed as an Euler product over primes. The fully proven 2001 Modularity theorem states all elliptic curves over the field of rational numbers are UNIQUELY related to modular forms in a particular way. We use notation $T = p^{-s} = \frac{1}{p^s}$ to indicate local polynomials of various degree. E.g., Sum-of-positive divisors function $\sigma_z(n)$ has local polynomial $F_p(T) = (1 - T)(1 - pT)$ of degree one. The functions $L_p(s)$ are called Euler factors (or local factors), and for $\sigma_z(n)$, this is $F_p(T)^{-1} = \frac{1}{F_p(T)} = \frac{1}{(1 - T)(1 - pT)}$. Local polynomials for elliptic curves are of degree 2.

Euler product of an L-function in details: It is expected that the Euler product of an L-function of

degree d and conductor N can be written as $L(s) = \prod_p L_p(s)$ where for $p \nmid N$ $L_p(s) = \prod_{n=1}^d \left(1 - \frac{\alpha_n(p)}{p^s}\right)^{-1}$

with $|\alpha_n(p)| = 1$ and for $p \mid N$, $L_p(s) = \prod_{n=1}^{d_p} \left(1 - \frac{\beta_n(p)}{p^s}\right)^{-1}$ where $d_p < d$ and $|\beta_n(p)| \leq 1$. Again, the functions $L_p(s)$ are called Euler factors (or local factors).

An eta quotient is any function f of the form $f(z) = \prod_{1 \leq i \leq s} \eta^{r_i}(m_i z)$, where $m_i \in \mathbb{N}$ and $r_i \in \mathbb{Z}$ and $\eta(z)$ is the Dedekind eta function. An eta product is an eta quotient in which all the r_i are non-negative. We define the Dedekind eta function $\eta(z)$ by the formula $\eta(z) = q^{1/24} \prod_{n \geq 1} (1 - q^n)$, where $q = e^{2\pi iz}$. The Dedekind eta function is a crucial example of a half-integral weight modular form, having weight $1/2$ and level 1. It is related to the Discriminant modular form via the formula $\Delta(z) = \eta^{24}(z)$.

All elliptic curves have UNIQUE local zeta functions, local L-functions and functional equations that can be equivalently expressed using Euler products, and fully satisfy Axioms I, II and III (Remark 1.2). Each elliptic curve over \mathbb{Q} has an integral Weierstrass model (or equation) of the form $y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6$, where a_1, a_2, a_3, a_4, a_6 are integers. Reducing these coefficients modulo p defines an elliptic curve over finite field \mathbb{F}_p (except for a finite number of primes p , where the reduced curve has a singularity and thus fails to be elliptic, in which case E is said to be of bad reduction at p). Each such equation, as unique minimal Weierstrass equation which satisfies the additional constraints $a_1, a_3 \in \{0, 1\}, a_2 \in \{-1, 0, 1\}$, has a discriminant Δ as nonzero integer divisible exactly by these "bad primes" p . Here, a minimal Weierstrass equation is one for which $|\Delta|$ is minimal among all Weierstrass models for the same curve. The zeta function of an elliptic curve over a finite field \mathbb{F}_p is given by $Z(E(\mathbb{F}_p), T) = \exp\left(\sum_{n=1}^{\infty} \# [E(\mathbb{F}_{p^n})] \frac{T^n}{n}\right)$.

This can also be given via a rational function in T as $Z(E(\mathbb{F}_p), T) = \frac{1 - a_p T + pT^2}{(1 - T)(1 - pT)}$, where the 'trace of Frobenius' term a_p is defined to be the difference between the 'expected' number $p + 1$ and the number of points on the elliptic curve E over \mathbb{F}_p , viz. $a_p = p + 1 - \#E(\mathbb{F}_p)$ or equivalently, $\#E(\mathbb{F}_p) = p + 1 - a_p$.

The L-function of E over \mathbb{Q} is then defined by collecting this information together, for all primes p and is defined by $L(E(\mathbb{Q}), s) = \prod_{p \nmid N} (1 - a_p p^{-s} + p^{1-2s})^{-1} \cdot \prod_{p \mid N} (1 - a_p p^{-s})^{-1}$ where N is the conductor of E ,

i.e. the product of primes with bad reduction. This product converges for $\Re(s) > \frac{3}{2}$ [which can be *analytically normalized* to converge for $\Re(s) > 1$ by using $\Gamma_{\mathbb{C}}(s + \frac{1}{2})$ instead of $\Gamma_{\mathbb{C}}(s)$ in deriving the functional equation]. Hasse's conjecture then affirms that the L-function admits an analytic continuation to the whole complex plane and satisfies a functional equation relating, for any s , $L(E, s)$ to $L(E, 2 - s)$.

This L-function, of a modular form whose analytic continuation is known, has valid values of $L(E, s)$ at any complex number s e.g. at $s = 1$ (where the conductor product can be discarded as it is finite), the L-function becomes $L(E(\mathbb{Q}), 1) = \prod_{p \nmid N} (1 - a_p p^{-1} + p^{-1})^{-1} = \prod_{p \nmid N} \frac{p}{p - a_p + 1} = \prod_{p \nmid N} \frac{p}{\#E(\mathbb{F}_p)}$.

The Analytic rank of an L-function $L(s)$ is its order of vanishing at its central point. When the Analytic rank r is positive, the value is typically an upper bound that is believed to be tight (in the sense that there are known to be r zeroes located very near to the central point). Examples of modular forms include classical modular forms, Maass waveforms, Hilbert modular forms, Bianchi modular forms, and Siegel modular forms. The Analytic rank of a cuspidal modular form f is actually the Analytic rank of the L-function $L(f, s) = \sum_{n \geq 1} a_n n^{-s}$ where a_n are complex coefficients that appear in the q -expansion of modular form: $f(z) = \sum_{n \geq 1} a_n q^n$, where $q = e^{2\pi iz}$. The complex coefficients a_n depend on a choice of embedding of the coefficient field of f into complex numbers. It is also conjectured the Analytic rank does not depend on this choice, and this conjecture has been verified for all classical modular forms. For modular forms, the Analytic ranks are provably correct whenever the listed Analytic rank is 0, or the listed Analytic rank is 1 and the modular form is self dual (in the self dual case the sign of functional equation determines the parity of the Analytic rank).

1.1. *Linking Generalized Riemann hypothesis to Birch and Swinnerton-Dyer conjecture. **Treatise on Polignac's and Twin prime conjectures that asserts the infinity nature of Odd Primes derived from each and every even Prime gaps 2, 4, 6, 8, 10... is supplied later on.*** Materials available in L-functions and modular forms database (LMFDB) are relevant to the Langlands program, Generalized Riemann hypothesis (GRH) [and its special case Riemann hypothesis (RH)], and Birch and Swinnerton-Dyer (BSD) conjecture. Appendix A contains additional supplementary materials, practical computations and **overall SUMMARY** on Predictability properties of Dirichlet L-series and Dirichlet L-functions relevant to open problems in Number theory. There are Completely Predictable infinitely-many trivial zeros and Incompletely Predictable infinitely-many nontrivial zeros in an L-function. We assert entire sets of unique Incompletely Predictable nontrivial zeros (spectrum) in all L-functions [\equiv GRH]; and in L-functions of Riemann zeta function / Dirichlet eta function [\equiv RH], must pass through Centroid (Origin) point whereby this 0-dimensional "geometric" point is equivalent to 1-dimensional "mathematical" Critical Line.

Supporting GRH and BSD conjecture to be true for all L-functions from elliptic curves, one could [analogically] derive computed **Central values at vertical line $s = 1$ in their functional equations** [relating $L(E, s)$ to $L(E, 2 - s)$ for any s ; see Remark 1.1], construct Polar graphs of $\zeta_E(\sigma \pm it)$ with all UNIQUE nontrivial zeros (as individualized E spectrum) being ONLY located at [**analytically normalized**] $\sigma = \frac{1}{2}$ -Critical Line for each elliptic curve, etc. Furthermore, validly applying *Principle of Equidistant for Multiplicative Inverse* (Remark 2.2) to elliptic functions and *Infinitesimal value $\frac{1}{\infty}$ at just above / below $\sigma = \frac{1}{2}$ -Origin point (Centroid point)* on graphed trajectories of elliptic curves for **analytically normalized $\zeta_E(s)$** [as similarly carried out for Riemann zeta function $\zeta(s)$ / Dirichlet eta function $\eta(s)$ e.g. in Figure 10] should also be possible for all elliptic curves that are defined for s over \mathbb{C} .

With complex variable $s = \sigma \pm it$, we further assert all L-functions with Analytic rank 0 [e.g. Dirichlet eta function (*proxy* function for Riemann zeta function) and those Elliptic curves having Analytic rank 0] DO NOT have first nontrivial zero located at $t = 0$ on actual / normalized $\sigma = \frac{1}{2}$ -Critical Line]. A caveat here is the L-function of Riemann zeta function, as exception, DO NOT have nontrivial zeros. The corollary is then true in that all L-functions with Analytic rank 1 or higher [e.g. Elliptic curves having Analytic ranks of 1, 2, 3, 4, 5... (to an arbitrarily large number value)] DO have first nontrivial zero located at $t = 0$ on the actual / normalized $\sigma = \frac{1}{2}$ -Critical Line. The [more complex] L-functions of elliptic curves having relatively higher Analytic ranks are expected to generally have higher frequency in appearances of nontrivial zeros. By the very definitions and constructions of Dirichlet L-series from Dirichlet L-functions [as various zeta functions], both mathematical objects must, by default, comply with three fundamental Axioms I, II and III of L-functions. We recognize this deduction allows us to derive $\zeta_E(s)$, $L_E(s)$, functional equations and (equivalent) Euler products expressions based on Dirichlet coefficients a_n obtained from associated $f(q)$ Modular forms. Intuitively, the mathematical objects of elliptic curves are "higher" analogues of MOST BASIC Riemann zeta function $\zeta(s)$ and its analytic continuation to Dirichlet eta function $\eta(s)$, simplified- $\eta(s)$ and Dirichlet Sigma-Power Law [= $\int \text{sim-}\eta(s)dn$], which can similarly be constructed for all elliptic curves. The first nontrivial zero of Dirichlet eta function [viz, *proxy* function

for Riemann zeta function] with Analytic rank 0, at height approximately 14.134, is higher than that of any other algebraic L-function. Then any other algebraic L-function [with Analytic rank 0, 1, 2, 3, 4, 5...] will always have comparatively more frequently occurring nontrivial zeros [that first occur at a relatively lower height [for L-functions with Analytic rank 0], up to and including the (lowest) height of 0 [for L-functions with Analytic rank 1 or higher].

Useful preliminary information on Generic Numbers, Generic Terms and Generic Equation with their definitions and classification role:

Hyperreal numbers extend real numbers to include certain classes of infinite and infinitesimal numbers. Surreal numbers is a totally ordered proper class containing the real numbers, infinite and infinitesimal numbers that are larger or smaller in absolute value than any positive real number. Quaternion number system extends the complex numbers. Quaternions have expression of the form $a + b\mathbf{i} + c\mathbf{j} + d\mathbf{k}$, where a, b, c, d are real numbers; $\mathbf{i}^2 = \mathbf{j}^2 = \mathbf{k}^2 = -1, \mathbf{i}\mathbf{j} = -\mathbf{j}\mathbf{i} = \mathbf{k}, \mathbf{j}\mathbf{k} = -\mathbf{k}\mathbf{j} = \mathbf{i}, \mathbf{k}\mathbf{i} = -\mathbf{i}\mathbf{k} = \mathbf{j}$. These higher or more abstract number systems, and the main number systems below, form the **Generic Numbers**.

Integer numbers $\mathbb{Z} \subset$ Rational numbers $\mathbb{Q} \subset$ Real numbers $\mathbb{R} \subset$ Complex numbers \mathbb{C} .

Natural numbers $\mathbb{N} \{1, 2, 3, 4, 5...\} \subset$ Whole numbers $\mathbb{W} \{0, 1, 2, 3, 4, 5...\} \subset$ Integer numbers $\mathbb{Z} \{...-3, -2, -1, 0, 1, 2, 3...\}$. The pairing of Even numbers $\mathbb{E} \{0, 2, 4, 6, 8, 10...\}$ and Odd numbers $\mathbb{O} \{1, 3, 5, 7, 9, 11...\}$, and the pairing of Prime numbers $\mathbb{P} \{2, 3, 5, 7, 11, 13, 17, 19, 23...\}$ and Composite numbers $\mathbb{C} \{4, 6, 8, 9, 10, 12, 14, 15, 16, 18, 20...\}$ can be separately combined to form \mathbb{W} whereby $\{0, 1\}$ are neither prime nor composite. Complex number $z = a + bi$ where imaginary unit $i = \sqrt{-1}; a, b \in \mathbb{R}$; and when $b = 0$, z becomes a real number. $\mathbb{Q} = \frac{p}{q}$ where $p, q \in \mathbb{Z}; \mathbb{Q}$ are \mathbb{Z} when $p = 1$; and $q = 0$ is undefined.

Irrational numbers $\mathbb{R} \setminus \mathbb{Q} \subset$ Real numbers \mathbb{R} or Complex numbers \mathbb{C} . Then $\mathbb{R} \setminus \mathbb{Q} =$ [I] Algebraic (irrational) numbers [viz, \mathbb{R} or \mathbb{C} that are the root of a non-zero polynomial of finite degree in one variable with integer or, equivalently, rational coefficients e.g. golden ratio $(1 + \sqrt{5})/2$, $\sqrt{2}$, $\sqrt[3]{2}$, etc] + [II] Transcendental (irrational) numbers [viz, \mathbb{R} or \mathbb{C} that are NOT the root of a non-zero polynomial of finite degree in one variable with integer or, equivalently, rational coefficients e.g. $\pi, e, \ln 2$]. The only even Prime number $\{2\}$ forms a Countably Finite Set (CFS). $\mathbb{E}, \mathbb{O}, \mathbb{P}, \mathbb{C}, \mathbb{N}, \mathbb{W}, \mathbb{Z}, \mathbb{Q}$ and Algebraic numbers form Countably Infinite Sets (CIS). Transcendental numbers, $\mathbb{R} \setminus \mathbb{Q}, \mathbb{R}$ and \mathbb{C} form Uncountably Infinite Sets (UIS).

We mathematically define the (finite) N terms in Generic Equation as: Generic Term $T_1 +$ Generic Term $T_2 +$ Generic Term $T_3 +$ Generic Term $T_4 + \dots$ Generic Term $T_{N-1} +$ Generic Term $T_N = 0$. The finite Generic Equation with finite $[n \rightarrow N]$ terms is $\sum_{n=1}^N T_n = 0$, and infinite Generic Equation with infinitely

many $[n \rightarrow \infty]$ terms is $\sum_{n=1}^{\infty} T_n = 0$. In particular, infinite series such as modular forms, power series and harmonic series are simply various types of Equations. **Importantly, Euler products [infinite-like expressions], when used for expansions of Dirichlet series resulting in infinite products [viz, NOT infinite sums] indexed by prime numbers, are NOT Equations *per se* but could, in principle, form a Term(s) or part of a Term(s) in an Equation.** The solutions or zeros (also sometimes called roots) of a **real-, complex-**

or generally **vector**-valued function f in a Generic Equation are members x of domain of f such that $f(x)$ vanishes at x . Observe the n index for Generic Equation to commence from 0 or 1 is an arbitrary choice that is mathematically valid as long as consistency is maintained (viz, standardized).

T_n , the n^{th} Term, is defined by either algebraic functions or non-algebraic (irrational or transcendental) functions; or a mixture of the two. Examples of [univariate or 1-variable] Terms involving x as variable (or indeterminate): $a_n x^n$, $a x^{-n} \equiv \frac{a}{x^n}$ [involving algebraic functions]; $a \sqrt[n]{x} \equiv a x^{\frac{1}{n}}$, $a \sin^n x$, e^{nx} , $a \log_n x$ [involving irrational or transcendental functions]; $a_n x^n \sin^n x$ [involving mixed functions]. Notations: a = coefficient, n = exponent [or base for logarithm function, whereby when $n = e$, $a \log_e x = a \ln x$ involves the natural logarithm]. The "generic" Infinite series are of two types: Convergent series or Divergent series. In general, non-alternating harmonic series are usually Divergent series but alternating harmonic series are usually Convergent series.

With infinitely many unit fractions as Terms [Infinite series], Harmonic series have T_n of the type $\frac{a}{x^n}$ e.g. alternating and non-alternating harmonic series, Riemann zeta function, Dirichlet eta function. Harmonic series could also involve more than one variables with coefficients a_n that could theoretically also be more complex and depend on variable x , etc. Egyptian fractions, being the finite sum of distinct unit fractions, are roughly like Finite harmonic series. Note 1: Not related to harmonic series *per se* is the Harmonic functions giving rise to non-polynomial (transcendental) terms e.g. second derivatives $e^x \sin y$ and $-e^x \sin y$ of a 2-variable harmonic function; viz, twice continuously differentiable function $f: U \rightarrow \mathbb{R}$ where U is an open subset of \mathbb{R}^n that satisfies this particular Laplace's equation $f(x, y) = e^x \sin y$. Note 2: **With the 1-variable now given here by $s [= \sigma \pm it]$ in various L -functions $L(s)$ as harmonic series associated with its L -series** [see Remark 1.3], s can now be complex numbers, or be positive / negative real numbers given by σ when $t = 0$.

With infinitely many Terms [Infinite series] that all have exponents (powers), Power series have T_n of the type $a_n x^n$ e.g. alternating and non-alternating power series, geometric series, Taylor series, Maclaurin series, exponential function formula, sine formula, etc. Power series can also involve more than one variables. Having n as negative powers and fractional powers give rise to variants of power series called Laurent series and Puiseux series. Formal power series capture the essence of power series without being restricted to the fields of real and complex numbers, and without the need to talk about convergence.

With finitely many Terms [Finite series] that all have exponents (powers), Polynomials have T_n of the type $a_n x^n$ e.g. integer polynomial, real polynomial, complex polynomial [as defined by their coefficients derived from various number systems]; rational fraction [being the quotient (algebraic fraction) of two polynomials], exponential polynomials [a bivariate polynomial where the second variable is substituted for an exponential function applied to the first variable such as $P(x, e^x)$], matrix polynomial [a polynomial with square matrices as variables], trigonometric polynomial [a finite linear combination of functions $\sin(nx)$ and $\cos(nx)$ with n taking on the values of one or more natural numbers, and having real- or complex-valued coefficients]. Polynomial is formally defined as a mathematical expression consisting of indeterminates (variables) and coefficients, that involves only the operations of addition, subtraction,

multiplication and exponentiation to nonnegative integer powers, and has a finite number of terms [viz, monomial with 1 term, binomial with 2 terms, polynomial with ≥ 2 terms (whereby each term can also be usefully labelled as a monomial)]. Polynomials can also involve more than one variables, as multivariate polynomial. Polynomials are roughly like "finite power series" [having finite degree], or equivalently, Power series are roughly like "infinite polynomials" [having infinite degree]. Laurent polynomials are like polynomials, but allow negative powers of the variable(s) to occur.

A number field can be defined by many different irreducible polynomials $f(x) \in \mathbb{Q}[x]$. The defining polynomial of a number field K is an irreducible polynomial $f \in \mathbb{Q}[x]$ such that $K \cong \mathbb{Q}(a)$, where a is a root of $f(x)$. A root $a \in K$ of the defining polynomial is a generator of K . Normalized polynomials are always monic with integer coefficients, such that the sum of the squares of the absolute values of all complex roots of $f(x)$ is minimized. In comparison, the defining polynomial of a p -adic field K is an irreducible polynomial $f(x) \in \mathbb{Q}_p[x]$ such that $K \cong \mathbb{Q}_p(a)$, where a is a root of $f(x)$. The defining polynomial can be chosen to be monic with coefficients in \mathbb{Z}_p ; by Krasner's lemma, we can further take $f(x) \in \mathbb{Z}[x]$.

Remark 1.3. Normalized defining polynomial for L -functions and Number fields:

Integer '0' as Equation $0 = 0$ is the zero polynomial $P(x) = 0$ of undefined degree arbitrarily assigned as either -1 or $-\infty$, with all coefficients = 0. It is the additive identity in set of polynomials; viz, $P(x) + 0 = P(x)$. Any nonzero integers c e.g. $-3, -2, -1, 1, 2, 3...$ as Equation $c = c$ is the constant polynomial $P(x) = c$ of degree 0 [since it can be written as $c \cdot x^0$], with coefficient of $x^0 = c$ & all other coefficients = 0.

Equation " P " $x = 0 \equiv x + 0 = 0$ is the [linear] defining polynomial $P(x) = x$ of degree 1 [since it can be written as $1 \cdot x^1$], with coefficient of $x^1 = 1$ & all other coefficients = 0. **It conceptually represents the "most basic" 1-variable infinite non-alternating harmonic series, an unique L -function $L_\zeta(s)$, called Riemann zeta function $\zeta(s)$ having infinitely many terms.**

Equation " D " $\pm x = 0 \equiv \pm x + 0 = 0$ is [linear] defining polynomial $P(x) = \pm x$ of degree 1 [since it can be written as $\pm 1 \cdot x^1$], with coefficient of $x^1 = \pm 1$ & all other coefficients = 0. Conveniently, it **conceptually represents the "most basic" 1-variable infinite alternating harmonic series, an unique L -function $L_\eta(s)$, called Dirichlet eta function $\eta(s)$ having infinitely many terms.**

Equation " K " $x = \pm \sqrt{-1} = \pm i \equiv x^2 + 1 = 0$ is [non-linear] defining polynomial $P(x) = x^2 + 1$ of degree 2, with coefficient of $x^2 = 1$, first coefficient = 1 & all other coefficients = 0, that connects Real numbers \mathbb{R} to Complex numbers \mathbb{C} and **conceptually represents an important 1-variable infinite non-alternating harmonic series, as an unique L -function $L_K(s)$ containing the field of Gaussian rational numbers, and is associated with Automorphic object " A " given by L -function $L_A(s)$. Both $L_K(s)$ and $L_A(s)$ have infinitely many terms.** Additional information: The ring of integers, $\mathbb{Z}[i]$, is a Euclidean domain, hence unique factorization domain, with norm $N(a + bi) = a^2 + b^2 = (a + bi)(a - bi)$. As a result, it is connected to the question of which positive integers can be written as the sum of two squares, and more specifically,

to the theorem of Fermat that a prime number p can be written as the sum of two squares if and only if $p \not\equiv 3 \pmod{4}$, and that if $p = a^2 + b^2$, then the representation is unique subject to $0 < a \leq b$.

Equation "E" $y^2 = x^3 + ax + b \equiv y^2 - x^3 - ax - b = 0$ is the [non-linear] 2-variable (bivariate) defining polynomial $P(x, y) = P(x)$ of degree 3 + $P(y)$ of degree 2 = $y^2 - x^3 - ax - b$ with coefficients 1, -1, -a, -b & all other coefficients = 0. It **conceptually give rise to important 1-variable infinite alternating harmonic series, as unique L -functions $L_E(s)$ associated with Modular forms** that act as (periodic) 'generating series or functions' based on elliptic functions / curves. Both $L_E(s)$ and Modular forms have infinitely many terms.

Riemann zeta function $\zeta(s)$ vs Dirichlet eta function $\eta(s)$: $\zeta(s)$ is the **1-variable prototypical non-alternating L -function**. Analytic continuation of $\zeta(s)$ and its L -function $L_\zeta(s)$ [with Convergence for complex number when $\text{Re}(s) > 1$] to proxy function $\eta(s)$ and its L -function $L_\eta(s)$ [with Convergence for any complex number when $\text{Re}(s) > 0$] is required to obtain nontrivial zeros. They are related via proportionality factor $\gamma = (1 - 2^{1-s})$ as $\eta(s) = \gamma \cdot \zeta(s)$. From Remark 1.3, we reiterate that $\eta(s)$ and $L_\eta(s)$ is the **1-variable prototypical alternating L -function** having infinitely many terms.

Infinitely-many Completely Predictable Trivial zeros of $\zeta(s)$ occurs at $s = -2, -4, -6, -8, -10, \dots$

Infinitely-many Completely Predictable Trivial zeros of $\eta(s)$ occurs at $s = 0, -2, -4, -6, -8, -10, \dots$

Nontrivial zeros of $\zeta(s)$ DO NOT exist. Note that $\eta(s)$ has "one extra" trivial zero at $s = 0$.

Infinitely-many Incompletely Predictable Nontrivial zeros in $\eta(s)$ occurs at Critical Line as defined by $\eta(s) = \frac{1}{2} \pm it$ with $\pm t$ values $\approx \pm 14.13, \pm 21.02, \pm 25.01, \pm 30.42, \pm 32.93, \pm 37.58, \dots$

Particular values from Dirichlet eta function $\eta(s)$:

In general, the k^{th} derivative of $\eta(s)$; viz, $\eta'(s), \eta''(s), \eta'''(s)$, etc with convergence for $\Re(s) > 0$ can be expressed as $\eta^{(k)}(s) = (-1)^k \sum_{n=1}^{\infty} \frac{(-1)^{n-1} (\ln n)^k}{n^s}$ for $k = 1, 2, 3, \dots$. The derivative with respect to parameter s for $s \neq 1$: $\eta'(s) = \sum_{n=1}^{\infty} \frac{(-1)^n \ln n}{n^s} = 2^{1-s} \ln(2) \zeta(s) + (1 - 2^{1-s}) \zeta'(s)$. Then, $\eta'(1) = \ln(2) \gamma - \ln(2)^2 2^{-1}$.

$\eta(0) = \frac{1}{2}$ [the Abel sum of Grandi's series $1 - 1 + 1 - 1 + \dots$]; $\eta(-1) = \frac{1}{4}$ [the Abel sum of $1 - 2 + 3 - 4 + \dots$].

For k an integer > 1 , if B_k is the k^{th} Bernoulli number then $\eta(1 - k) = \frac{2^k - 1}{k} B_k$. Also, $\eta(1) = \ln 2$ as an alternating harmonic series. $\eta(2) = \frac{\pi^2}{12}$, $\eta(4) = \frac{7\pi^4}{720} \approx 0.94703283$, $\eta(6) = \frac{31\pi^6}{30240} \approx 0.98555109$, $\eta(8) = \frac{127\pi^8}{1209600} \approx 0.99623300$, $\eta(10) = \frac{73\pi^{10}}{6842880} \approx 0.99903951$, $\eta(12) = \frac{1414477\pi^{12}}{1307674368000} \approx 0.99975769$.

The general form for even positive integers is: $\eta(2n) = (-1)^{n+1} \frac{B_{2n} \pi^{2n} (2^{2n-1} - 1)}{(2n)!}$. As $n \rightarrow \infty$, we obtain $\eta(\infty) = 1$. Via its functional equation, $\eta(s)$ has Completely Predictable infinitely many trivial zeros at each even negative integer $s = -2n$ for $n = 0, 1, 2, 3, 4, 5, \dots$; and also have infinitely-many nontrivial zeros.

Particular values from Riemann zeta function $\zeta(s)$:

In general, the k^{th} derivative of $\zeta(s)$; viz, $\zeta'(s)$, $\zeta''(s)$, $\zeta'''(s)$, etc with convergence for $\Re(s) > 1$ can be expressed as $\zeta^{(k)}(s) = (-1)^k \sum_{n=1}^{\infty} \frac{(\ln n)^k}{n^s}$ for $k = 1, 2, 3, \dots$. Derivative of $\zeta(s)$ when $s = 0$, for example, is $\zeta'(0) = -\frac{1}{2} \ln(2\pi)$. When s is considered for (purely) real number values: $\zeta(0) = -\frac{1}{2}$, $\zeta(\frac{1}{2}) = -1.4603545\dots$, etc. Taking the limit $s \rightarrow +\infty$ through the real numbers, one obtains $\zeta(+\infty) = 1$. But at complex infinity on the Riemann sphere the zeta function has an essential singularity.

For any positive even integer $2n$, $\zeta(2n) = \frac{|B_{2n}|(2\pi)^{2n}}{2(2n)!}$, where B_{2n} is the $(2n)^{\text{th}}$ Bernoulli number. For odd positive integers, no such simple expression is known, although these values are thought to be related to the algebraic K -theory of the integers.

For nonpositive integers, one has $\zeta(-n) = -\frac{B_{n+1}}{n+1}$ for $n \geq 0$ (using the convention that $B_1 = \frac{1}{2}$).

In particular, $\zeta(s)$ vanishes at negative even integers because $B_m = 0$ for all odd m other than 1. These are the trivial zeros. The point $s = 1$ in $\zeta(s)$ corresponds to a simple pole with complex residue 1. Even though $\zeta(1)$ is undefined as it diverges to ∞ , its Cauchy principal value $\lim_{\varepsilon \rightarrow 0} \frac{\zeta(1 + \varepsilon) + \zeta(1 - \varepsilon)}{2}$ exists and is equal to Euler-Mascheroni constant $\gamma = 0.577218\dots$ [a transcendental number].

For special values of $\zeta(s)$ [as an L -function] for nonpositive integers [and where B_{n+1} is the $(n+1)^{\text{th}}$ Bernoulli number], one has $\zeta(-n) = -\frac{B_{n+1}}{n+1}$ for $n \geq 0$ (using the convention that $B_1 = \frac{1}{2}$). In particular, $\zeta(s)$ vanishes at the negative even integers because $B_m = 0$ for all odd m other than 1. Then $\zeta(-1) = -\frac{1}{12}$, $\zeta(-3) = -\frac{1}{120}$, $\zeta(-5) = -\frac{1}{252}$, ..., $\zeta(-11) = -\frac{691}{32760}$, $\zeta(-13) = -\frac{1}{12}$, For other special values

of $\zeta(s)$ [as an L -function] involving small positive integer values of s : $\zeta(1) = \infty$, $\zeta(2) = \frac{\pi^2}{6}$, $\zeta(3) = 1.2020569032\dots$, $\zeta(4) = \frac{\pi^4}{90}$, $\zeta(5) = 1.0369277551\dots$, $\zeta(6) = \frac{\pi^6}{945}$, $\zeta(7) = 1.0083492774\dots$, $\zeta(8) = \frac{\pi^8}{9450}$, $\zeta(9) = 1.0020083928\dots$, $\zeta(10) = \frac{\pi^{10}}{93555}$, etc. When $s = 2, 4, 6, 8, 10\dots$; computed $\zeta(s)$ values all contain transcendental irrational number π . When $s = 3, 5, 7, 9, 11\dots$; computed $\zeta(s)$ values are

"likely" all algebraic irrational numbers. In fact, only $\zeta(3)$ or Apéry's constant is proven to be an irrational number but it is unknown whether it is also a transcendental number derived from (e.g.) π^3 or another unrelated transcendental number. Here $\zeta(3) = \frac{7}{180}\pi^3 - 2 \sum_{k=1}^{\infty} \frac{1}{k^3(e^{2\pi k} - 1)}$ [as series representation found

by Ramanujan] and $\zeta(3) = \int_0^1 \int_0^1 \int_0^1 \frac{1}{1-xyz} dx dy dz$, [as integrand of the known triple integral for $\zeta(3)$]. Despite these unknowns, the computed $\zeta(s)$ solutions from substituting $s =$ even numbers 2, 4, 6, 8, 10... versus $s =$ odd numbers 3, 5, 7, 9, 11... should all be irrational numbers that are, crucially, *mutually exclusive* and **mathematically, geometrically and topologically** different from each other.

Via its functional equation, $\zeta(s)$ has Completely Predictable infinitely many trivial zeros at each even negative integer $s = -2n$ for $n = 1, 2, 3, 4, 5\dots$; but DO NOT have any nontrivial zeros.

Via its functional equation, $\zeta(s)$ has Completely Predictable infinitely many trivial zeros at each even negative integer $s = -2n$ for $n = 1, 2, 3, 4, 5\dots$; but DO NOT have any nontrivial zeros.

More information on Elliptic curves E over \mathbb{Q} of conductor N : E has good reduction at all primes p not dividing N , has multiplicative reduction at the primes p that exactly divide N (i.e. such that p divides N , but p^2 does not, with this written as $p||N$), and has additive reduction elsewhere (i.e. at the primes

where p^2 divides N). Then Hasse-Weil zeta function of E is of the form $Z_{V,\mathbb{Q}}(s) = \frac{\zeta(s)\zeta(s-1)}{L(E,s)}$. Here, $\zeta(s)$ is the usual Riemann zeta function and $L(E,s)$ is called the L -function of E/\mathbb{Q} , which takes the form

$$L(E,s) = \prod_p L_p(E,s)^{-1} \text{ where, for a given prime } p, L_p(E,s) = \begin{cases} (1 - a_p p^{-s} + p^{1-2s}), & \text{if } p \nmid N \\ (1 - a_p p^{-s}), & \text{if } p \mid N \text{ and } p^2 \nmid N \\ 1, & \text{if } p^2 \mid N \end{cases}$$

where in the case of good reduction a_p is $p + 1 -$ (number of points of $E \bmod p$), and in the case of multiplicative reduction a_p is ± 1 depending on whether E has split (plus sign) or non-split (minus sign) multiplicative reduction at p . A multiplicative reduction of curve E by the prime p is said to be split if $-c_6$ is a square in the finite field with p elements. There is a useful relation not using the conductor:

1. If p doesn't divide Δ (where Δ is the discriminant of the elliptic curve) then E has good reduction at p .
2. If p divides Δ but not c_4 then E has multiplicative bad reduction at p .
3. If p divides both Δ and c_4 then E has additive bad reduction at p .

Duality: In mathematics, a duality translates concepts, theorems or mathematical structures into other concepts, theorems or structures in a one-to-one fashion, often (but not always) by means of an involution operation: if the dual of A is B , then the dual of B is A . Such involutions sometimes have fixed points, so that the dual of A is A itself. Any vector space V has a corresponding dual vector space consisting of all linear forms on V together with the vector space structure of pointwise addition and scalar multiplication by constants. In any finite group, the number of nonisomorphic irreducible representations over the complex numbers is precisely the number of conjugacy classes. A **ket** is of the form $|v\rangle$ whereby it mathematically denotes a vector v in an abstract (complex) vector space V and physically represents a state of some quantum system; and a **bra** is of the form $\langle f|$ whereby it mathematically denotes a linear form $f : V \rightarrow \mathbb{C}$, i.e. a linear map that maps each vector in V to a number in the complex plane \mathbb{C} . Then letting the linear functional $\langle f|$ act on a vector $|v\rangle$ is written as $\langle f|v\rangle \in \mathbb{C}$. L -functions are fundamental mathematical objects in Number theory that are dual to prime numbers; viz, each L -function can be viewed as a vector in certain Hilbert space, and each prime can then be viewed as a vector in the *dual* Hilbert space.

From above discussion, the ubiquitous deep theme of duality exist in Linear algebra [viz, vector space $V \rightleftharpoons$ dual vector space V^* having elements called functionals], Quantum mechanics [viz, **bra** \rightleftharpoons **ket**], Group theory [conjugacy class \rightleftharpoons irreducible representations] and Number theory [viz, prime numbers \rightleftharpoons L -functions]. In particular, the duality present in Number theory is inevitably connected to Theory of Symmetry from Langlands program whereby various power series and harmonic series, L -series, Dirichlet series, Dirichlet eta function (proxy function for Riemann zeta function as the generating function for all nontrivial zeros), Sieve of Eratosthenes (as the generating algorithm for all prime numbers), etc are usefully regarded as variants of *infinite series*.

The complex number $z = a + bi$. Its real part a and imaginary part b are real numbers. Its imaginary unit i satisfy power-series expansions $\sum_{n=0}^{\infty} i^n$ [as well as basic facts about powers of i] with given terms:

$$i^0 = 1, \quad i^1 = i, \quad i^2 = -1, \quad i^3 = -i,$$

$$i^4 = 1, \quad i^5 = i, \quad i^6 = -1, \quad i^7 = -i$$

$$\vdots \quad \quad \quad \vdots \quad \quad \quad \vdots \quad \quad \quad \vdots$$

Using power-series definition, we prove Euler's formula for real values of x :

$$\begin{aligned} e^{ix} &= 1 + ix + \frac{(ix)^2}{2!} + \frac{(ix)^3}{3!} + \frac{(ix)^4}{4!} + \frac{(ix)^5}{5!} + \frac{(ix)^6}{6!} + \frac{(ix)^7}{7!} + \frac{(ix)^8}{8!} + \dots \\ &= 1 + ix - \frac{x^2}{2!} - \frac{ix^3}{3!} + \frac{x^4}{4!} + \frac{ix^5}{5!} - \frac{x^6}{6!} - \frac{ix^7}{7!} + \frac{x^8}{8!} + \dots \\ &= 1 \left(\frac{x^0}{0!} - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \dots \right) + i \left(\frac{x^1}{1!} - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots \right) \\ &= \cos x + i \sin x. \text{ [Note that when } x = \pi, e^{i\pi} = -1 \text{ (Euler's identity).]} \end{aligned}$$

In the last step above we recognize $\frac{x^0}{0!} = 1$ and the two terms are Maclaurin series [**alternating power series** or, broadly, alternating infinite series] for $\cos x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!}$ and $\sin x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!}$ with the rearrangement of terms justified because each series is absolutely convergent. Recall that $\cos x$ & $\cosh x$ are even functions, so $\cos(-x) = \cos(x)$ & $\cosh(-x) = \cosh(x)$; and $\sin x$ & $\sinh x$ are odd functions, so $\sin(-x) = -\sin(x)$ & $\sinh(-x) = -\sinh(x)$.

$$\sinh i = \frac{e^i - e^{-i}}{2} = i \sin 1, \cosh i = \frac{e^i + e^{-i}}{2} = \cos 1 \quad \& \quad \tanh i = \frac{\sinh i}{\cosh i} = \frac{(e^i - e^{-i})}{(e^i + e^{-i})} = i \tan 1.$$

$$\sin i = i \frac{e^1 - e^{-1}}{2} = i \sinh 1, \cos i = \frac{e^1 + e^{-1}}{2} = \cosh 1 \quad \& \quad \tan i = \frac{\sin i}{\cos i} = \frac{i(e^1 - e^{-1})}{(e^1 + e^{-1})} = i \tanh 1.$$

$$\cos i = \sum_{n=0}^{\infty} \frac{1}{(2n)!} = \frac{1}{0!} + \frac{1}{2!} + \frac{1}{4!} + \frac{1}{6!} + \frac{1}{8!} + \dots \quad \& \quad \sin i = i \sum_{n=0}^{\infty} \frac{1}{(2n+1)!} = i \left(\frac{1}{1!} + \frac{1}{3!} + \frac{1}{5!} + \frac{1}{7!} + \dots \right)$$

[Note: For $n = 0$ to ∞ , $(i)^{2n} = (i^2)^n = (-1)^n$].

Euler's formula produces following analytical identities for sine, cosine and tangent in terms of e and i :

$$\sin x = \frac{e^{ix} - e^{-ix}}{2i}, \cos x = \frac{e^{ix} + e^{-ix}}{2} \quad \& \quad \tan x = \frac{\sin x}{\cos x} = \frac{(e^{ix} - e^{-ix})}{i(e^{ix} + e^{-ix})}.$$

The related or extended Lindemann-Weierstrass theorem, Gelfond-Schneider theorem, Baker's theorem, four exponentials conjecture or Schanuel's conjecture could be used to establish transcendence of a large class of numbers constituted from the (algebraic) irrational numbers, transcendental (irrational) numbers and rational numbers. Natural logarithm of any natural number other than 0 and 1 (more generally, of any positive algebraic number other than 1) e.g. $\ln 2$ and $\ln \sqrt{2} = \ln 2^{\frac{1}{2}} = \frac{1}{2} \ln 2$ are transcendental numbers by the Lindemann-Weierstrass theorem. By the Gelfond-Schneider theorem, e^π [Gelfond's constant], $2^{\sqrt{2}}$ [Gelfond-Schneider constant as an example of a^b where a is algebraic but not 0 or 1, and b is (algebraic) irrational number], $e^{-\frac{\pi}{2}} = i^i$, etc are all transcendental numbers.

As sum of infinite (power) series, Euler's number $e = \sum_{n=0}^{\infty} \frac{1}{(n)!} = \sum_{n=1}^{\infty} \frac{1}{(n-1)!} = 1 + \frac{1}{1} + \frac{1}{1 \cdot 2} + \frac{1}{1 \cdot 2 \cdot 3} + \dots$
 $\cong 2.71828$ is the limit $\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n} \right)^n$. It can be characterized using integral $\int_1^e \frac{dx}{x} = 1$. As sum of

infinite series, $\ln 2 = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} = \frac{1}{1} - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots \approx 0.693147$. This infinite series can also be

expressed using Riemann zeta function as $\sum_{n=1}^{\infty} \frac{1}{n} [\zeta(2n) - 1] = \ln 2$. Some explicit formulas for $\ln 2$ as

a result of integration include $\int_0^1 \frac{dx}{1+x} = \int_1^2 \frac{dx}{x} = \ln 2$, $\int_0^{\infty} e^{-x} \frac{1-e^{-x}}{x} dx = \ln 2$, $\int_0^{\infty} 2^{-x} dx = \frac{1}{\ln 2}$,

$\int_0^{\frac{\pi}{3}} \tan x dx = 2 \int_0^{\frac{\pi}{4}} \tan x dx = \ln 2$, $-\frac{1}{\pi i} \int_0^{\infty} \frac{\ln x \ln \ln x}{(x+1)^2} dx = \ln 2$. In principal branch of logarithm, $\ln(-1) = 0 + i\pi = i\pi$. The analytic identity using natural logarithm $-\ln(1-i)$ is analogous to Euler's formula for

chosen transcendental (real) number values as based on inverse functions $\ln i = \ln(e^{i\frac{\pi}{2}}) = 0 + \frac{\pi}{2}i = 1.57079632679i$ & $e^i = \cos(1) + i \sin(1) = 0.540302306 + 0.841470985i$. It conforms to *Langlands program's Theory of Symmetry* w.r.t. imaginary number (point) $i = \sqrt{-1} = 0 + i = \cos(\frac{\pi}{2}) + i \sin(\frac{\pi}{2})$; viz,

$\ln(e^i) = i$ & $e^{(\ln i)} = i$ [c.f. Figure 8 manifesting (perfect) diagonal symmetry via $\ln(e^x) = x$ & $e^{(\ln x)} = x$]. Then $-\ln(1-i) = -\ln \sqrt{2} + i\frac{\pi}{4}$

$$= 0 + i + \frac{(i)^2}{2} + \frac{(i)^3}{3} + \frac{(i)^4}{4} + \frac{(i)^5}{5} + \frac{(i)^6}{6} + \frac{(i)^7}{7} + \frac{(i)^8}{8} + \dots$$

$$= 0 + \frac{i}{1} - \frac{1}{2} - \frac{i}{3} + \frac{1}{4} + \frac{i}{5} - \frac{1}{6} - \frac{i}{7} + \frac{1}{8} \dots$$

$$= 1 \left(0 - \frac{1}{2} + \frac{1}{4} - \frac{1}{6} + \frac{1}{8} - \dots \right) + i \left(\frac{1}{1} - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots \right)$$

Transcendental numbers $-\ln \sqrt{2} = -\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{2n} \approx -0.3465\dots$ and $\frac{\pi}{4} = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{2n-1} \approx 0.7853\dots$ [aka

Leibniz formula for π] as two **alternating power series** [or, broadly, alternating infinite series] are recognizably related to each other as they represent the two terms in the last step above. As expected, our

additive identity 0 in $-\ln(1-i)$ is analogous to **multiplicative identity 1** [viz, $\frac{x^0}{0!}$] in Euler's formula e^{ix} .

A formal series is an infinite series (sum) that is considered independently from any notion of convergence, and is manipulated with usual algebraic operations on series such as addition, subtraction, multiplication, division, partial sums, etc. A power series defines a function by taking numerical values for the variable WITHIN a radius of convergence. In contrast with NO requirements of convergence, a *formal* power series is a special kind of formal series whose terms are of the form ax^n where x^n is the n^{th} power of a variable x (n is a non-negative integer), and a is called the coefficient.

Not actually regarded as a function *per se* with its "variable" remaining an indeterminate, a generating function (or series) is a representation of infinite sequences of numbers as coefficients of a formal power series. More generally, a formal power series can include series with any finite (or countable) number of variables, and with coefficients in an arbitrary ring. Rings of formal power series are complete local rings, and this allows using calculus-like methods in the purely algebraic framework of algebraic geometry and commutative algebra. They are analogous in many ways to p -adic integers which can be defined as formal series of the powers of p (see Page 22 – 23 of [8]). Various types of generating functions include ordinary generating functions, exponential generating functions, Lambert series, Bell series, and Dirichlet series. Sieve of Eratosthenes (as generating algorithm for all prime numbers) and Dirichlet eta function (the *proxy* function for Riemann zeta function as generating function for all nontrivial zeros) are infinite series since

they both encapsulate "infinite sequences of numbers". In this sense, *generating functions and generating algorithms are literally synonymous with infinite series*. By the same token, harmonic series formed by summing all positive [or alternating positive and negative] unit fractions, are infinite series and can thus also be conveniently regarded as generating functions.

Remark 1.4. L -functions, which can also be denoted by \mathbb{L} [e.g. Dedekind zeta function, Riemann zeta function $\zeta(s)$, Dirichlet eta function $\eta(s)$, Dirichlet L -functions, Hecke L -functions, Artin L -functions, automorphic L -functions, elliptic functions, etc] are meromorphic functions on the complex plane, associated to one out of several categories of mathematical objects [viz, anything that has been or could be formally defined, and with which one may do deductive reasoning and mathematical proofs] e.g. Dirichlet character, Hecke character, Artin representations of Galois group G , modular form, λ -ring, Hilbert space, dual vector space, elliptic curve E (abelian variety / group) defined over field K (which can be general field, finite fields, quadratic field $\mathbb{Q}\sqrt{d}$ with d a square-free integer, field of p -adic numbers \mathbb{Q}_p , rational numbers \mathbb{Q} , real numbers \mathbb{R} or complex numbers \mathbb{C}), etc.

A 'general' Dirichlet series is an infinite series of the form $\sum_{n=1}^{\infty} a_n e^{-\lambda_n s}$ where a_n, s are complex numbers and $\{\lambda_n\}$ is a strictly increasing sequence of nonnegative real numbers that tends to infinity. An 'ordinary' Dirichlet series $\sum_{n=1}^{\infty} \frac{a_n}{n^s}$ is obtained by substituting $\lambda_n = \ln n$ while a power series $\sum_{n=1}^{\infty} a_n (e^{-s})^n$ is obtained when $\lambda_n = n$. **Riemann zeta function $\zeta(s)$ as non-alternating harmonic series Eq. (1) is the **most basic** 'ordinary' Dirichlet series with complex sequence $a_n = 1$ for $n = 1$ to ∞ **. Hurwitz zeta function is one of many zeta functions formally defined for complex variables s with $\text{Re}(s) > 1$ and $a \neq 0, -1, -2, -3, \dots$ by $\zeta(s, a) = \sum_{n=0}^{\infty} \frac{1}{(n+a)^s}$. This series is absolutely convergent for given values of s & a , and can be extended to a meromorphic function defined for all $s \neq 1$. **Riemann zeta function is then $\zeta(s, 1)$ **.

In more details: Dirichlet L -series is a function of the form $L(s, \chi) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s}$ where χ is a Dirichlet character and s a complex variable with $\text{Re}(s) > 1$. It is a special case of a Dirichlet series. By analytic continuation, it can be extended to a meromorphic function on whole complex plane, and is then called Dirichlet L -function and also denoted $L(s, \chi)$. Since Dirichlet character χ is completely multiplicative, its L -function can also be written as an Euler product in the half-plane of absolute convergence $L(s, \chi) = \prod_p (1 - \chi(p)p^{-s})^{-1}$ for $\text{Re}(s) > 1$ where the product is over all prime numbers. Dirichlet L -functions may be written as a linear combination of Hurwitz zeta function at rational values. Fixing an integer $k \geq 1$, Dirichlet L -functions for characters modulo k are linear combinations, with constant coefficients, of $\zeta(s, a)$ where $a = \frac{r}{k}$ and $r = 1, 2, 3, \dots, k$. This means Hurwitz zeta function for rational a has analytic properties that are closely related to Dirichlet L -functions. Specifically, let χ be a character modulo k . Then we can write its Dirichlet L -function as $L(s, \chi) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s} = \frac{1}{k^s} \sum_{r=1}^k \chi(r) \zeta\left(s, \frac{r}{k}\right)$.

In more details: Dirichlet L -functions satisfy a functional equation, which provides a way to analytically continue them throughout the complex plane. The functional equation relates the value of $L(s, \chi)$ to

the value of $L(1-s, \bar{\chi})$. Let χ be a primitive character modulo q , where $q > 1$. One way to express functional equation is $L(s, \chi) = \varepsilon(\chi) 2^s \pi^{s-1} q^{1/2-s} \sin\left(\frac{\pi}{2}(s+a)\right) \Gamma(1-s) L(1-s, \bar{\chi})$. In this equation, Γ denotes gamma function; a is 0 if $\chi(-1) = 1$, or 1 if $\chi(-1) = -1$; and $\varepsilon(\chi) = \frac{\tau(\chi)}{i^a \sqrt{q}}$ where $\tau(\chi)$ is a Gauss sum $\tau(\chi) = \sum_{n=1}^q \chi(n) \exp(2\pi i n/q)$. It is a property of Gauss sums that $|\tau(\chi)| = q^{1/2}$, so $|\varepsilon(\chi)| = 1$. Another way to state functional equation is in terms of $\xi(s, \chi) = \left(\frac{q}{\pi}\right)^{(s+a)/2} \Gamma\left(\frac{s+a}{2}\right) L(s, \chi)$. The functional equation is expressed as $\xi(s, \chi) = \varepsilon(\chi) \xi(1-s, \bar{\chi})$. The functional equation implies $L(s, \chi)$ and $\xi(s, \chi)$ are entire functions of s . Again, this assumes χ is primitive character modulo q with $q > 1$. If $q = 1$, then $L(s, \chi) = \zeta(s)$ has a pole at $s = 1$.

In mathematics and theoretical physics, techniques of zeta function regularization, dimensional regularization and analytic regularization are types of regularization or summability methods that assigns finite values to divergent sums or products. They are then used to define determinants and traces of some self-adjoint operators [which admit orthonormal eigenbasis with real eigenvalues]. Inspired by Method of Smoothed asymptotics developed by Prof. Terence Tao in 2010, we broadly base some deductions in this paper on introduction in 2024 by Prof. Antonio Padilla and Prof. Robert Smith of a new ultra-violet regularization scheme for loop integrals in Quantum field theory dubbed η regularization. We outline in section 4 rich underlying connections between analytic number theory and perturbative quantum field theory.

The functoriality conjecture states that a suitable homomorphism of L -groups is expected to give a correspondence between automorphic forms (in the global case) or representations (in the local case). Roughly speaking, the Langlands reciprocity conjecture is the special case of the functoriality conjecture when one of the reductive groups is trivial. *Broadly viewed as vast "resource materials" that support completed 2001 proofs on modularity theorem*, we have bi-directional correspondences (bridges) existing between Number theory \leftrightarrow Harmonic analysis forming "framework" for L -functions and modular forms database (LMFDB, launched on May 10, 2016)[3] involving reciprocity conjecture, functoriality conjecture, etc: (i) {Elliptic curves \leftrightarrow Modular forms}; (ii) {Counting problem $1+p$ -number of solutions mod p [in *finite series* Elliptic curves] \leftrightarrow Coefficients of q^p [in *infinite series* Modular forms]} whereby nome $q = e^{\pi i \tau}$ & $p =$ prime numbers from Modular forms act as (periodic) 'generating series or functions' having Group of symmetry = $SL_2(\mathbb{Z})$ [involving unit disk in complex plane], which is analogous to Group of symmetry = Group of integers \mathbb{Z} [involving real number line present in general solutions such as $\sin(x + 2\pi n) = \sin(x)$ with $n = \dots -3, -2, -1, 0, 1, 2, 3, \dots$]; viz, these properties conform to the *Langlands program "Theory of Symmetry"* [for Transformations of Rotation, Translation, Dilation and Reflection]; and (iii) {Representations of Galois groups \leftrightarrow Automorphic forms} whereby Modular forms are classified as a specific type of these [more general] Automorphic forms, which are ultimate objects in Harmonic analysis.

Diophantine equations are effectively various "*finite series*" polynomial equations that generally involve operation of adding finitely many terms e.g. Fermat's equation $x^n + y^n = z^n$ and elliptic curve $y^2 = x^3 + ax + b$. Proposed by Pierre de Fermat in 1637, Fermat's Last Theorem states that no three positive

integers a , b and c can satisfy Fermat's equation for any integer value of n greater than 2. The 2001 modularity theorem asserts that every elliptic curve is modular. This meant that all elliptic curves are associated with unique "infinite series" modular forms. In a nutshell, this was broadly a crucial step in proving Fermat's Last Theorem because it famously allowed Prof. Andrew Wiles to prove the theorem in 1994 by establishing a deep connection between [semistable] elliptic curves and modular forms. Sir Andrew Wiles was deservedly awarded the 2016 Abel Prize for this work.

In simpler language, Birch and Swinnerton-Dyer (BSD) conjecture asserts an elliptic curve E has either an infinite number or a finite number of rational points (solutions) according to whether $\zeta(1) = 0$ or $\zeta(1) \neq 0$, respectively [when these rational points (solutions) are points of an abelian variety and $\zeta(1)$ is an associated zeta function $\zeta(s)$ near point $s = 1$]. The rank of $E(\mathbb{Q})$ [\equiv subset of $E(\mathbb{Q})$ with its elements having infinite order] is the finite number of copies of \mathbb{Z} in $E(\mathbb{Q})$ or, equivalently, the finite number of independent points on an elliptic curve mod p . It is the rank of E (denoted here by r_E). Rank 0 [having zero independent basis point with infinite order]: There is a subset of $E(\mathbb{Q})$ in E either with zero finite integral points and zero finite $E(\mathbb{Q})$ solutions or with non-zero finite integral points and non-zero finite $E(\mathbb{Q})$ solutions]. Rank 1 [having one independent basis point with infinite order]: There is a subset of $E(\mathbb{Q})$ in E with non-zero finite integral points and infinite $E(\mathbb{Q})$ solutions. Higher Ranks 2 or more [having two or more independent basis point with infinite order]: There are subsets of $E(\mathbb{Q})$ with non-zero finite integral points and infinite $E(\mathbb{Q})$ solutions. The simplest ("strong") version of BSD conjecture: **Proposition.** In an elliptic curve E , there are infinitely many $E(\mathbb{Q})$ solutions when $L_E(1) = 0$; viz, when Central value [at $s = 1$] = 0. **Corollary.** In an elliptic curve E , there are finitely many (or zero) $E(\mathbb{Q})$ solutions when $L_E(1) \neq 0$; viz, when Central value [at $s = 1$] $\neq 0$. The standard ("weak") version of BSD conjecture asserts r_E can be arbitrarily large, and Algebraic r_E [Order of zero at $s = 1$ in $L(E, s)$] = Analytic r'_E [related to leading coefficient of Taylor expansion of $L(E, s)$ at $s = 1$].

We have infinities or infinitely large numbers as the unbounded and limitless quantities (∞) at the big end, and infinitesimals or infinitely small numbers as the extremely small but nonzero quantities ($\frac{1}{\infty}$) at the small end. Applying infinitesimals to their corresponding outputs in section 6 allow us to prove 1859-dated Riemann hypothesis [viz, the proposal that relevant outputs as infinitely many nontrivial zeros or Origin intercept points of Riemann zeta function are all located on its $\sigma = \frac{1}{2}$ -critical line or $\sigma = \frac{1}{2}$ -Origin point], and Polignac's and Twin prime conjectures [viz, the proposal that relevant outputs as subsets of Odd Primes derived from every even Prime gaps 2, 4, 6, 8, 10... all contain infinitely many unique elements]. Referring to even Prime gap 2, 1846-dated Twin prime conjecture is simply a subset of 1849-dated Polignac's conjecture [which refers to all even Prime gaps 2, 4, 6, 8, 10...]. Altered terminology on cardinality of Odd Primes being *arbitrarily large number* (ALN) instead of *infinitely many* was previously used by us to denote *Modified* Polignac's and Twin prime conjectures.

Our *generic mathematical approaches* for solving Riemann hypothesis, Polignac's and Twin prime conjectures is applicable to selected branches of science such as relativistic quantum mechanics, quantum gravity or string theory. When usefully construed as a self-sufficient research paper, correct and complete

mathematical arguments condensed in this current paper are the major (core) arguments from publications [6], [7] & [8] whereby Riemann zeta function [= function that faithfully generates output of all nontrivial zeros via its *proxy* Dirichlet eta function] and Sieve of Eratosthenes [= algorithm that faithfully generates output of all prime numbers] are treated as de novo or derived infinite series in order to prove their connected open problems in Number theory. These infinite series are either convergent series or divergent series where partial sums of sequence from the former tends to a finite limit, while that from the later do not tend to a finite limit [viz, it tends to infinity]. Prime number theorem for Arithmetic Progressions [as Axiom 1], Generic Squeeze theorem [as Theorem 3.1] and Theorem of Divergent-to-Convergent series conversion for Prime numbers [as Theorem 4.1] are outlined (respectively) in section 2, section 3 and section 4. Lemma 5.1 and Lemma 5.2 in section 5 (respectively) introduce novel concept of Incompletely Predictable entities and innovatively classifying countably infinite sets into accelerating, linear or decelerating subtypes. To the extent that some associated minor (peripheral) arguments were not included in this paper, we advocate their absence do not adversely reflect the rigorous nature of our derived proofs but, rather, helps disseminate mathematical knowledge to the lay person and scientific community.

A function [sometimes loosely termed an operator or an equation] is usefully defined as relation between a set of inputs (called domain) and a set of possible outputs (called codomain) where each input is related to EXACTLY one output. More precisely, classical example of [linear] operator performed on [eligible] functions is differentiation. An algorithm is usefully defined as finite sequence of rigorous instructions typically used to solve a class of specific problems or to perform a computation. Functions or algorithms as infinite-dimensional vectors: A function or algorithm defined on real numbers \mathbb{R} can be represented by an uncountably infinite set of vectors (as a vector field) while a function or algorithm defined on natural numbers \mathbb{N} [or any other countably infinite domain such as prime numbers and composite numbers] can be represented by a countably infinite set of vectors (as a vector field). One could also use the later countably infinite set of vectors involving [discrete] \mathbb{N} {e.g. all nontrivial zeros of Riemann zeta function interpolated as "nearest" t -valued \mathbb{N} 14, 21, 25, 30, 33, 38, 41, 43...} to approximate the former uncountably infinite set of vectors that "pseudo-represent" [continuous] \mathbb{R} {for the same nontrivial zeros when precisely given as t -valued transcendental numbers} \cong Law of continuity: If a quantity changes "continuously", then its value at any point between two given values can be determined by the process of interpolation.

Based on Figure 1 and Figure 2 that accommodate both positive (+ve) parts and negative (-ve) counterparts of prime numbers, composite numbers and nontrivial zeros, **we can represent eligible functions with complex vector space [having +ve and -ve complex vectors pointing in opposite directions] and eligible algorithms with real vector space [having +ve and -ve real vectors pointing in opposite directions]**: Recall that a row vector or a column vector is, respectively, a one-row matrix or a one-column matrix. Real numbers \mathbb{R} [and natural numbers \mathbb{N}] are exactly one-dimensional vectors (on a line) and complex numbers \mathbb{C} are exactly two-dimensional vectors (in a plane). A complex vector (or complex matrix) as Cartesian representation $z = x + iy$ or Polar representation $z = r(\cos \theta + i \cdot \sin \theta)$ is simply a vector (matrix) of the complex numbers. A two-dimensional real vector (or real matrix) in a plane is

One-dimensional integer number line containing Prime and Composite numbers




Figure 1. Narrow range of positive & negative prime and composite numbers plotted together on integer number line generated using Sieve-of-Eratosthenes and complement-Sieve-of-Eratosthenes. The combined [positive] image and [negative] mirror image will conceptually represent a one-dimensional line (state) having perfect Mirror symmetry with integer number 0 acting as the Point of symmetry.




Figure 2. OUTPUT for $\sigma = \frac{1}{2}$ as Gram points. Polar graph of $\zeta(\frac{1}{2} + it)$ depicted as a two-dimensional figure (state) plotted along critical line for real values of t between -30 and $+30$ [viz, for $s = \sigma \pm t$ range], horizontal axis: $Re\{\zeta(\frac{1}{2} + it)\}$, and vertical axis: $Im\{\zeta(\frac{1}{2} + it)\}$. Origin intercept points are present. There is manifestation of perfect Mirror symmetry about horizontal x-axis acting as the line of symmetry.

given by Cartesian representation as $v = x + iy$ or Polar representation as $v = r(\cos \theta + i \sin \theta)$. x & y are \mathbb{R} , modulus $r = |z|$ or $|v| = \sqrt{x^2 + y^2}$, multi-valued $\arg(z)$ or $\arg(v)$ or principal-valued $\text{Arg}(z)$ or $\text{Arg}(v) = \theta = \arctan(y/x)$, and imaginary unit $i = \sqrt{-1}$.

Integers $\{0, 1\}$ are neither prime nor composite. Prime & composite numbers form distinct countably infinite sets of integers as two subsets in uncountably infinite set of real numbers. Both [algorithmic] inputs Sieve-of-Eratosthenes and Complement-Sieve-of-Eratosthenes in section 2 that faithfully generate outputs prime & composite numbers are visually represented by countably infinite set of real vectors. We recognize all real vector sub-spaces for even Prime gaps 2, 4, 6, 8, 10... with each unique sub-space

constituted by its corresponding countably infinite set of real vectors, must imply Modified Polignac's and Twin prime conjectures are true.

Where $\sigma, t, Re\{\zeta(s)\}, Im\{\zeta(s)\}, Re\{\eta(s)\}$ and $Im\{\eta(s)\}$ are \mathbb{R} , (input) parameter $s = \sigma \pm it$ used in (output) functions from section 2 such as non-alternating Riemann zeta function Eq. 1 $\zeta(s) = Re\{\zeta(s)\} + i \cdot Im\{\zeta(s)\}$ and alternating Dirichlet eta function Eq. 2 $\eta(s) = Re\{\eta(s)\} + i \cdot Im\{\eta(s)\}$ are recognized to all be given in $z = x + iy$ format, thus allowing uncountably infinite set of complex vectors to visually represent them. Next consider the two derived functions from section 2: simplified Dirichlet eta function or $sim-\eta(s)$ and Dirichlet Sigma-Power Law or DSPL [= $\int sim-\eta(s)dn \equiv$ "signed area under a curve" for this Riemann integrable function] with their corresponding horizontal and vertical axes being perpendicular to each other or, equivalently, being $\frac{\pi}{2}$ out-of-phase with each other (as per Page 12 of [6]). Complex vectors representing $sim-\eta(s)$ and DSPL when combined together form an orthonormal set in the inner product space since all these vectors in the set are mutually orthogonal ("perpendicular") and can be depicted using their ("normalized") unit length. When equivalently expressed using countably infinite set of complex vectors; we recognize nontrivial zeros of $\zeta(s), \eta(s), sim-\eta(s)$ or DSPL that only exist in unique $\sigma = \frac{1}{2}$ complex vector sub-space, must imply Riemann hypothesis is true.

Non-alternating power series $\sum_{n=0}^{\infty} a_n x^n = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \dots$

Alternating power series $\sum_{n=0}^{\infty} (-1)^n a_n x^n = a_0 - a_1 x + a_2 x^2 - a_3 x^3 + \dots$

Non-alternating harmonic series $\sum_{n=1}^{\infty} \frac{1}{n} = \frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \dots$

Alternating harmonic series $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} = \frac{1}{1} - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \dots$
--

When $s = 1$ in Eq. 1 $\zeta(s)$ & Eq. 2 $\eta(s)$ with $n = +ve$ integers, we (respectively) obtain the above **most basic Non-alternating harmonic series and Alternating harmonic series**. An infinite series [listed above as various types of power series and harmonic series] (or a finite series) is sum of $[\geq 1]$ infinite (or finite) sequence of terms constituted by numbers, scalars, or anything *e.g. functions, vectors, matrices*.

As previously discussed, power series [with VARYING coefficients a_n] are infinite polynomials. Sieve-of-Eratosthenes & Complement-Sieve-of-Eratosthenes as well-defined infinite algorithms give rise to [infinite] n solutions of all primes & composites; viz, they are the "analogs" of power or harmonic series as well-defined infinite functions. With SAME coefficients a , the (non-alternating) geometric series

$\sum_{n=0}^{\infty} ax^n = a + ax + ax^2 + ax^3 + \dots$ having +ve common ratio x between successive terms, is simply a special case of (non-alternating) power series e.g. when $a = \frac{1}{2}$ & $\frac{1}{2}$ for +ve common ratio. **Cf** when $a = \frac{1}{2}$ & $-\frac{1}{2}$ for -ve common ratio in an "inverse" (alternating) geometric series, which is simply a special case of (alternating) power series (Page 56 of [8]):

$$\sum_{n=0}^{\infty} \frac{1}{2} \left(-\frac{1}{2}\right)^n = \frac{1}{2} - \frac{1}{4} + \frac{1}{8} - \frac{1}{16} + \dots = \frac{\frac{1}{2}}{1 - (-\frac{1}{2})} = \frac{1}{3}$$

cf $\sum_{n=0}^{\infty} \frac{1}{2} \left(\frac{1}{2}\right)^n = \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \dots = \frac{\frac{1}{2}}{1 - (+\frac{1}{2})} = 1.$

Power and Harmonic (infinite) series defined over prime numbers p , with $a_{p_n} = p_n$, for example:

Non-alternating power series $\sum_{n=1}^{\infty} a_{p_n} x^{p_n} = \sum_{n=1}^{\infty} p_n x^{p_n} = 2x^2 + 3x^3 + 5x^5 + 7x^7 + 11x^{11} + \dots$

Alternating power series $\sum_{n=1}^{\infty} (-1)^{n-1} a_{p_n} x^{p_n} = \sum_{n=1}^{\infty} (-1)^{n-1} p_n x^{p_n} = 2x^2 - 3x^3 + 5x^5 - 7x^7 + 11x^{11} - \dots$

Non-alternating harmonic series $\sum_{n=1}^{\infty} \frac{1}{p} = \frac{1}{2} + \frac{1}{3} + \frac{1}{5} + \frac{1}{7} + \frac{1}{11} + \dots$

Alternating harmonic series $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{p} = \frac{1}{2} - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{11} - \dots$

An expression is in closed form if it is formed with constants, variables and a finite set of basic functions connected by arithmetic operations (viz, +, -, ×, ÷, and integer powers) and function composition. The commonly allowed functions are [I] the algebraic functions [viz, defined as the root of an irreducible polynomial equation] e.g. n^{th} root or raising to a fractional power and [II] the transcendental (non-algebraic) functions e.g. exponential function, logarithmic function, Γ function, trigonometric functions and their inverses. Algebraic and transcendental (non-algebraic) solutions form two subsets of closed-form expressions. Thus, a solution in radicals or algebraic solution is a closed-form expression, and more specifically a closed-form algebraic expression, that is the solution of a polynomial equation, and relies only on addition, subtraction, multiplication, division, raising to integer powers, and the extraction of n^{th} roots (square roots, cube roots, and other integer roots). Following directly from Galois theory using polynomial $f(x) = x^5 - x - 1$ as one of the simplest examples of a non-solvable quintic polynomial, Abel-Ruffini theorem states that there is no solution in radicals to SOME general (finite) polynomial equations of degree five or higher with arbitrary coefficients. Here, *general* meant the coefficients of a polynomial equation are viewed and manipulated as indeterminates. We extrapolate: Any power series [e.g. e^x , $\sin x$, $\sinh x$, $\ln x$, etc] as general (infinite) polynomial equations having infinitely many coefficients should have no solution in radicals [viz, have transcendental solutions]. However some power series with coefficients involving (infinite) polynomials [e.g. geometric series, binomial series, etc] can have solutions expressible in terms of radicals, provided the series converges within the domain where such expressions are valid. Similar to, but not categorized as, power series are various hypergeometric series [as defined by the generalized hypergeometric function] that could have either transcendental solutions or solutions in radicals.

Eq. 1 $\zeta(s)$ & Eq. 2 $\eta(s)$ have complex variable $s = \sigma \pm it$. In $0 < \sigma < 1$ critical strip containing $\sigma = \frac{1}{2}$ critical line, $\eta(s)$ must act as proxy function for $\zeta(s)$ [with both \equiv infinite series]. Useful relationship: z as a complex number \mathbb{C} is defined by $z = a + bi$ with i being the imaginary unit, and a & b being real numbers \mathbb{R} . Then $\mathbb{R} \subset \mathbb{C}$ since when $b = 0$, $z = a + 0i = a$ will always be \mathbb{R} . Our "amalgated" *generic Fundamental Theorem of Algebra* heuristically \implies (eligible) general [finite or infinite or ALN] algorithms and functions (of degree n with real or complex coefficients) have exactly [finite or infinite or ALN] n roots or n solutions as real or complex numbers, counting *multiplicities* {e.g. $\sin(x + 2\pi n)$ with $n = \dots -3, -2, -1, 0, 1, 2, 3, \dots$; \pm nontrivial zeros; \pm Primes; \pm Composites; etc}. Riemann hypothesis is




Figure 3. INPUT for $\sigma = \frac{1}{2}$ (for Figure 4), $\frac{2}{5}$ (for Figure 5), and $\frac{3}{5}$ (for Figure 6). Riemann zeta function $\zeta(s)$ has two countable infinite sets of firstly, Completely Predictable trivial zeros located at $s =$ all negative even numbers and secondly, Incompletely Predictable nontrivial zeros located at $\sigma = \frac{1}{2}$ as various t -valued transcendental numbers.

true when nontrivial zeros as Origin point intercepts are the infinitely many n roots that only occur when parameter $\sigma = \frac{1}{2}$ resulting in [optimal] "formula symmetry" for $\eta(s)$ [as **infinite series**]. Polignac's and Twin prime conjectures are true when Sieve-of-Eratosthenes algorithm and its derived sub-algorithms [as "**infinite series**" via $\sum_{n=i}^{ALN} p_{n+1} = 3 + \sum_{i=2}^n g_i$] have ALN of n solutions represented by the Set [\equiv total] of Odd Primes and Subsets [\equiv subtotals] of Odd Primes derived from all even Prime gaps.

2. General notations including Prime number theorem for Arithmetic Progressions and creating *de novo* Infinite Series

Common abbreviations used in this paper: CP = Completely Predictable, IP = Incompletely Predictable, FL = Finite-Length, IL = Infinite-Length, CFS = countably finite set, CIS = countably infinite set, IM = infinitely-many, ALN = arbitrarily large number. We treat eligible algorithms and functions as *de novo* infinite series. Critical strip $\equiv \{s \in \mathbb{C} : 0 < \text{Re}(s) < 1\}$ & Critical line $\equiv \{s \in \mathbb{C} : \text{Re}(s) = \frac{1}{2}\}$ in Figure 3. Phrase "*inside* the critical strip" refers to parameter $s [= \sigma \pm it$ with $0 < \sigma < 1$; viz, $0 < \text{Re}(s) < 1$] having complex number values defined for $\eta(s)$ as given by parameter t over \pm real numbers. Phrase "*outside* the critical strip" refers to parameter $s [= \sigma \pm it$ with $\sigma > 1$; viz, $\text{Re}(s) > 1$] having complex number values defined for $\zeta(s)$ as given by parameter t over \pm real numbers.

List of abbreviations incorporating relevant definitions:

- CP entities**: These entities manifest CP *independent* properties.
- IP entities**: These entities manifest IP *dependent* properties.
- $\zeta(s)$: $f(n)$ Riemann zeta function [\equiv **infinite (converging) series** for $\text{Re}(s) > 1$] – see Eq. (1) below containing variable n , and parameters t and σ will generate [via its *proxy* Dirichlet eta function] Zeroes when $\sigma = \frac{1}{2}$ and virtual Zeroes when $\sigma \neq \frac{1}{2}$.
- $\eta(s)$: $f(n)$ Dirichlet eta function [\equiv **infinite (converging) series** for $\text{Re}(s) > 0$] – see Eq. (2) below as the analytic continuation of $\zeta(s)$, containing variable n , and parameters t and σ will generate Zeroes when




Figure 4. OUTPUT for $\sigma = \frac{1}{2}$ as Gram points. Polar graph of $\zeta(\frac{1}{2} + it)$ plotted along critical line for real values of t running from 0 to 34. Horizontal axis: $Re\{\zeta(\frac{1}{2} + it)\}$. Vertical axis: $Im\{\zeta(\frac{1}{2} + it)\}$. Presence of Origin intercept points. **Nil-shift w.r.t. Origin point when $\sigma = \frac{1}{2}$.**




Figure 5. OUTPUT for $\sigma = \frac{2}{5}$ as virtual Gram points. Varying Loops are shifted to left of Origin with horizontal axis: $Re\{\zeta(\frac{2}{5} + it)\}$, and vertical axis: $Im\{\zeta(\frac{2}{5} + it)\}$. Nil Origin intercept points. **Left-shift w.r.t. Origin point when $\sigma < \frac{1}{2}$; viz, $0 < \sigma < \frac{1}{2}$.**




Figure 6. OUTPUT for $\sigma = \frac{3}{5}$ as virtual Gram points. Varying Loops are shifted to right of Origin with horizontal axis: $Re\{\zeta(\frac{3}{5} + it)\}$, and vertical axis: $Im\{\zeta(\frac{3}{5} + it)\}$. Nil Origin intercept points. **Right-shift w.r.t. Origin point when $\sigma > \frac{1}{2}$; viz, $\frac{1}{2} < \sigma < 1$.**




Figure 7. Close-up view of virtual Origin points when $\sigma = \frac{1}{3}$. OUTPUT for $\sigma = \frac{1}{3}$ [$\sigma < \frac{1}{2}$ situation] as virtual Gram points. Polar graph of $\zeta(\frac{1}{3} + it)$ plotted along non-critical line for real values of t running between 0 and 100, horizontal axis: $Re\{\zeta(\frac{1}{3} + it)\}$, and vertical axis: $Im\{\zeta(\frac{1}{3} + it)\}$. Total absence of all Origin intercept points at "static" Origin point. Total presence of all virtual Origin intercept points (as additional negative virtual Gram[$y=0$] points on x -axis) at "varying" [infinitely many] virtual Origin points. **With respect to $\sigma = \frac{1}{2}$ -Origin point being analogically the $\sigma = \frac{1}{2}$ -'Centroid', then the [depicted] "left-shifted" $\sigma = \frac{1}{3}$ as being $\frac{1}{3} - \frac{1}{2} = -\frac{1}{6}$ and the [undepicted] "right-shifted" $\sigma = \frac{2}{3}$ as being $\frac{2}{3} - \frac{1}{2} = +\frac{1}{6}$ are BOTH equidistant from 'Centroid' [thus fully satisfying (Remark 2.2) Principle of Equidistant for Multiplicative Inverse – see last paragraph discussion in section 6 Conclusions.**

$\sigma = \frac{1}{2}$ and virtual Zeroes when $\sigma \neq \frac{1}{2}$.

·**sim- $\eta(s)$** : $f(n)$ simplified Dirichlet eta function [\equiv infinite (converging) series for $Re(s) > 0$], derived by applying Euler formula to $\eta(s)$, containing variable n , and parameters t and σ will generate Zeroes when $\sigma = \frac{1}{2}$ – see Eq. (4) below and virtual Zeroes when $\sigma \neq \frac{1}{2}$ – see Eq. (5) below.

·**DSPL**: $F(n)$ Dirichlet Sigma-Power Law [\equiv "continuous" infinite (converging) series for $Re(s) > 0$] = $\int \text{sim-}\eta(s)dn$ containing variable n , and parameters t and σ will generate Pseudo-zeroes when $\sigma = \frac{1}{2}$ – see Eq. (6) below and virtual Pseudo-zeroes when $\sigma \neq \frac{1}{2}$ whereby the (virtual) Zeros = (virtual) Pseudo-zeros $-\frac{\pi}{2}$ relationship allows (virtual) Pseudo-zeros to (virtual) Zeros conversion and *vice versa*.

·**NTZ**: Nontrivial zeros located on the one-dimensional (mathematical) $\sigma = \frac{1}{2}$ -critical line are precisely equivalent to **G[x=0,y=0]P**: Gram[x=0,y=0] points as Origin intercept points which are located at the zero-dimensional (geometrical) $\sigma = \frac{1}{2}$ -Origin point [as per Figure 4]. These entities, mathematically defined by $\sum ReIm\{\eta(s)\} = Re\{\eta(s)\} + Im\{\eta(s)\} = 0$, are generated by equation G[x=0,y=0]P- $\eta(s)$ containing exponent $\frac{1}{2}$ when $\sigma = \frac{1}{2}$.

·**GP or G[y=0]P**: 'usual' or 'traditional' Gram points = Gram[y=0] points = x -axis intercept points that are [multiple-positioned] located on one-dimensional x -axis line are generated by equation G[y=0]P- $\eta(s)$ when $\sigma = \frac{1}{2}$. These entities are mathematically defined by $\sum ReIm\{\eta(s)\} = Re\{\eta(s)\} + 0$, or simply $Im\{\eta(s)\} = 0$. Riemann hypothesis is usefully stated as none of the [additional] virtual G[x=0]P generated by equation G[x=0]P- $\eta(s)$ when $\sigma \neq \frac{1}{2}$ – as demonstrated by Figure 7 for $\sigma = \frac{1}{3}$ – can be constituted by

t transcendental number values that [incorrectly] coincide with t transcendental number values for NTZ when $\sigma = \frac{1}{2}$.

·**G[x=0]P**: Gram[x=0] points = y-axis intercept points that are [multiple-positioned] located on one-dimensional y-axis line are generated by equation $G[x=0]P-\eta(s)$ when $\sigma = \frac{1}{2}$. These entities are mathematically defined by $\sum \text{ReIm}\{\eta(s)\} = 0 + \text{Im}\{\eta(s)\}$, or simply $\text{Re}\{\eta(s)\} = 0$.

·**virtual NTZ**: virtual nontrivial zeros or **virtual G[x=0,y=0]P**: virtual Gram[x=0,y=0] points. These are virtual Origin intercept points located at the multiple-positioned virtual Origin points which are generated by equation virtual- $G[x=0,y=0]P-\eta(s)$ containing exponent values $\neq \frac{1}{2}$ when $\sigma \neq \frac{1}{2}$. We note that each virtual NTZ when $\sigma < \frac{1}{2}$ in Figure 5 equates to an [additional] negative virtual $G[y=0]P$ located at IP varying positions on horizontal axis, and each virtual NTZ when $\sigma > \frac{1}{2}$ in Figure 6 equates to an [additional] positive virtual $G[y=0]P$ located at IP varying positions on horizontal axis. We observe overall less virtual $G[x=0]P$ when $\sigma > \frac{1}{2}$, and overall more virtual $G[x=0]P$ when $\sigma < \frac{1}{2}$.

·**Sieve-of-Eratosthenes (S-of-E)**: For $i = 1, 2, 3, 4, 5\dots$ and with $p_1 = 2$ [\equiv even prime number 2 forming a CFS with cardinality of 1] as the first term in S-of-E; the algorithm S-of-E as symbolically denoted by $p_{n+1} = 2 + \sum_{i=1}^n g_i$ with $g_n = p_{n+1} - p_n$ and its derived sub-algorithms faithfully generate the set of all prime numbers 2, 3, 5, 7, 11, 13... and subsets of Odd Primes derived from even Prime gaps 2, 4, 6, 8, 10... We now ignore even prime number 2 by changing variable i to instead commence from 2^{nd} position. For $i = 2, 3, 4, 5, 6\dots$ and with $p_2 = 3$ [\equiv first Odd Prime 3] as the first term in Modified-S-of-E; the altered algorithm Modified-S-of-E as symbolically denoted by $p_{n+1} = 3 + \sum_{i=2}^n g_i$ with $g_n = p_{n+1} - p_n$ and its derived sub-algorithms will faithfully generate the set of all Odd Primes 3, 5, 7, 11, 13, 17... and subsets of Odd Primes derived from even Prime gaps 2, 4, 6, 8, 10... By performing summation [viz, conducting repeated addition of sequence from ALN of prime gaps and prime numbers that are arranged in a unique order] on above two algorithms as $\sum_{n=i}^{ALN} p_{n+1} = 2 + \sum_{i=1}^n g_i$ and $\sum_{n=i}^{ALN} p_{n+1} = 3 + \sum_{i=2}^n g_i$, we obtain (de novo) infinite series. These infinite series are all **diverging series** for this two algorithms [and their derived sub-algorithms]. In contrast, Brun's constants as outlined in section 4 are **converging series**. The cardinality of CIS-ALN-decelerating is applicable for (i) set of all prime numbers, (ii) set of all Odd Primes, (iii) subsets of Odd Primes, and (iv) set of all even Prime gaps \implies Modified Polignac's and Twin prime conjectures are true.

·**Complement-Sieve-of-Eratosthenes**: For $i = 1, 2, 3, 4, 5\dots$ and with $c_1 = 4$; this algorithm as symbolically denoted by $c_{n+1} = 4 + \sum_{i=1}^n c_i$ with $g_n = c_{n+1} - c_n$ and its derived sub-algorithms will faithfully generate all composite numbers. Parallel arguments to construct de novo infinite series as **diverging series** for (sub)sets of composite numbers are also possible.

In general, we note the infinite-length sequence of a given converging series or diverging series can theoretically be constituted by either positive terms e.g. $\zeta(s)$ as non-alternating harmonic series Eq. (1) OR

alternating positive and negative terms e.g. $\eta(s)$ as alternating harmonic series Eq. (2).

$$(1) \quad \zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} \\ = \frac{1}{1^s} + \frac{1}{2^s} + \frac{1}{3^s} + \frac{1}{4^s} + \frac{1}{5^s} + \dots \\ = \prod_{p \text{ prime}} \frac{1}{1-p^{-s}} = \frac{1}{1-2^{-s}} \cdot \frac{1}{1-3^{-s}} \cdot \frac{1}{1-5^{-s}} \cdot \frac{1}{1-7^{-s}} \cdot \frac{1}{1-11^{-s}} \dots \frac{1}{1-p^{-s}} \dots$$

Eq. (1) non-alternating harmonic series Riemann zeta function $\zeta(s)$ is a function of complex variable s ($= \sigma \pm it$) that continues sum of infinite series $\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = \frac{1}{1^s} + \frac{1}{2^s} + \frac{1}{3^s} + \dots$ for $\text{Re}(s) > 1$, and its analytic continuation elsewhere for $0 < \text{Re}(s) < 1$. Containing no nontrivial zeros, $\zeta(s)$ is defined only in $1 < \sigma < \infty$ region where it is absolutely convergent. The common convention is to write s as $\sigma + it$ with $t = \sqrt{-1}$, and with σ and t real. Valid for $\sigma > 1$, we write $\zeta(s)$ as $\text{Re}\{\zeta(s)\} + i\text{Im}\{\zeta(s)\}$ and note that $\zeta(\sigma + it)$ when $0 < t < +\infty$ is the complex conjugate of $\zeta(\sigma - it)$ when $-\infty < t < 0$. In Eq. (1), the equivalent **Euler product formula** with product over all prime numbers implies the presence of Sieve of Eratosthenes. Also note that for all $s \in \mathbb{C}$, $s \neq 1$, the integral relation $\zeta(s) = \frac{1}{s-1} + \frac{1}{2} + 2 \int_0^{\infty} \frac{\sin(s \arctan t)}{(1+t^2)^{s/2} (e^{2\pi t} - 1)} dt$ holds true, which may be used for a numerical evaluation of the zeta function.

$$(2) \quad \eta(s) = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^s} = \frac{1}{1^s} - \frac{1}{2^s} + \frac{1}{3^s} - \frac{1}{4^s} + \frac{1}{5^s} - \dots \\ = \prod_{p \text{ prime}} \frac{1}{1 \pm p^{-s}} = \frac{1}{1+2^{-s}} \cdot \frac{1}{1-3^{-s}} \cdot \frac{1}{1+5^{-s}} \cdot \frac{1}{1-7^{-s}} \cdot \frac{1}{1+11^{-s}} \dots \frac{1}{1 \pm p^{-s}} \dots$$

Eq. (2) alternating harmonic series Dirichlet eta function $\eta(s)$ that faithfully generates all three types of Gram points as three dependent CIS-IM-linear Incompletely Predictable entities when $\sigma = \frac{1}{2}$ must represent and act as *proxy* function for Eq. (1) in $0 < \sigma < 1$ -critical strip [viz, for $0 < \text{Re}(s) < 1$] containing $\sigma = \frac{1}{2}$ -critical line because $\zeta(s)$ only converges when $\sigma > 1$. In Eq. (2), the equivalent **Euler product formula** with product over all prime numbers also implies the presence of Sieve of Eratosthenes. They are related to each other [via Analytic continuation] as $\zeta(s) = \gamma \cdot \eta(s)$ or equivalently as $\eta(s) = \frac{1}{\gamma} \cdot \zeta(s)$ with proportionality factor $\gamma = \frac{1}{(1-2^{1-s})}$.

$$(3) \quad \zeta(s) = 2^s \pi^{s-1} \sin\left(\frac{\pi s}{2}\right) \Gamma(1-s) \zeta(1-s)$$

[The reflection functional equation for $\eta(s)$ is already given previously and not reproduced here.] $\zeta(s)$ satisfies Eq. (3) as the reflection functional equation whereby Γ is the gamma function. [Note that derived for complex numbers with a positive real part, Γ is defined via a convergent improper integral $\Gamma(z) = \int_0^{\infty} t^{z-1} e^{-t} dt$, $\Re(z) > 0$. Γ is then defined as analytic continuation of this integral function to a meromorphic function that is holomorphic in whole complex plane except zero and negative integers, where the function has simple poles. The main motivation for its development is $\Gamma(x+1)$ interpolates factorial function $x! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot x$ to non-integer values.] As an equality of meromorphic functions

valid on whole complex plane, Eq. (3) relates values of $\zeta(s)$ at points s and $1 - s$; in particular, it relates even positive integers with odd negative integers. Owing to zeros of sine function, the functional equation implies $\zeta(s)$ has a simple zero at each even negative integer $s = -2n = -2, -4, -6, -8, -10, \dots$ known as trivial zeros of $\zeta(s)$. When s is an even positive integer, product $\sin(\frac{\pi s}{2})\Gamma(1 - s)$ on right is non-zero because $\Gamma(1 - s)$ has a simple pole, which cancels simple zero of sine factor. With perfect line symmetry at the vertical line $s = \frac{1}{2}$, we have a **symmetric version** of this functional equation applied to the Lambda-function given by $\Lambda(s) = \pi^{-\frac{s}{2}}\Gamma\left(\frac{s}{2}\right)\zeta(s)$, which satisfies $\Lambda(s) = \Lambda(1 - s)$ OR to the Riemann xi-function given by $\xi(s) = \frac{1}{2}\pi^{-\frac{s}{2}}s(s - 1)\Gamma\left(\frac{s}{2}\right)\zeta(s)$, which satisfies $\xi(s) = \xi(1 - s)$. $\Lambda(s)$ OR $\xi(s)$ is thus the 'completed zeta function' whereby $\pi^{-\frac{s}{2}}\Gamma\left(\frac{s}{2}\right) = \Gamma_{\mathbb{R}}(s)$ is "Gamma factor" as the local L-factor corresponding to the Archimedean place, with the other factors in the Euler product expansion being the local L-factors of the non-Archimedean places. The conductor of L-function is the positive integer N from $N^{\frac{s}{2}}$. For Riemann zeta function, its conductor N as derived from $\pi^{-\frac{s}{2}} = \left(\frac{1}{\pi} \cdot 1\right)^{\frac{s}{2}}$ is 1.

Remark 2.1. Consider sequence $\lambda_n = \frac{1}{(n - 1)!} \frac{d^n}{ds^n} [s^{n-1} \log \xi(s)] \Big|_{s=1}$. As a possible pathway to solve Riemann hypothesis, Li's criterion states that this hypothesis is equivalent to the statement $\lambda_n > 0$ for every positive integer n [viz, positivity of λ_n].

The numbers λ_n (as the Additive invariants denoted by \mathbb{L}^* and sometimes defined with a slightly different normalization) are called Keiper-Li coefficients or Li coefficients. They may also be expressed in terms of nontrivial zeros of Riemann zeta function $\lambda_n = \sum_{\rho} \left[1 - \left(1 - \frac{1}{\rho} \right)^n \right]$ where the sum extends over ρ , the nontrivial zeros of the zeta function. This conditionally convergent sum should be understood in the sense usually used in Number theory; namely, that $\sum_{\rho} = \lim_{N \rightarrow \infty} \sum_{|\text{Im}(\rho)| \leq N}$. [Re(s) and Im(s) denote real and imaginary parts of s].

At $\sigma = \frac{1}{2}$, $\text{sim-}\eta(s) =$

$$(4) \quad \sum_{n=1}^{\infty} (2n)^{-\frac{1}{2}} 2^{\frac{1}{2}} \cos(t \ln(2n) + \frac{1}{4}\pi) - \sum_{n=1}^{\infty} (2n - 1)^{-\frac{1}{2}} 2^{\frac{1}{2}} \cos(t \ln(2n - 1) + \frac{1}{4}\pi)$$

At $\sigma = \frac{2}{5}$, $\text{sim-}\eta(s) =$

$$(5) \quad \sum_{n=1}^{\infty} (2n)^{-\frac{2}{5}} 2^{\frac{1}{2}} \cos(t \ln(2n) + \frac{1}{4}\pi) - \sum_{n=1}^{\infty} (2n - 1)^{-\frac{2}{5}} 2^{\frac{1}{2}} \cos(t \ln(2n - 1) + \frac{1}{4}\pi)$$

For any real number n , $e^{in} = \cos n + i \cdot \sin n$ is Euler's formula where e [\approx transcendental number 2.71828] is base of natural logarithm, $i = \sqrt{-1}$ is imaginary unit. When $n = \pi$ [\approx transcendental number 3.14159], then $e^{i\pi} + 1 = 0$ or $e^{i\pi} = -1$, known as Euler's identity. Applying this formula to f(n) $\eta(s)$ results in Eq. (4) f(n) simplified $\eta(s)$ at $\sigma = \frac{1}{2}$ that incorporate all nontrivial zeros [as Zeroes]. There is total absence of (non-existent) virtual nontrivial zeros [as virtual Zeroes]. Eq. (5) f(n) simplified $\eta(s)$ at $\sigma = \frac{2}{5}$ will incorporate all (non-existent) virtual nontrivial zeros [as virtual Zeroes]. There is total absence of nontrivial zeros [as Zeroes].




Figure 8. The natural logarithm function $\log_e x$ or $\ln(x)$ and natural exponential function $\exp(x)$ or e^x . The graphs of $\log_e x$ and its inverse e^x are symmetric with respect to line $y = x$ thus geometrically denoting diagonal symmetry of these two functions; viz, $\ln(e^x) = x$ and $e^{(\ln x)} = x$.

At $\sigma = \frac{1}{2}$, DSPL =

$$(6) \quad \frac{1}{2^{\frac{1}{2}}} \left(t^2 + \frac{1}{4} \right)^{\frac{1}{2}} \left[(2n)^{\frac{1}{2}} \cos(t \ln(2n)) - \frac{1}{4} \pi - (2n-1)^{\frac{1}{2}} \cos(t \ln(2n-1)) - \frac{1}{4} \pi + C \right]_1^{\infty}$$

$F(n)$ Dirichlet Sigma-Power Law, denoted by DSPL, refers to $\int \text{sim-}\eta(s) \text{dn}$. Eq. (6) is $F(n)$ DSPL at $\sigma = \frac{1}{2}$ that will incorporate all nontrivial zeros [as Pseudo-zeroes to Zeroes conversion].

Remark 2.2. Given $\delta = \frac{1}{10}$, the left-shifted $\sigma = \frac{1}{2} - \delta = \frac{2}{5}$ -non-critical line (Figure 5) and right-shifted $\sigma = \frac{1}{2} + \delta = \frac{3}{5}$ -non-critical line (Figure 6) are **equidistant** from nil-shifted $\sigma = \frac{1}{2}$ -critical line (Figure 4). Let $x = (2n)$ or $\frac{1}{(2n)}$ or $(2n-1)$ or $\frac{1}{(2n-1)}$. With multiplicative inverse operation of $x^\delta \cdot x^{-\delta} = 1$ or $\frac{1}{x^\delta} \cdot \frac{1}{x^{-\delta}} = 1$ that is applicable, this imply intrinsic presence of **Multiplicative Inverse** in $\text{sim-}\eta(s)$ or DSPL for all σ values with this function or law rigidly obeying relevant trigonometric identity. Then both $f(n) \text{sim-}\eta(s)$ and $F(n)$ DSPL will manifest **Principle of Equidistant for Multiplicative Inverse** (as per Page 41 of [8]).

The dissertation based on Figure 8 with inverse functions $\ln(x)$ & $e(x)$ in Page 30 – 35 of [8] confirms Asymptotic law of distribution for prime numbers as $\lim_{x \rightarrow \infty} \frac{\text{Prime-}\pi(x)}{\left[\frac{x}{\ln(x)} \right]} = 1$ and Asymptotic law of distribution for composite numbers as $\lim_{x \rightarrow \infty} \frac{\text{Composite-}\pi(x)}{\left[\frac{x}{e(x)} \right]} = 1$. This fully supports Prime number theorem

[viz, $\text{Prime-}\pi(x) \approx \frac{x}{\ln(x)}$] & derived Composite number theorem [viz, $\text{Composite-}\pi(x) \approx \frac{x}{e(x)}$].

A number base, consisting of any whole number greater than 0, is number of digits or combination of digits that a number system uses to represent numbers e.g. decimal number system or base 10, binary number system or base 2, octal number system or base 8, hexa-decimal number system or base 16. Prime counting function, $\text{Prime-}\pi(x)$ = number of primes $\leq x$ and Composite counting function, $\text{Composite-}\pi(x)$ = number of composites $\leq x$. As $x \rightarrow \infty$, derived properties of $\text{Prime-}\pi(x)$ occur in, for instance, Prime number theorem for Arithmetic Progressions, $\text{Prime-}\pi(x; b, a)$ [= number of primes $\leq x$ with last

digit of primes given by a in base b]. For any choice of digit a in base b with $\gcd(a,b) = 1$: Prime- $\pi(x; b, a) \sim \frac{\text{Prime-}\pi(x)}{\phi(b)}$. Here, Euler's totient function $\phi(n)$ is defined as the number of positive integers $\leq n$ that are relatively prime to (i.e., do not contain any factor in common with) n , where 1 is counted as being relatively prime to all numbers. Then each of the last digit of primes given by digit a in base b as $x \rightarrow \infty$ is equally distributed between the permitted choices for digit a with this result being valid for, and is independent of, any chosen base b .

Numbers with their last digit ending in (i) 1, 3, 7 or 9 [which can be either primes or composites] constitute $\sim 40\%$ of all integers; and (ii) 0, 2, 4, 5, 6 or 8 [which must be composites] constitute $\sim 60\%$ of all integers. We validly ignore the only single-digit even prime number 2 and odd prime number 5. We note ≥ 2 -digit Odd Primes can only have their last digit ending in 1, 3, 7 or 9 but not in 0, 2, 4, 5, 6 or 8. These are given as the **complete List**:

The last digit of Odd Primes having their Prime gaps with last digit ending in 2 [viz, Gap 2, Gap 12, Gap 22, Gap 32...] can only be 1, 3 or 9 [but not (5) or 7] as three choices.

The last digit of Odd Primes having their Prime gaps with last digit ending in 4 [viz, Gap 4, Gap 14, Gap 24, Gap 34...] can only be 1, 3 or 7 [but not (5) or 9] as three choices.

The last digit of Odd Primes having their Prime gaps with last digit ending in 6 [viz, Gap 6, Gap 16, Gap 26, Gap 36...] can only be 3, 7 or 9 [but not (5) or 1] as three choices.

The last digit of Odd Primes having their Prime gaps with last digit ending in 8 [viz, Gap 8, Gap 18, Gap 28, Gap 38...] can only be 1, 7 or 9 [but not (5) or 3] as three choices.

The last digit of Odd Primes having their Prime gaps with last digit ending in 0 [viz, Gap 10, Gap 20, Gap 30, Gap 40...] can only be 1, 3, 7 or 9 [but not (5)] as four choices.

AXIOM 1. *Applications of the Prime number theorem for Arithmetic Progressions will confirm Modified Polignac's and Twin prime conjectures to be true (as per Page 31 – 32 in [8]).*

Proof. We use decimal number system (base $b = 10$), and ignore the only single-digit even prime number 2 and odd prime number 5. For $i = 1, 2, 3, 4, 5, \dots$; the last digit of all Gap $2i$ -Odd Primes can only end in 1, 3, 7 or 9 that are each proportionally and equally distributed as $\sim 25\%$ when $x \rightarrow \infty$, whereby this result is consistent with Prime number theorem for Arithmetic Progressions. The 100%-Set of, and its derived four unique 25%-Subsets of, Gap $2i$ -Odd Primes based on their last digit being 1, 3, 7 or 9 must all be CIS-ALN-decelerating. "Different Prime numbers literally equates to different Prime gaps" is a well-known intrinsic property. Since the ALN of Gap $2i$ as fully represented by all Prime gaps with last digit ending in 0, 2, 4, 6 or 8 are associated with various permitted combinations of last digit in Gap $2i$ -Odd Primes being 1, 3, 7 and/or 9 as three or four choices [outlined above in **List** from preceding paragraph]; then these ALN unique subsets of Prime gaps based on their last digit being 0, 2, 4, 6 or 8 together with their correspondingly derived ALN unique subsets constituted by Gap $2i$ -Odd Primes having last digit 1, 3, 7 or 9 must also all be CIS-ALN-decelerating. The Probability (any Gap $2i$ abruptly terminating as $x \rightarrow \infty$) = Probability (any Gap $2i$ -Odd Primes abruptly terminating as $x \rightarrow \infty$) = 0. Thus Modified Polignac's and

Twin prime conjectures is confirmed to be true. With ordinary Riemann hypothesis being a special case, we also note the generalized Riemann hypothesis formulated for Dirichlet L-function holds once $x > b^2$, or base $b < x^{\frac{1}{2}}$ as $x \rightarrow \infty$. *The ["statistical" or "probabilistic"] proof is now complete for Axiom 1* \square .

3. Generic Squeeze theorem as a novel mathematical tool in Number theory

We adopt abbreviations \mathbb{P} = Prime numbers, \mathbb{C} = Composite numbers, NTZ = nontrivial zeros, $G[y=0]P$ = Gram[$y=0$] points (usual / traditional Gram points), and $G[x=0]P$ = Gram[$x=0$] points.

Gram's Law and Rosser's Rule for Riemann zeta function via its proxy Dirichlet eta function at $\sigma = \frac{1}{2}$ are perpetually associated with recurring violations (failures). Violations of Gram's Law equates to intermittently observing various geometric variants of two consecutive (positive first and then negative) $G[y=0]P$ that is alternatingly followed by two consecutive NTZ. Violations of Rosser's Rule equates to intermittently observing various geometric variants of reduction in expected number of certain x-axis intercept points. Both types of violations may give rise to intermittent or cyclical events of two missing $G[y=0]P$ or, equivalently, to two extra NTZ.

We hereby artificially and conveniently regard the $G[y=0]P \leq G[x=0]P \leq \text{NTZ}$ inequality as being applicable for Theorem 3.1 below. Observe that this particular inequality has never been definitively confirmed to be true *over the large range of numbers*. With full analysis, one of the following alternative inequalities $G[x=0]P \leq G[y=0]P \leq \text{NTZ}$ or $\text{NTZ} \leq G[y=0]P \leq G[x=0]P$ or $\text{NTZ} \leq G[x=0]P \leq G[y=0]P$ or $G[x=0]P \leq \text{NTZ} \leq G[y=0]P$ or $G[y=0]P \leq \text{NTZ} \leq G[x=0]P$ *over the large range of numbers* could instead be true. Even the equality $G[y=0]P = G[x=0]P = \text{NTZ}$ *over the large range of numbers* could instead also be true. It may even be the case that all types of inequalities mentioned above could cyclically co-exist *over the large range of numbers*. In principle, Theorem 3.1 should intuitively be validly applicable to the correctly chosen inequality [or equality].

THEOREM 3.1. *(Generic Squeeze theorem). Crucially applicable to all prime numbers, composite numbers and nontrivial zeros, our devised Theorem 3.1 is formally stated as follows (as per Page 51 – 53 in [8]).*

Let I be an interval containing point a . Let g , f , and h be algorithms or functions defined on I , except possibly at a itself. Suppose for every x in I not equal to a , we have $g(x) \leq f(x) \leq h(x)$ and also suppose $\lim_{x \rightarrow a} g(x) = \lim_{x \rightarrow a} h(x) = L$. Then $\lim_{x \rightarrow a} f(x) = L$. The algorithms or functions g and h are said to be lower and upper bounds (respectively) of f . Here, a is not required to lie in the interior of I . Indeed, if a is an endpoint of I , then the above limits are left- or right-hand limits. A similar statement holds for infinite intervals e.g. applicable to the IM t -valued NTZ (as CIS-IM-linear) obtained from Riemann zeta function via its proxy Dirichlet eta function, and the ALN of \mathbb{P} (as CIS-ALN-decelerating) obtained from Sieve-of-Eratosthenes and IM \mathbb{C} (as CIS-IM-accelerating) obtained from Complement-Sieve-of-Eratosthenes. In particular, if $I = (0, \infty)$ or $(0, \text{ALN})$, then the conclusion holds, taking the limits as $x \rightarrow \infty$ or ALN.

Let a_n, c_n be two sequences converging to ℓ , and b_n a sequence. If $\forall n \geq N, N \in \mathbb{N}$ we have $a_n \leq b_n \leq c_n$, then b_n also converges to ℓ . From above arguments, we logically notice Generic Squeeze theorem is valid for carefully selected sequences e.g. those precisely derived from algorithm Sieve-of-Eratosthenes generating set of all unique \mathbb{P} 2, 3, 5, 7, 11, 13, 17, 19, 23, 29... with progressive "cumulative" cardinality $\equiv c_n$ and sub-algorithms from Complement-Sieve-of-Eratosthenes generating two subsets of all unique pre-prime-Gap 2-Even \mathbb{C} 4, 6, 10, 12, 16, 18, 22, 28... with progressive "cumulative" cardinality $\equiv b_n$ and of all unique 1st post-prime-Gap 1-Even \mathbb{C} 8, 14, 20, 24, 32, 38, 44... with progressive "cumulative" cardinality $\equiv a_n$. We recognize even \mathbb{P} 2 is not a pre-prime-Gap 2-Even \mathbb{C} , and 1st \mathbb{P} 3, 5, 11, 17, 29, 41, 59... from all twin prime pairings (3, 5), (5, 7), (11, 13), (17, 19), (29, 31), (41, 43), (59, 61)... are never associated with 1st post-prime-Gap 1-Even \mathbb{C} as these even numbers 4, 6, 12, 18, 30, 42, 60... [which must be ***eternally ubiquitous***, not least, to comply with Law of Continuity] are all pre-prime-Gap 2-Even \mathbb{C} . Incorporating mixtures of \mathbb{P} & \mathbb{C} , our findings on twin prime pairings $\implies \{c_n \text{ representing progressive total of all } \mathbb{P}\} > \{b_n \text{ representing progressive total of all pre-prime-Gap 2-Even } \mathbb{C}\} > \{a_n \text{ representing progressive total of all 1}^{st} \text{ post-prime-Gap 1-Even } \mathbb{C}\}$. Since $\lim_{n \rightarrow ALN} a_n = \lim_{n \rightarrow ALN} c_n = \text{CIS-ALN-decelerating}$, then $\lim_{n \rightarrow ALN} b_n = \text{CIS-ALN-decelerating}$. Stated in another insightful way: The perpetual recurrence of intermittent inevitable DISAPPEARANCE of 1st post-prime-Gap 1-Even \mathbb{C} is solely due to coinciding intermittent inevitable APPEARANCE of twin primes \implies Twin prime conjecture is true.

*The 1st post-prime-Gap 1-Even \mathbb{C} precisely forms OEIS sequence A014574 *Average of twin prime pairs* 4, 6, 12, 18, 30, 42, 60, 72, 102, 108, 138, 150, 180, 192, 198, 228, 240, 270, 282, 312, 348, 420, 432, 462, 522, 570, 600, 618... by R. K. Guy, N. J. A. Sloane & E. W. Weisstein (June 11, 2011) <https://oeis.org/A014574> whereby

(i) With an initial 1 added, these numbers form part of the complement of closure of $\{2\}$ under the operations $a * b + 1$ and $a * b - 1$ within the set of all non-zero positive even numbers $U = \{2, 4, 6, 8, 10, \dots\}$. For $a * b + 1$: $2 * 2 + 1 = 5$. For $a * b - 1$: $2 * 2 - 1 = 3$. Under both operations, we obtain the set $S = \{2, 3, 5\}$. Therefore the complement of S within U would be all even numbers except 2 [and 5 & 3]; viz, $S' = \{4, 6, 8, 10, 12, 14, 16, \dots\}$.

(ii) These numbers are also the square root of the product of twin prime pairs + 1. Two consecutive odd numbers can be written as $2k + 1, 2k + 3$. Then $(2k + 1)(2k + 3) + 1 = 4(k^2 + 2k + 1) = 4(k + 1)^2$, a perfect square [where the countably infinite set of all perfect squares \equiv product of an integer multiplied by itself = 1, 4, 9, 16, 25, 36, 49, 64, 81, 100...]. Since twin prime pairs are two consecutive odd numbers, the statement is true for all CIS-ALN-decelerating twin prime pairs.

(iii) These numbers are single (or isolated) composites. Nonprimes k such that neither $k - 1$ nor $k + 1$ is nonprime.

(iv) These form the numbers n such that $\sigma(n - 1) = \phi(n + 1)$. This equation involves two arithmetic functions: the sum of divisors function σ [which calculates the sum of all positive divisors of n e.g. when $n = 30$: Prime factorization of $(n - 1) = 29$ is $29 = 29^1$, and $\sigma(29) = 1 + 29 = 30$] and Euler's totient function ϕ [which gives the count of positive integers less than n that are coprime to n e.g. Prime factorization of $(n + 1) = 31$ is $31 = 31^1$, and $\phi(31) = 31 - 1 = 30$].

(v) Aside from first term 4 in the sequence, all remaining terms 6, 12, 18, 30, 42, 60, 72, 102, 108, 138, 150... have digital root 3, 6, or 9 e.g. digital root of 138 is 3 since $138 = 1 + 3 + 8 = 12$ and $1 + 2 = 3$.

(vi) These form the numbers n such that $n^2 - 1$ is a semiprime [a natural number that is the product of two prime numbers].

(vii) Every term but the first term 4 is a multiple of 6 [and all the multiple of 6 clearly constitute a countably infinite set].

From above synopsis that is valid for [mixed] prime & composite numbers as $x \rightarrow \text{ALN}$, we conclude: Since there is an ALN of all prime numbers as (c_n) and also an ALN of all 1st post-prime-Gap 1-Even composite numbers as (a_n) , then by the Generic Squeeze theorem, there must also be an ALN of all Gap 2-Even composite numbers as (b_n) . Thus ℓ must have the value of ALN. In theory, even if there are [incorrectly] only finitely many twin primes, the mathematical relationship of $a_n \leq b_n \leq c_n$ will still hold except that the Generic Squeeze theorem is no longer applicable as there will be inevitable "errors" present in the computed a_n, b_n and c_n .

By applying Generic Squeeze theorem [only] to Odd \mathbb{P} , we now prove Polignac's and Twin prime conjectures are true: We ignore even \mathbb{P} 2. Let algorithm Sieve-of-Eratosthenes that generate the set of all unique Total Odd \mathbb{P} 3, 5, 7, 11, 13, 17, 19, 23, 29... with progressive "cumulative" cardinality $\equiv c_n$ and sub-algorithms from Sieve-of-Eratosthenes that generate the two [randomly selected] subsets of all unique Gap 4-Odd \mathbb{P} 7, 13, 19, 37, 43, 67... with progressive "cumulative" cardinality $\equiv a_n$ and of all unique Gap 2, 6, 8, 10, 12...-Odd \mathbb{P} 3, 5, 11, 17, 23, 23, 29, 31, 41, 47, 53, 59, 61... [viz, not including Gap 4-Odd \mathbb{P}] with progressive "cumulative" cardinality $\equiv b_n$. Instead of choosing b_n to be even Prime gap 4, one could choose any other eligible even Prime gap derived from the set of all even Prime gaps [which will inevitably also include Zhang's landmark result of an unknown even Prime gap $N < 70$ million]. Since $\lim_{n \rightarrow \text{ALN}} a_n = \lim_{n \rightarrow \text{ALN}} c_n = \text{CIS-ALN-decelerating}$, then $\lim_{n \rightarrow \text{ALN}} b_n = \text{CIS-ALN-decelerating}$. Stated in another insightful way: In order for novel method Generic Squeeze theorem to be ubiquitously applicable for Odd \mathbb{P} , all even Prime gaps 2, 4, 6, 8, 10... must be associated with corresponding ALN of Odd \mathbb{P} .

On 17 April 2013, Yitang Zhang announced an incredible proof that there are infinitely many pairs of prime numbers that differ by less than 70 million[9]; viz, there is an arbitrarily large number of Odd Primes with an unknown even Prime gap N of less than 70 million. By optimizing Zhang's bound, subsequent Polymath Project collaborative efforts using a new refinement of GPY sieve in 2014 lowered N to 246; and assuming Elliott-Halberstam conjecture and its generalized form further lower N to 12 and 6, respectively. Intuitively, N has more than one valid values such that the same condition holds for each N value. Using different methods, we can at most lower N to 2 and 4 in regards to Odd Primes having small even Prime gaps 2 & 4 with each uniquely generating CIS-ALN-decelerating Odd Primes. We anticipate there are all remaining even Prime gaps w.r.t. Odd Primes with large even Prime gaps ≥ 6 as denoted by corresponding $N \geq 6$ values whereby each large even Prime gap generates its unique CIS-ALN-decelerating Odd Primes.

We justify "**Zhang's optimized result ≥ 3 up to ALN even Prime gaps with each having ALN of elements**": Always as finite [but NOT infinite] length, we observe as side note that two or more consecutive

Odd Primes are validly and rarely constituted by [same] even Prime gap of 6 or multiples of 6. With just one or two existing even Prime gaps that have ALN of elements being simply "insufficient" in the large range of prime numbers, then the landmark result by Zhang on this unknown even Prime gap N of less than 70 million is usefully extrapolated as "There must be at least one subset of Odd Primes having ALN of elements". Hence there are aesthetically at least two, if not three, existing even Prime gaps generating corresponding CIS-ALN-decelerating Odd Primes. Modified Polignac's and Twin prime conjectures equates to all even Prime gaps 2, 4, 6, 8, 10... generating corresponding CIS-ALN-decelerating Odd Primes.

Near-identical arguments can be made for three types of Gram points located at $\sigma = \frac{1}{2}$ -critical line of Riemann zeta function but we leave out the full exercise of applying Generic Squeeze theorem to NTZ as progressive "cumulative" cardinality $\equiv c_n$, $G[x=0]P$ as progressive "cumulative" cardinality $\equiv b_n$ and $G[y=0]P$ as progressive "cumulative" cardinality $\equiv a_n$. We immediately recognize the [trivial] conclusion: Since $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} c_n = \text{CIS-IM-linear}$, then $\lim_{n \rightarrow \infty} b_n = \text{CIS-IM-linear}$.

Eq. (4) manifests *exact* Dimensional analysis homogeneity when $\sigma = \frac{1}{2}$ whereby $\Sigma(\text{all fractional exponents}) = 2(-\sigma) = \text{exact negative whole number } -1$ [c.f. Eq. (5) manifests *inexact* Dimensional analysis homogeneity when $\sigma = \frac{2}{3}$ whereby $\Sigma(\text{all fractional exponents}) = 2(-\sigma) = \text{inexact negative fractional number } -\frac{4}{3}$]. Only Dirichlet eta function having parameter $\sigma = \frac{1}{2}$ will mathematically depict [optimal] "formula symmetry" on $\Sigma(\text{all fractional exponents})$ as an exact negative whole number, whereby absolute values of all fractional exponents $= \frac{1}{2}$ when associated with constant 2 and variable $(2n)$ or $(2n-1)$. This formula symmetry is not equivalent to geometrical symmetry about X-axis, Y-axis, Diagonal, or Origin point that do not exist for any Dirichlet eta function when considered for either $-\infty < t < 0$ or $0 < t < +\infty$ from full range $-\infty < t < +\infty$; whereby we conventionally adopt the positive range. Simple observation of [optimal] "formula symmetry" implies only $\sigma = \frac{1}{2}$ -Dirichlet eta function will perpetually & geometrically intercept $\sigma = \frac{1}{2}$ -Origin point as Origin intercept points or Gram $[x=0, y=0]$ points (i.e. will perpetually & mathematically lie on $\sigma = \frac{1}{2}$ -critical line as nontrivial zeros) an infinite number of times.

Conforming to Langlands program "**Theory of Symmetry**", IL (sub-)algorithms or IL (sub-)equations and FL (sub-)algorithms or FL (sub-)equations will respectively generate infinitely-many and finitely-many entities. All the FL (sub-)algorithms or FL (sub-)equations are CP but the IL (sub-)algorithms or IL (sub-)equations can be either CP or IP. Here, we validly regard equation Dirichlet eta function (*proxy* for Riemann zeta function that generate nontrivial zeros when $\sigma = \frac{1}{2}$), and algorithms Sieve-of-Eratosthenes [for prime numbers] and Complement-Sieve-of-Eratosthenes [for composite numbers] as **non-overlapping** "IP IL number generators".

Remark 3.1. Not least to maintain Dimensional analysis homogeneity and to conserve Total number of elements (cardinality), it is a crucial ***sine qua non*** *Pre-requisite Mathematical Condition that a parent IP IL algorithm which is precisely constituted by its IP IL sub-algorithms or a parent IP IL equation which is precisely constituted by its IP IL sub-equations must generally all be wholly IP IL [and not be mixed IP IL and CP FL]*. Useful self-explanatory analogy using CP IL (sub)algorithms or (sub)equations: Set "twin" even numbers 0, 2, 4, 6, 8, 10... with Even gap 2, Subset "cousin" even numbers 0, 4, 8, 12, 16,

20... with Even gap 4, Subset "sexy" even numbers 0, 6, 12, 18, 24, 30... with Even gap 6, etc must all be constituted by CP IL [and not mixed CP IL and IP IL] even numbers that are derived from, paradoxically, **overlapping** "CP IL number generators".

Remark 3.2. It was correctly asserted on Page 3 – 4 of [8] that any created Prime-tuplet or Prime-tuple is not able to be used to either prove or disprove Modified Polignac's and Twin prime conjectures. The main reason is Prime-tuplets or Prime-tuples are **"overlapping and incomplete"** (Sub)Tuples Classification of consecutive primes. In contrast, we can use **"non-overlapping and complete"** (Sub)Sets Classification of grouped primes to prove these conjectures. Thus even Prime gap 2 = Prime 2-tuplets of diameter 2 and even Prime gaps 4, 6, 8, 10, 12... = Prime 2-tuples of diameter 4, 6, 8, 10, 12....

4. Applying the Theorem of Divergent-to-Convergent series conversion for Prime numbers to Polignac's and Twin prime conjectures

Recall from section 2 the algorithms Sieve-of-Eratosthenes (S-of-E) and Modified-S-of-E. Both algorithms and their derived sub-algorithms faithfully generate set of all prime numbers 2, 3, 5, 7, 11, 13...; set of all Odd Primes 3, 5, 7, 11, 13, 17...; and subsets of Odd Primes derived from even Prime gaps 2, 4, 6, 8, 10.... By performing summation given by $\sum_{n=i}^{ALN} p_{n+1} = 2 + \sum_{i=1}^n g_i$ and $\sum_{n=i}^{ALN} p_{n+1} = 3 + \sum_{i=2}^n g_i$, we obtain (de novo) infinite series as diverging series for these two algorithms [and their derived sub-algorithms]. For Polignac's and Twin prime conjectures to be true, we deduce the cardinality for (i) set of all prime numbers, (ii) set of all Odd Primes, (iii) subsets of Odd Primes, and (iv) set of all even Prime gaps must all be CIS-ALN-decelerating. In contrast, we deduce below after Theorem 4.1 that all Brun's constants as (derived) infinite series are, in fact, converging series.

Helpful preliminary information about Theorem 4.1: Four basic arithmetic operations of addition [and complementary subtraction] and multiplication [and complementary division] obey Axioms of Addition and Multiplication, and Axioms of Order. Division of one number by another is equivalent to multiplying first number by reciprocal (or multiplicative inverse) of second number, whereby division by 0 is always undefined. Subtraction of one number from another is equivalent to adding additive inverse of second number (viz, a negative number) to first number (viz, a positive number). Below are Completely Predictable properties arising from (non-)alternating addition of any Even numbers (\mathbb{E}) 0, 2, 4, 6, 8, 10, 12... and any Odd numbers (\mathbb{O}) 1, 3, 5, 7, 9, 11, 13...:

(1) $\mathbb{E} + \mathbb{E} + \mathbb{E} + \mathbb{E}...$ when involving any number of terms = \mathbb{E} .

(2) $\mathbb{O} + \mathbb{O} + \mathbb{O} + \mathbb{O}...$ when involving an even number of terms = \mathbb{E} ; and when involving an odd number of terms = \mathbb{O} .

The alternating sum $\mathbb{E} + \mathbb{O} + \mathbb{E} + \mathbb{O} + \mathbb{E} + \mathbb{O}...$ when involving $(1 + n)$ terms for $n = 1, 2, 3, 4, 5...$ = repeating patterns of $\mathbb{O}, \mathbb{O}, \mathbb{E}, \mathbb{E}, \mathbb{O}, \mathbb{O}, \dots$

A convergent series (CS) as an infinite series having its partial sums of sequence that tends to a finite limit is validly represented by the [defined] value of this finite limit. A divergent series (DS) as an infinite series

having its partial sums of sequence that tends to a infinite limit is validly represented by the [undefined] value of this infinite limit. As previously discussed in section 2, the infinite-length sequence of a given CS or DS can theoretically be constituted by either positive terms OR alternating positive and negative terms. The following are Completely Predictable properties arising from addition of any infinite series constituted by ≥ 1 CS and/or ≥ 1 DS:

- I. DS+DS+DS+... when involving any number of DS terms = DS.
- II. CS+CS+...+DS+DS+... when involving any number of CS terms & any number of DS terms = DS.
- III. CS+CS+CS+... when involving a finite number of CS terms = CS.
- IV. CS+CS+CS+... when involving an infinite number of CS terms or arbitrarily large number (ALN) of CS terms = DS.

THEOREM 4.1. (*Theorem of Divergent-to-Convergent series conversion for Prime numbers*) (as per Page 53 – 54 in [8]).

We validly ignore even prime number 2. Theorem 4.1, aka **Smoothed asymptotics for Prime numbers with an enhanced regulator**, as given in next two paragraphs is further expanded below using three Brun's constants computed for twin primes, cousin primes and sexy primes.

For [eligible] homogenous entities of prime numbers with application of divergent series (DS) to convergent series (CS) conversion relationship, we obtain CS + CS + CS +... when involving arbitrarily large number (ALN) of CS terms [that faithfully "represent" all Subsets of Odd Primes] = DS [that faithfully "represent" the Set of all Odd Primes]. We recognize the ALN of computed CS terms will precisely correspond to Brun's constants. The correctly chosen **enhanced regulator for prime numbers** \equiv *sine qua non* condition [that must be fully complied with by all Odd Primes]: *Derived from the set of all Odd Primes, there must be an ALN of subsets of Odd Primes derived from even Prime gaps 2, 4, 6, 8, 10... with each subset of Odd Primes containing an ALN of unique elements.*

The elimination of a DS to CS under our novel *Divergent-to-Convergent series theorem for Prime numbers* fully supports Polignac's and Twin prime conjectures to be true. As alluded to in section 1, this procedure is reminiscent of invoking 'Method of Smooth asymptotics' and 'regularization of divergent series or integrals' to enable elimination of divergences in analytic number theory and preservation of gauge invariance at one loop in a wide class of non-abelian gauge theories coupled to Dirac fermions that preserves Ward identity for vacuum polarisation tensor [viz, a regularized quantum field theory]. This is achieved by Padilla and Smith via adopting suitable choices from their proposed families of enhanced regulators[4] with analytic continuation that converge to Riemann zeta function value $\zeta(-1) = -\frac{1}{12}$ of extra relevance to quantum gravity, string theory, etc.

Considering Euler products $\sum_{n=1}^{\infty} \frac{1}{n} = \prod_p \frac{1}{1-p^{-1}}$ when taken over the set of all infinitely many primes, Leonhard Euler in 1737 showed the [harmonic] **infinite series** of all infinitely many primes (as sum of the reciprocals of all infinitely many primes) **diverges** very slowly; viz, $\sum_{p \text{ prime}} \frac{1}{p} = \frac{1}{2} + \frac{1}{3} + \frac{1}{5} + \frac{1}{7} + \frac{1}{11} + \frac{1}{13}$

$+\frac{1}{17} + \frac{1}{19} + \dots = \infty$. If it were the case that this sum of the reciprocals of twin primes (Prime gap 2), cousin primes (Prime gap 4), sexy primes (Prime gap 6), etc all diverged; then that fact would imply that there are infinitely many of twin primes, cousin primes, sexy primes, etc. However twin primes are less frequent than all infinitely many prime numbers by nearly a logarithmic factor with this bound giving the intuition that the sum of the reciprocals of twin primes **converges** very slowly, or stated in other words, twin primes form a small set. The sum $\sum_{p: p+2 \in \mathbb{P}} \left(\frac{1}{p} + \frac{1}{p+2}\right) = \left(\frac{1}{3} + \frac{1}{5}\right) + \left(\frac{1}{5} + \frac{1}{7}\right) + \left(\frac{1}{11} + \frac{1}{13}\right) + \left(\frac{1}{17} + \frac{1}{19}\right) + \dots = 1.902160583104\dots$ in explicit terms either has finitely many terms or has infinitely many terms but is very slowly convergent with its value known as Brun's constant for (consecutive) twin primes. Similar deductive arguments can be developed for the sum of the reciprocals of cousin primes, sexy primes, etc that also **converges** very slowly with their associated Brun's constant for (consecutive) cousin primes [≈ 1.19705479], (consecutive) sexy primes [≈ 1.13583508], etc. All these heuristically computed Brun's constants are irrational (transcendental) numbers ONLY IF there are infinitely many twin primes, cousin primes, sexy primes, etc. Based on Zhang's result[9], there must be at least one computed Brun's constant that is irrational (transcendental) associated with infinitely many Odd Primes having an even Prime gap < 70 million. We ignore solitary even prime number 2. We use "Arbitrarily Large Number" to denote "infinitely many". As an absolutely indispensable condition, there are ALN of subsets of Odd Primes with each subset of Odd Primes containing ALN of elements – this is akin to choosing the correct "enhanced regulator". From above discussions, we heuristically deduce very slowly **diverging** sum (series) of the reciprocals of all ALN Odd Primes are fully constituted by very slowly **converging** sum (series) of the reciprocals of ALN Odd Primes derived from each and every subsets of Odd Primes.

Erdos primitive set conjecture, now proven as a theorem by Prof. Jared Lichtman[2], is the assertion that for any primitive set S with exactly k prime factors (with repetition), $\sum_{n \in S} \frac{1}{n \log n} \leq \sum_p \frac{1}{p \log p} = \frac{1}{2 \log 2} + \frac{1}{3 \log 3} + \frac{1}{5 \log 5} + \frac{1}{7 \log 7} + \frac{1}{11 \log 11} + \dots = 1.6366\dots$ [as a very slowly converging sum when $k = 1$ over infinitely-many integers 1, 2, 3, 4, 5...] $\implies f_k$ is maximized by the prime sum $f_1 = \sum_p \frac{1}{p \log p} = 1.6366\dots$ [representing the unique "largest" primitive set that ONLY contains all infinitely-many prime numbers 2, 3, 5, 7, 11, 13...]. As supporting Modified Polignac's and Twin prime conjectures to be true [with **all Odd Primes belonging to CIS-ALN-decelerating**]; one can calculate the equivalent $f_1 = \sum_p \frac{1}{p \log p}$ [also as very slowly converging sums with values $< 1.6366\dots$] for individual subsets of Odd Primes obtained from even Prime gaps 2, 4, 6, 8, 10... and notice these [derived] "**infinite series**" calculations must all, in principle and in synchrony, incorporate corresponding **CIS-ALN-decelerating Odd Primes from each subset**. This last statement is heavily supported by Yitang Zhang's result[9] which can be extrapolated as "There must be at least one subset of Odd Primes [obtained from an even Prime gap < 70 million] having infinitely many elements".

5. Subtypes of Countably Infinite Sets with Incompletely Predictable entities from Riemann zeta function and Sieve of Eratosthenes

The sets of numbers generated using power (exponent) such as 2 or $\frac{1}{2}$, even numbers, odd numbers, etc are morphologically constituted by *Completely Predictable (CP) numbers* in the sense that these sets of numbers are actually *not* random and do not behave like one. The sets of nontrivial zeros, primes, composites, etc are morphologically constituted by *Incompletely Predictable (IP) numbers* [or *pseudo-random numbers*] in the sense that these sets of numbers are actually *not* random but behave like one; thus giving rise to so-called "Mathematics for Incompletely Predictable Problems". The word *number* [singular noun] or *numbers* [plural noun] in reference to CP even and odd numbers, IP prime and composite numbers, IP Gram points and virtual Gram points can be interchanged with the word *entity* [singular noun] or *entities* [plural noun].

LEMMA 5.1. *We can formally define the elements from (sub)sets and (sub)tuples as Completely Predictable or Incompletely Predictable entities (as per Page 18 in [8]). Please also see Remark 3.1 & Remark 3.2 above in section 3 indicating the important significances arising from Lemma 5.1.*

Proof. A set is a collection of zero (viz, the empty set) or more elements (viz, a finite set with a finite number of elements or an infinite set with an infinite number of elements). A singleton refers to a finite set with a single element. A set can be any kind of mathematical objects: numbers, symbols, points in space, lines, other geometrical shapes, variables, or even other sets whereby these [mutable] non-repeating elements are not arranged in an unique order. A subset can be a [smaller] finite set derived from its [larger] parent finite set or its [larger] parent infinite set. A subset can also be a [smaller] infinite set derived from its [larger] parent infinite set. A tuple, which can potentially be subdivided into subtuples, is a finite ordered list (sequence) of elements whereby these [immutable] non-repeating elements are arranged in an unique order. Thus a tuple or a subtuple is regarded as a special type of finite set with the extra imposed restriction. As shown below using worked examples:

CP simple equation or algorithm generates CP numbers e.g. even numbers 0, 2, 4, 6, 8, 10... or odd numbers 1, 3, 5, 7, 9, 11.... Thus a generated CP number is **locationally defined** as a number whose i^{th} position is *independently* determined by simple calculations without needing to know related positions of all preceding numbers – this is a **Universal Property**.

IP complex equation or algorithm generates IP numbers e.g. prime numbers 2, 3, 5, 7, 11, 13... or composite numbers 4, 6, 8, 9, 10, 12.... Thus a generated IP number is **locationally defined** as a number whose i^{th} position is *dependently* determined by complex calculations with needing to know related positions of all preceding numbers – this is a **Universal Property**.

We clearly note the elements in (sub)sets and (sub)tuples when generated by equations or algorithms will precisely constitute relevant entities or numbers of interest. *The proof is now complete for Lemma 5.1*□.

A formula for primes in Number theory is a formula generating all prime numbers 2, 3, 5, 7, 11, 13, 17, 19, 23... exactly and without exception. Computationally slow and inefficient formulas for calculating

primes exist e.g. 1964 Willans formula $p_n = 1 + \sum_{i=1}^{2^n} \left[\left(\frac{n}{\sum_{j=1}^i \left[\left(\cos \frac{(j-1)! + 1}{j} \pi \right)^2 \right] \right)^{1/n} \right]$ which is

based on Wilson's theorem $n + 1$ is prime iff $n! \equiv n \pmod{n + 1}$. Then critics may ask the question "For $n = 1, 2, 3, 4, 5, \dots$; does Willans formula that faithfully compute corresponding n^{th} prime number p_n for all infinitely-many primes contradict Sieve-of-Eratosthenes algorithm as being an Infinite Length (IL) and Incompletely Predictable (IP) algorithm?" The answer is categorically 'no' based on carefully analyzing this formula using following arguments [which lend further support to Polignac's and Twin prime conjectures being true]: Willans formula has two sub-components $\left[\left(\cos \frac{(j-1)! + 1}{j} \pi \right)^2 \right] = \begin{cases} 1 & \text{if } j \text{ is prime or } 1 \\ 0 & \text{if } j \text{ is composite} \end{cases}$

& $\sum_{j=1}^i \left[\left(\cos \frac{(j-1)! + 1}{j} \pi \right)^2 \right] = (\# \text{ primes } \leq i) + 1$. We recognize this second sub-component stipulating $(\# \text{ primes } \leq i) + 1$ meant the actual position of every n^{th} prime number will have to be fully and indirectly computed each time, thus confirming the infinitely-many prime numbers are IP and of IL. Note all [complementary] composite numbers 4, 6, 8, 9, 10, 12, 14, 15, 16, 18... are simply obtained by discarding all prime numbers from integers 2, 3, 4, 5, 6, 7, 8, 9, 10... whereby "special" integers 0 & 1 are neither prime nor composite. We ignore even prime number 2. Zhang's landmark result[9] states there are infinitely many Odd Primes having an even Prime gap < 70 million. One could extrapolate Zhang's result to: There must be at least two or three up to all even Prime gaps being each associated with infinitely many Odd Primes. All obtained consecutive Odd Primes p_n and p_{n+1} can have their calculated $p_{n+1} - p_n$ values grouped together as belonging to even Prime gaps 2, 4, 6, 8, 10... whereby when the Zhang's result is maximally extrapolated, Polignac's and Twin prime conjectures are supported to be true.

LEMMA 5.2. *We can validly classify countably infinite sets as accelerating, linear or decelerating subtypes (as per Page 18 – 19 in [8]).*

Proof. We provide the following required mathematical arguments.

Cardinality: With increasing size, arbitrary Set [or Subset] \mathbf{X} can be countably finite set (CFS), countably infinite set (CIS) or uncountably infinite set (UIS). Denoted as $\|\mathbf{X}\|$ in this paper, the cardinality of Set \mathbf{X} measures *number of elements* in Set \mathbf{X} . E.g., Set **negative Gram[y=0] point** as constituted by a [solitary] rational (\mathbb{Q}) t-value of 0 instead of a usual transcendental ($\mathbb{R} - \mathbb{A}$) t-value has CFS of negative Gram[y=0] point with this particular $\|\mathbf{negative\ Gram[y=0]\ point}\| = 1$, Set even Prime number (\mathbb{P}) has CFS of solitary even $\mathbb{P} 2$ with $\|\mathbf{even\ P}\| = 1$, Set Natural numbers (\mathbb{N}) has CIS of \mathbb{N} with $\|\mathbb{N}\| = \aleph_0$, and Set Real numbers (\mathbb{R}) has UIS of \mathbb{R} with $\|\mathbb{R}\| = \mathfrak{c}$ (cardinality of the continuum). Then with $\|\mathbf{CIS}\| = \aleph_0 =$ [countably] infinitely many elements; we provide a novel classification for CIS based on its number of elements (cardinality) manifesting linear, accelerating or decelerating property constituting three subtypes of CIS.

CIS-IM-accelerating: CIS with cardinality = $\|\mathbf{CIS-IM-accelerating}\| = \aleph_0$ -accelerating = [countably]

infinitely many elements that (overall) acceleratingly reach an *infinity value*. Examples: CP integers 0, 1, 4, 9, 16... generated by simple equation $y = x^2$ for $x = 0, 1, 2, 3, 4...$ and CP values obtained from natural exponential function $y = e(x)$; and IP composite numbers 4, 6, 8, 9, 10... faithfully generated by complex Complement-Sieve-of-Eratosthenes algorithm [which is equivalent to simply discarding 0, 1, and all generated prime numbers via Sieve-of-Eratosthenes algorithm from the set of integers 0, 1, 2, 3, 4, 5...].

CIS-IM-linear: CIS with cardinality = $\|\text{CIS-IM-linear}\| = \aleph_0$ -linear = [countably] infinitely many elements that (overall) linearly reach an *infinity value*. Examples: CP entities 0, 1, 2, 3, 4, 5... [representing all positive integer numbers] generated by simple equation $y = x$ for $x = 0, 1, 2, 3, 4...$; CP entities 0, 2, 4, 6, 8, 10... [representing all positive even numbers] generated by simple equation $y = 2x$ for $x = 0, 1, 2, 3, 4...$; CP entities 1, 3, 5, 7, 9, 11... [representing all positive odd numbers] generated by simple equation $y = 2x - 1$ for $x = 1, 2, 3, 4, 5...$; and IP nontrivial zeros, Gram[$y=0$] points and Gram[$x=0$] points (all given as $\mathbb{R} - \mathbb{A}$ t-values) generated from complex equation Riemann zeta function via its *proxy* Dirichlet eta function. These IP entities will inevitably manifest IP perpetual repeating violations (failures) in Gram's Law and Rosser's Rule occurring infinitely many times. E.g., the former give rise to Set **negative Gram[$y=0$] points** whereby CIS negative Gram[$y=0$] points is constituted by $\mathbb{R} - \mathbb{A}$ t-values classified as having $\|\text{negative Gram}[y=0] \text{ points}\| = \|\text{CIS-IM-linear}\| = \aleph_0$ -linear.

CIS-ALN-decelerating: CIS with cardinality = $\|\text{CIS-ALN-decelerating}\| = \aleph_0$ -decelerating = [countably] arbitrarily large number of elements that (overall) deceleratingly reach an *Arbitrarily Large Number* value. Examples: CP entities 0, 1, $\sqrt{2}$, $\sqrt{3}$, 2, $\sqrt{5}$... generated by simple equation $y = \sqrt{x}$ for $x = 0, 1, 2, 3, 4, 5...$ and CP values obtained from natural logarithm function $y = \ln(x)$; and IP prime numbers 2, 3, 5, 7, 11... faithfully generated by complex Sieve-of-Eratosthenes algorithm.

The proof is now complete for Lemma 5.2□.

6. Conclusions including applying infinitesimals to outputs from Sieve of Eratosthenes and Riemann zeta function

NOTE: Additional materials regarding L-functions of relevance to Riemann hypothesis, and Birch and Swinnerton-Dyer conjecture are supplied in Appendix A [containing important computations and an overall SUMMARY provided at the end of this Appendix].

Figure 1 [depicting positive & negative prime numbers and composite numbers] and Figure 2 [depicting the Co-linear Riemann zeta function for positive & negative range] will manifest perfect Mirror symmetry and fully comply with Law of Continuity. Valid comments: Whereas the continuous-like equation Riemann zeta function $\zeta(s)$ Eq. (1) [via *proxy* Dirichlet eta function $\eta(s)$ Eq. (2)] for $s = \sigma \pm t$ range that generate mutually exclusive CIS-IM-linear σ -valued co-lines be mathematically regarded as smoothly *continuous everywhere* thus obeying Law of continuity; so must the discrete-like algorithms Sieve-of-Eratosthenes and Complement-Sieve-of-Eratosthenes that generate mutually exclusive Primes and Composites be conceptually regarded as jaggedly *continuous everywhere* thus also obeying Law of continuity.

CIS-ALN-decelerating Primes and CIS-IM-accelerating Composites are dependent complementary entities. In $\zeta(s)$ Eq. (1), the equivalent Euler product formula with product over prime numbers [instead of summation over natural numbers] represents $\zeta(s) \implies$ all primes and, by default, [complementary] composites are intrinsically encoded in $\zeta(s)$. Since via analytic continuation, $\eta(s) = \frac{1}{\gamma} \cdot \zeta(s)$ [proxy function for $\zeta(s)$ in $0 < \sigma < 1$ - critical strip]; then all primes and, by default, [complementary] composites are also intrinsically encoded in $\eta(s)$ Eq. (2).

For $i = 1, 2, 3, 4, 5, \dots, n$ (Page 14 of [8]): Recurring *Accelerating primes as Prime gap_{i+2} – Prime gap_{i+1} > Prime gap_{i+1} – Prime gap_i*, *Decelerating primes as Prime gap_{i+2} – Prime gap_{i+1} < Prime gap_{i+1} – Prime gap_i* and *Steady primes as Prime gap_{i+2} – Prime gap_{i+1} = Prime gap_{i+1} – Prime gap_i* [≡ "**Alternating Prime Gaps series**" with Prime gaps alternatingly \uparrow & \downarrow] are computed by (sub-)algorithms to obtain mutually exclusive (solitary) even prime number 2 with odd Prime gap 1; odd Twin primes, odd Cousin primes & odd Sexy primes with even Prime gaps 2, 4 & 6.

(a) For IP IL algorithm [Gap 2, 4, 6, 8, 10...]-Sieve of Eratosthenes $p_{n+1} = 3 + \sum_{i=1}^n g_i$ [where $n = \text{ALN}$] that faithfully generates all Odd \mathbb{P} {3, 5, 7, 11, 13, 17, 19...} with cardinality \aleph_0 -decelerating, the n^{th} even Prime gap between two successive Odd \mathbb{P} is denoted by $g_n = (n + 1)^{\text{st}}$ Odd $\mathbb{P} - (n)^{\text{th}}$ Odd \mathbb{P} , i.e. $g_n = p_{n+1} - p_n = 2, 2, 4, 2, 4, 2, \dots$

(b) For CP FL sub-algorithm [Gap 1]-Sieve of Eratosthenes $p_{n+1} = 2 + \sum_{i=1}^n g_i$ [where $n = 1$ and not ALN] that faithfully generates the first and only Even \mathbb{P} {2} \equiv first and only paired Even \mathbb{P} {(2,3)} with cardinality CFS of 1, the solitary n^{th} odd prime gap between two successive primes is denoted by $g_n = (n + 1)^{\text{st}}$ Odd $\mathbb{P} - (n)^{\text{th}}$ Even \mathbb{P} , i.e. $g_n = p_{n+1} - p_n = 3 - 2 = 1$.

(c) For IP IL sub-algorithm [Gap 2]-Sieve of Eratosthenes $p_{n+1} = 3 + \sum_{i=1}^n g_i$ [where $n = \text{ALN}$] that faithfully generates all Odd twin \mathbb{P} {3, 5, 11, 17, 29, 41, 59...} \equiv all paired Odd twin \mathbb{P} {(3,5), (5,7), (11,13), (17,19), (29,31), (41,43), (59,61)...} with cardinality \aleph_0 -decelerating, the n^{th} even Prime gap between two successive Odd twin \mathbb{P} is denoted by $g_n = (n + 1)^{\text{st}}$ Odd twin $\mathbb{P} - (n)^{\text{th}}$ Odd twin \mathbb{P} , i.e. $g_n = p_{n+1} - p_n = 2, 6, 6, 12, 12, 18, \dots$

(d) For IP IL sub-algorithm [Gap 4]-Sieve of Eratosthenes $p_{n+1} = 7 + \sum_{i=1}^n g_i$ [where $n = \text{ALN}$] that faithfully generates all Odd cousin \mathbb{P} {7, 13, 19, 37, 43, 67...} \equiv all paired Odd cousin \mathbb{P} {(7,11), (13,17), (19,23), (37,41), (43,47), (67,71)...} with cardinality \aleph_0 -decelerating, the n^{th} even Prime gap between two successive Odd cousin \mathbb{P} is denoted by $g_n = (n + 1)^{\text{st}}$ Odd cousin $\mathbb{P} - (n)^{\text{th}}$ Odd cousin \mathbb{P} , i.e. $g_n = p_{n+1} - p_n = 6, 6, 8, 6, 24, \dots$

(e) For IP IL sub-algorithm [Gap 6]-Sieve of Eratosthenes $p_{n+1} = 23 + \sum_{i=1}^n g_i$ [where $n = \text{ALN}$] that faithfully generates all Odd sexy \mathbb{P} {23, 31, 47, 53, 61, 73, 83...} \equiv all paired Odd sexy \mathbb{P} {(23,29), (31,37), (47,53), (53,59), (61,67), (73,79), (83,89)...} with cardinality \aleph_0 -decelerating, the n^{th} even Prime gap between two successive Odd sexy \mathbb{P} is denoted by $g_n = (n + 1)^{\text{st}}$ Odd sexy $\mathbb{P} - (n)^{\text{th}}$ Odd sexy \mathbb{P} , i.e. $g_n = p_{n+1} - p_n = 8, 16, 6, 8, 12, 10, \dots$

With $n = \text{ALN}$ or, traditionally, ∞ ; rigorous algorithm-type proof for Modified Polignac's and Twin prime conjectures can be stated here as two statements. Statement 1: All known prime numbers = IP IL algorithm (a) + CP FL sub-algorithm (b). Statement 2: IP IL algorithm (a) = IP IL sub-algorithm (c) + IP IL sub-algorithm (d) + IP IL sub-algorithm (e) +... [that involves all even Prime gaps 2, 4, 6, 8, 10...].

As *proxy* function for Riemann zeta function in $0 < \sigma < 1$ critical strip, Dirichlet eta function when treated as equation and sub-equation at (unique) $\sigma = \frac{1}{2}$ -critical line will faithfully generate all x-axis intercept points as *usual* Gram points or Gram[y=0] points, all y-axis intercept points as Gram[x=0] points, and all Origin intercept points as Gram[x=0,y=0] points or nontrivial zeros. Confirming Riemann hypothesis to be true, these entities that constitute the three types of Gram points are mutually exclusive, dependent and endowed with t -valued irrational (transcendental) numbers except for initial Gram[y=0] point endowed with a t -valued rational number:

(a) Considered for $t = 0$ to $+\infty$ at $\sigma = \frac{1}{2}$, Dirichlet eta function as IP IL equation will faithfully generate all above-mentioned three types of Gram points that are endowed with t -valued irrational (transcendental) numbers except for first Gram[y=0] point.

(b) Considered only for $t = 0$ at $\sigma = \frac{1}{2}$, Dirichlet eta function as CP FL sub-equation will faithfully generate the first and only Gram[y=0] point that is endowed with t -valued rational number 0.

We analyze the data of all CIS-IM-linear computed nontrivial zeros (NTZ) when extrapolated out over a wide range of $t \geq 0$ real number values. Akin to Prime counting function Prime- $\pi(x)$ = number of primes $\leq x$, we can symbolically define nontrivial zeros counting function NTZ- $\pi(t)$ = number of NTZ $\leq t$ with t assigned to having real number values which are conveniently designated by 10^n whereby $n = 1, 2, 3, 4, 5, \dots$. The *cumulative Prevalence of nontrivial zeros* = NTZ- $\pi(t) / t = \text{NTZ-}\pi(t) / (10^n)$ when $t = 0$ to 10^n , whereby denominator t is [artificially] regarded as having integer number values. We conceptually define all consecutive NTZ gaps as i^{th} t -valued NTZ - $(i-1)^{\text{th}}$ t -valued NTZ. Thus there are CIS-IM-linear computed NTZ gaps. The numbers of NTZ between $10^0 - 10^1$ [interval = 9], $10^1 - 10^2$ [interval = 90], $10^2 - 10^3$ [interval = 900], $10^3 - 10^4$ [interval = 9000], $10^4 - 10^5$ [interval = 90000], $10^5 - 10^6$ [interval = 900000], $10^6 - 10^7$ [interval = 9000000], $10^7 - 10^8$ [interval = 90000000]... are 0, 29, 620, 9493, 127927, 1609077, 19388979, 226871900... with corresponding *rolling Prevalence of nontrivial zeros* = 0, 0.322, 0.689, 1.055, 1.421, 1.788, 2.154, 2.521... \implies *rolling Prevalence of nontrivial zeros* seems to overall fluctuatingly increase by around 0.366 in a "linear" manner. This limited observation alone suggests Cardinality of nontrivial zeros = $\|\text{CIS-IM-linear}\| = \aleph_0$ -linear.

In comparison, we further notice here the numbers of NTZ between $10^0 - 10^1$ [interval = 9], $10^0 - 10^2$ [interval = 99], $10^0 - 10^3$ [interval = 999], $10^0 - 10^4$ [interval = 9999], $10^0 - 10^5$ [interval = 99999], $10^0 - 10^6$ [interval = 999999], $10^0 - 10^7$ [interval = 9999999], $10^0 - 10^8$ [interval = 99999999]... are 0, 29, 649, 10142, 138069, 1747146, 21136125, 248008025... with corresponding *cumulative Prevalence of nontrivial zeros* = 0, 0.293, 0.650, 1.014, 1.381, 1.747, 2.114, 2.480...

On the overall objective to rigorously derive Algorithm-type proofs for Modified Polignac's and Twin prime conjectures [as based on Figure 9] and Equation-type proof for Riemann hypothesis [as based




Figure 9. Proportion (Prevalence) of Twin primes, Cousin primes [as partial calculations] and Sexy Primes [as partial calculations] with Proportion (Prevalence) of all Primes included. These Proportions (Prevalences) are essentially *self-similar fractal objects*. The $n = 1, 2, 3, 4, 5, 6, 7, 8, \dots$ in 10^n that is denoted with horizontal x-axis \implies the scale of this axis is non-linearly depicted using increasing powers of 10.




Figure 10. Simulated dynamic trajectories showing Origin intercept points when $\sigma = \frac{1}{2}$ and virtual Origin intercept points when $\sigma = \frac{2}{5}$ and $\sigma = \frac{4}{5}$. Horizontal axis: $Re\{\zeta(\sigma + it)\}$, and vertical axis: $Im\{\zeta(\sigma + it)\}$. Total presence of all Origin intercept points at the [static] Origin point. Total presence of all virtual Origin intercept points as additional negative virtual Gram[$y=0$] points on the x-axis (e.g. when using $\sigma = \frac{2}{5}$ value) at the [infinitely many varying] virtual Origin points; viz, these negative virtual Gram[$y=0$] points on the x-axis cannot exist at the solitary Origin point since the two trajectories form two colinear lines (or co-lines) [two parallel lines that never cross over].

on Figure 10], we apply infinitesimal numbers $\frac{1}{\infty}$ at two places using the following colloquially-stated propositions with their formal proofs given in Page 44 – 45 of [8].

PROPOSITION 6.1. *In the limit of never reaching a [nonexisting] zero hereby conceptually visualized as Prevalences of both even Prime gaps and the associated [positive and negative] Odd Primes never becoming zero whereby arbitrarily large number of different even Prime gaps that uniquely accompany all Odd Primes in totality will never stop recurring. Foundation Figure 9 is roughly and analogically based on cohomology as an algebraic tool in topology allowing Geometrical-Mathematical interpretation for positive Odd Primes. We note these Prevalences can only have $\frac{1}{\infty}$ values above zero in the large range of prime numbers [but must never have zero values].*

PROPOSITION 6.2. *In the limit of reaching an [existing] zero hereby conceptually visualized as the entire $-\infty < t < +\infty$ trajectory of Dirichlet eta function, proxy for Riemann zeta function, touching (symbolic) zero-dimensional $\sigma = \frac{1}{2}$ -Origin point only when parameter $\sigma = \frac{1}{2}$ whereby all nontrivial zeros [mathematically] located on (symbolic) one-dimensional $\sigma = \frac{1}{2}$ -critical line will [geometrically] declare themselves in totality as corresponding Origin intercept points. Foundation Figure 10 is roughly and analogically based on cohomology as an algebraic tool in topology allowing Geometrical-Mathematical interpretation for $0 < t < +\infty$ range. Our **Corollary** is: Any $\sigma \neq \frac{1}{2}$ co-lines that are $\frac{1}{\infty}$ above or below the zero-dimensional $\sigma = \frac{1}{2}$ -Origin point must never be classified as having nontrivial zeros. Then the **Proposition** must be: Only one unique $\sigma = \frac{1}{2}$ co-line that [repeatedly] touch the zero-dimensional $\sigma = \frac{1}{2}$ -Origin point must always be classified as having [infinitely-many] nontrivial zeros.*

$(0 < \sigma < 1) \equiv (0 < \sigma < \frac{1}{2}) + (\sigma = \frac{1}{2}) + (\frac{1}{2} < \sigma < 1)$. Usefully regarded as *variants* of infinite series are various power series and harmonic series [e.g. (with $s = \sigma \pm it$) Riemann zeta function $\zeta(s)$ via Dirichlet eta function $\eta(s)$ generating infinitely-many $0 < \sigma < 1$ -associated trajectories that are all of $-\infty < t < +\infty$ infinite length such as depicted by Figure 2 when $\sigma = \frac{1}{2}$], and various (sub)algorithms [e.g. Sieve of Eratosthenes generating Set of (\pm) prime numbers in its entirety and Subsets of (\pm) Odd Primes from even Prime gaps 2, 4, 6, 8, 10... that all have cardinality of ALN]. **Note that each $0 < \sigma < 1$ -associated trajectory represents a unique infinite series that is, crucially, *mutually exclusive* by being **mathematically, geometrically and topologically** different from other infinite series**. Analogous to term 'centroid' referring to fixed invariant (0-dimensional) point with PERFECT Point Symmetry representing *center of a geometric object in (n-dimensional) Euclidean space*; there must be: (i) [being valid for entire range +ve & -ve integers] the easily deduced integer number 0 in (1-dimensional) Figure 1 as **Centroid point** and (ii) [being valid for entire range $-\infty < t < +\infty$] Origin point in (2-dimensional) $\sigma = \frac{1}{2}$ Figure 4 when combined together with (2-dimensional) $0 < \sigma < \frac{1}{2}$ Figure 5 and (2-dimensional) $\frac{1}{2} < \sigma < 1$ Figure 6 [while fully satisfying (Remark 2.2) Principle of Equidistant for Multiplicative Inverse as previously discussed in Figure 7 with ONLY $\sigma = \frac{1}{2}$ containing the **most frequently & infinitely-often traversed or visited Centroid (Origin) point**]. Our unique Centroid (Origin) point for $\eta(s)$ is conceptually the Point Symmetry with ASSIGNED Central value as $\eta(\frac{1}{2} \pm it) = 0.0 + 0.0i = 0$ at intersection of horizontal real axis & vertical imaginary axis [and having two Line Symmetry of horizontal real axis as depicted by Figure 2 and vertical line $\sigma = \frac{1}{2}$ as depicted by Figure 3]. In comparison, COMPUTED Central value for $\zeta(s)$ via its functional equation having Line Symmetry of vertical line $s = \frac{1}{2}$ [that intersect horizontal real axis] is $\zeta(\frac{1}{2}) \approx -1.4603545 + 0.0i \approx -1.4603545$. As overall summary, we insightfully conclude mutually exclusive (sub)sets arising from prime numbers, composite numbers, Gram points and virtual Gram points MUST all conceptually comply in full with *Theory of Symmetry from Langlands program* and *Inclusion-Exclusion Principle* when "extended to the infinite (sub)sets".

Acknowledgements and Declarations

The author is grateful to Editors and Reviewers for feedbacks. Relevant generated data are included. This paper is dedicated to his daughter Jelena born 13 weeks early on May 14, 2012 with Very Low Birth

Weight of 1010 grams. AUS \$5,000 research grant was provided by Mrs. Connie Hayes and Mr. Colin Webb on January 20, 2020. He discloses receiving an additional AUS \$3,250 reimbursement from Q-Pharm for participating in the EyeGene Shingles trial commencing on March 10, 2020. From Doctor of Philosophy (PhD) viewpoint on Ageing, Dementia, Sleep, Learning, Memory and Number theory; he possesses average level of working, short-term and long-term memory, and *Concrete Mathematics* ability. While conducting research that requires advance *Abstract Mathematics*, he routinely practices behavioral augmentation on personal Stage 3 Deep Sleep which contributes to insightful thinking, creativity and memory, and Stage 4 REM Sleep which is essential to cognitive functions memory, learning and creativity.

References

- [1] Gross B.H. & Zagier D. (1980). On the critical values of Hecke L-series. *Bulletin de la Societe Mathematique de France*, 108(2), pp. 49 – 54.
- [2] Lichtman J. (2023). A proof of the Erdos primitive set conjecture. *Forum of Mathematics, Pi*, 11, E18, pp. 1 – 21. <http://dx.doi.org/10.1017/fmp.2023.16>
- [3] LMFDB Collaboration. (2024). *The L-functions and modular forms database*. <https://www.lmfdb.org> [Online accessed on 5 May 2024]
- [4] Padilla A. & Smith R.G.C. (2024). Smoothed asymptotics: from number theory to QFT. *arXiv:2401.10981v1*, pp. 1–28. <https://doi.org/10.48550/arXiv.2401.10981>
- [5] Rodriguez-Villegas F. & Zagier D. (1993). Square roots of central values of Hecke L-series. In *Advances in number theory (Kingston, ON, 1991)*, pp. 81 – 99. Oxford Sci. Publ., Oxford Univ. Press, New York.
- [6] Ting J.Y.C. (2020). Mathematical Modelling of COVID-19 and Solving Riemann Hypothesis, Polignac’s and Twin Prime Conjectures Using Novel Fic-Fac Ratio With Manifestations of Chaos-Fractal Phenomena. *J. Math. Res.*, 12(6) pp. 1 – 49. <https://doi.org/10.5539/jmr.v12n6p1>
- [7] Ting J.Y.C. (2023). Origin Point Must Represent Critical Line as Location for Nontrivial Zeros of Riemann Zeta Function, and Set Prime Gaps With Subsets Odd Primes Are Arbitrarily Large in Number. *J. Math. Res.*, 15(4) pp. 1 – 55. <https://doi.org/10.5539/jmr.v15n4p1>
- [8] Ting J.Y.C. (2024). On the Universal Presence of Mathematics for Incompletely Predictable Problems in Rigorous Proofs for Riemann Hypothesis, Modified Polignac’s and Twin Prime Conjectures. *J. Math. Res.*, 16(2) pp. 1 – 61. <https://doi.org/10.5539/jmr.v16n2p1>
- [9] Zhang Y. (2014). Bounded gaps between primes. *Ann. of Math.*, 179, pp. 1121 – 1174. <http://dx.doi.org/10.4007/annals.2014.179.3.7>

Appendix A. Predictability properties of Dirichlet L-series from Dirichlet L-functions that are relevant to Riemann hypothesis, and Birch and Swinnerton-Dyer conjecture

One can derive an [motivic] L-function from a **normalized defining polynomial** and obtain the coefficient sequence of this L-function associated to middle cohomology of the projective closure of hyperspace defined by the given polynomial equation. Recall that the 'general' Dirichlet series is an infinite series of the form $\sum_{n=1}^{\infty} a_n e^{-\lambda_n s}$ where a_n , $s [= \sigma \pm it]$ are complex numbers and $\{\lambda_n\}$ is a strictly increasing

sequence of nonnegative real numbers that tends to infinity. An 'ordinary' Dirichlet series $\sum_{n=1}^{\infty} \frac{a_n}{n^s}$ is obtained by substituting $\lambda_n = \ln n$ while a power series $\sum_{n=1}^{\infty} a_n(e^{-s})^n$ is obtained when $\lambda_n = n$. Parallel to 'ordinary' Dirichlet series, we further define a useful 'extra-ordinary' Dirichlet series [for non-polynomial (transcendental) equations such as $\cos s$ and e^s] below.

We characterize the Predictability property of Eq. (7) Generic L -function [referring to non-polynomial equations with having denominator = $f(n)$ thus forming any mathematical expressions that do not involve s], denoted by Generic- $L(s)$, which is our *****useful 'extra-ordinary' Dirichlet series [(non-)alternating power series]***** where $a_n = a_0, a_1, a_2, a_3, a_4, \dots$ are the Dirichlet coefficients that are, in theory, either Completely Predictable or Incompletely Predictable entities.

$$(7) \quad \text{Generic-}L(s) = \sum_{n=0}^{\infty} \frac{a_n}{f(n)} = \frac{a_0}{f(0)} + \frac{a_1}{f(1)} + \frac{a_2}{f(2)} + \frac{a_3}{f(3)} + \frac{a_4}{f(4)} + \dots$$

The Generic- $L(s)$ for $\cos s = \sum_{n=0}^{\infty} \frac{(-1)^n s^{2n}}{(2n)!} = \frac{s^0}{0!} - \frac{s^2}{2!} + \frac{s^4}{4!} - \frac{s^6}{6!} + \frac{s^8}{8!} - \dots$. When $s = 1 + 0i = 1$, we obtain $\cos 1 = \frac{1^0}{0!} - \frac{1^2}{2!} + \frac{1^4}{4!} - \frac{1^6}{6!} + \frac{1^8}{8!} - \dots$ as alternating power series where $a_n = (-1)^{2n} = a_0, a_1, a_2, a_3, a_4, \dots = 1, -1, 1, -1, 1, \dots$ are [computed] Dirichlet coefficients that are Completely Predictable entities. When $s = 0 + i = i$, we obtain $\cos i = \frac{1^0}{0!} + \frac{1^2}{2!} + \frac{1^4}{4!} + \frac{1^6}{6!} + \frac{1^8}{8!} + \dots$ as non-alternating power series where $a_n = (1)^{2n} = a_0, a_1, a_2, a_3, a_4, \dots = 1, 1, 1, 1, 1, \dots$ are [computed] Dirichlet coefficients that are Completely Predictable entities.

The Generic- $L(s)$ for $e^s = \sum_{n=0}^{\infty} \frac{s^n}{n!} = \frac{s^0}{0!} + \frac{s^1}{1!} + \frac{s^2}{2!} + \frac{s^3}{3!} + \frac{s^4}{4!} + \frac{s^5}{5!} + \dots$. When $s = 1 + 0i = 1$, we obtain $e^1 = \sum_{n=0}^{\infty} \frac{1^n}{n!} = \frac{1^0}{0!} + \frac{1^1}{1!} + \frac{1^2}{2!} + \frac{1^3}{3!} + \frac{1^4}{4!} + \frac{1^5}{5!} + \dots$ as non-alternating power series where $a_n = (1)^n = a_0, a_1, a_2, a_3, a_4, \dots = 1, 1, 1, 1, 1, \dots$ are [computed] Dirichlet coefficients that are Completely Predictable entities. When $s = 0 + i = i$, we obtain $e^i = \sum_{n=0}^{\infty} \frac{i^n}{n!} = \frac{i^0}{0!} + \frac{i^1}{1!} + \frac{i^2}{2!} + \frac{i^3}{3!} + \frac{i^4}{4!} + \frac{i^5}{5!} + \dots = \frac{1}{0!} + \frac{i}{1!} - \frac{1}{2!} - \frac{i}{3!} + \frac{1}{4!} + \frac{i}{5!} - \dots$ as alternating power series where $a_n = (i)^n = a_0, a_1, a_2, a_3, a_4, a_5, \dots = +1, +i, -1, -i, +1, +i, \dots$ [as perpetual repeating (periodic) patterns of $+1, +i, -1, -i$] are [computed] Dirichlet coefficients that are Completely Predictable entities.

We next characterize the Predictability property of Eq. (8) General L -function [referring to polynomial equations with having denominator = $f(n, s)$], denoted by General- $L(s)$, which is *****the 'ordinary' Dirichlet series [(non-)alternating harmonic series]***** where $a_n = a_1, a_2, a_3, a_4, a_5, \dots$ are the Dirichlet

coefficients that are, in theory, either Completely Predictable or Incompletely Predictable entities.

$$(8) \quad \text{General-}L(s) = \sum_{n=1}^{\infty} \frac{a_n}{n^s} = \frac{a_1}{1^s} + \frac{a_2}{2^s} + \frac{a_3}{3^s} + \frac{a_4}{4^s} + \frac{a_5}{5^s} + \dots$$

The [reproduced] Eq. (9) Riemann zeta function $\zeta(s)$ is the **most basic** General L -function, denoted here by $L_{\zeta}(s)$ [as non-alternating harmonic series], where $a_n = (1)^n = a_1, a_2, a_3, a_4, a_5, \dots = 1, 1, 1, 1, 1, \dots$ are [computed] Dirichlet coefficients that are Completely Predictable entities. In Eq. (9), the equivalent **Euler product formula** with product over all prime numbers implies the presence of Sieve of Eratosthenes.

$$(9) \quad \zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = \frac{1}{1^s} + \frac{1}{2^s} + \frac{1}{3^s} + \frac{1}{4^s} + \frac{1}{5^s} + \dots$$

$$= \prod_{p \text{ prime}} \frac{1}{1 - p^{-s}} = \frac{1}{1 - 2^{-s}} \cdot \frac{1}{1 - 3^{-s}} \cdot \frac{1}{1 - 5^{-s}} \cdot \frac{1}{1 - 7^{-s}} \cdot \frac{1}{1 - 11^{-s}} \dots \frac{1}{1 - p^{-s}} \dots$$

The [reproduced] Eq. (10) Dirichlet eta function $\eta(s)$ is the **most basic** General L -function, denoted here by $L_{\eta}(s)$ [as alternating harmonic series], where $a_n = (-1)^{n+1} = a_1, a_2, a_3, a_4, a_5, \dots = 1, -1, 1, -1, 1, -1, \dots$ are [computed] Dirichlet coefficients that are Completely Predictable entities.

$$(10) \quad \eta(s) = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^s} = \frac{1}{1^s} - \frac{1}{2^s} + \frac{1}{3^s} - \frac{1}{4^s} + \frac{1}{5^s} - \dots = \frac{1}{1^s} + \frac{-1}{2^s} + \frac{1}{3^s} + \frac{-1}{4^s} + \frac{1}{5^s} + \dots$$

$$= \prod_{p \text{ prime}} \frac{1}{1 \pm p^{-s}} = \frac{1}{1 + 2^{-s}} \cdot \frac{1}{1 - 3^{-s}} \cdot \frac{1}{1 + 5^{-s}} \cdot \frac{1}{1 - 7^{-s}} \cdot \frac{1}{1 + 11^{-s}} \dots \frac{1}{1 \pm p^{-s}} \dots$$

Reiterating: The complex number $s = \sigma \pm it$ with parameter σ , variable t and imaginary unit $i [= \sqrt{-1}]$. Its real part σ and imaginary part t are real numbers. Eq. (10), as Analytic continuation of Eq. (9), are related to each other as $\zeta(s) = \gamma \cdot \eta(s)$ or equivalently as $\eta(s) = \frac{1}{\gamma} \cdot \zeta(s)$ with proportionality factor $\gamma = \frac{1}{(1 - 2^{1-s})}$.

Denote \mathbb{P} = prime numbers, \mathbb{Z} = integer numbers, \mathbb{R} = real numbers and \mathbb{C} = complex numbers [represented by $z = a + bi$]. Based on the complex unit equation $i^2 + 1 = 0$ that define imaginary unit i , Gaussian integers are the set $Z[i] = \{a + bi \mid a, b \in \mathbb{Z}\}$ and Complex numbers are the set $\mathbb{C} = R[i] = \{a + bi \mid a, b \in \mathbb{R}\}$. The " \mathbb{R} -to- \mathbb{C} " polynomial equation K is a case of motivic L -function [which is a non-alternating harmonic series] referred to as Dedekind zeta function of the number field K defined by this specific one-variable equation: $x^2 + 1 = 0 \equiv x^2 = -1 \equiv x = \pm \sqrt{-1} \equiv x = \pm i$. Its L -function, $L_K(s)$, has [computed] Dirichlet coefficients $a_n = a_1, a_2, a_3, a_4, a_5, \dots = 1, 1, 0, 1, 2, 0, 0, 1, 1, 2, 0, 0, 2, \dots$ that are Incompletely Predictable entities [viz, $L_K(s) = \zeta_K(s) = \frac{1}{1^s} + \frac{1}{2^s} + \frac{0}{3^s} + \frac{1}{4^s} + \frac{2}{5^s} + \dots$]. The norm is defined as $N(a + bi) = a^2 + b^2$. Then the computed a_n values for Gaussian integers are precisely the Number of Points corresponding to the norm $N(a + bi)$ of values $n = 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, \dots$ [Note: $L_K(0) = -\frac{1}{4}$ is a special case of class number formula that relates many important invariants of an algebraic number field to a special value of its Dedekind zeta function.] However, when combined with $a_n = (1)^n = a_1, a_2, a_3, a_4, a_5, \dots = 1, 1, 1, 1, 1, \dots$ from $\zeta(s)$'s $L_{\zeta}(s)$, one can construct the associated ["diagonal"] Automorphic L -function [which is an alternating harmonic series] for $L_K(s)$ having eternally repeating (periodic) patterns given by [Completely Predictable entities] $1, 0, -1, 0$ to recursively derive all the a_n values in Motivic L -function

$L_K(s)$. This fundamental Automorphic L -function for $L_K(s)$, denoted by $L_A(s)$ [with its unique eternally repeating (periodic) a_n pattern of 1, 0, -1, 0], is one of the simplest Automorphic L -function in nature.

Here $L_A(s) = \zeta_A(s) = \frac{1}{1^s} + \frac{0}{2^s} - \frac{1}{3^s} + \frac{0}{4^s} + \frac{1}{5^s} + \dots$.

"Elliptic curve with LMFDB label 11.a.2" <https://www.lmfdb.org/EllipticCurve/Q/11/a/2> having analytic rank 0 can be equivalently written as $y^2 + y = x^3 - x^2 - 10x - 20$ (Minimal Weierstrass equation; viz, $y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6$ whereby $a_1, a_3 = 0$ or 1 and $a_2 = -1, 0$ or 1 are Weierstrass coefficients in \mathbb{Z}) OR $y^2z + yz^2 = x^3 - x^2z - 10xz^2 - 20z^3$ (Minimal Weierstrass equation, homogenize with extra variable z) OR $y^2 = x^3 - 13392x - 1080432$ (simplified equation; viz $y^2 = x^3 + Ax + B$). The [proven] modularity theorem asserts that every elliptic curve [viz, fundamental mathematical objects defined by Genus 1 cubic (or degree 3) polynomial diophantine equations in two variables] over the rational numbers is modular, meaning it is associated with an "infinite series" modular form. The unique correspondence in Langland program is given as {Counting problem 1 + p -number of solutions mod p [in finite series Elliptic curves] \leftrightarrow Coefficients of q^p [in infinite series Modular forms]} whereby nome $q = e^{\pi i \tau}$ & $p =$ prime numbers from Modular forms act as the (periodic) 'generating series or functions' having Group of symmetry = $SL_2(\mathbb{Z})$ [involving unit disk in complex plane]. Let E be an elliptic curve, and let N_p denote the number of points on $E \pmod{p}$. Set $a_p = p + 1 - N_p$. We can define the incomplete L -function of E [viz, Hasse-Weil L -function $L(E, s)$ of E]. We hereby provide the modern formulation of BSD conjecture that relates arithmetic data associated with E over a number field K to the behavior of this $L(E, s)$ of E at $s = 1$. More specifically, it is conjectured that algebraic $r_E = \text{ord}_{s=1} L(E, s)$; viz, the rank of abelian group $E(K)$ of points of E is the order of the zero of $L(E, s)$ at $s = 1$. By the modularity theorem, for any E , $L(E, s)$ has a holomorphic continuation to the entire complex plane. Then there exist the analytic r'_E as an integer such that the Taylor expansion of $L(E, s)$ at $s = 1$ is of a certain form that involves r'_E . BSD conjecture asserts that $r_E = r'_E$. Known results involving r_E or r'_E : If $r'_E = 0$ or 1 for an elliptic curve E , then BSD conjecture is true for E , whereby most E [viz, at least 83%] have rank 0 or 1. BSD conjecture is true for > 66% of all E [of rank 0, 1 or > 1]. Eventhough " $\approx 100\%$ of E " are conjectured to have rank 0 or 1 [infinite in numbers], the remaining " $\approx 0\%$ of E " having rank at least 2, while extremely rare, will also be infinite in number. **Analogous to distribution of prime numbers being deceleratingly infinitely-many or arbitrarily large in number (ALN), then rank $r_E \geq 2$ should heuristically be ALN associated with increasing rank size of 2, 3, 4, 5....** Large exact rank 20 (by Noam Elkies & Zev Klagsbrun in 2020) and 28 (by Noam Elkies in 2006) for elliptic curves E :

$$r_E = 20: y^2 + xy + y = x^3 - x^2 - 244537673336319601463803487168961769270757573821859853707x + 961710182053183034546222979258806817743270682028964434238957830989898438151121499931$$

$$r_E = 28: y^2 + xy + y = x^3 - x^2 - 20067762415575526585033208209338542750930230312178956502x + 34481611795030556467032985690390720374855944359319180361266008296291939448732243429$$

Example of a Family of L -function $L_n(s)$ from elliptic curves: For $n = 1, 2, 3, 4, 5, \dots$, $y^2 = x^3 - n^2x$. This family is related to the **congruent number problem**; viz, finding a congruent number which is a positive integer [or positive rational number] that is the area of a right triangle with three rational number sides.

Example "Elliptic curve with LMFDB label 14.a5" <https://www.lmfdb.org/EllipticCurve/Q/14/a/5> ["mixed" polynomial equation E as two-variable equation having analytic rank 0 is an alternating harmonic series that can be expressed in terms of a simpler periodic sequence; viz, an Automorphic object called a Dirichlet L -function or an L -function of a modular form]: $y^2 + xy + y = x^3 - x \equiv$ [factorized] $y(y + x + 1) = x(x + 1)(x - 1)$. Its degree 2 Euler product $L_E(s) = \prod_p F_p(p^{-s})^{-1}$ has [finitely-many] "bad primes" of 2 and 7 corresponding to $F_p T$ of $(1 + T)$ and $(1 - T)$, and [infinitely-many] "good primes" 3, 5, 11... corresponding to $F_p T$ of $(1 + 2T + pT^2)$, $(1 + pT^2)$, $(1 + pT^2)$ Its $L_E(s)$ [which is a (complex) alternating harmonic series] has "Counting solutions mod p " as $a_n = a_1, a_2, a_3, a_4, a_5, \dots = 1, -1, -2, 1, 0, 2, 1, -1, 1, 0, 0, -2, -4, -1, 0, 1, 6, -1, 2, \dots$. These [computed] Dirichlet coefficients are Incompletely Predictable entities **with perpetually alternating increasing and decreasing +ve and -ve integer values**. All a_n values can now be precisely obtained from the particular generating series below for this $L_E(s)$ [with FINITE unique numbers 1, 2, 7, 14 in the exponents of Euler product formula]: $f(z) = \eta(z)\eta(2z)\eta(7z)\eta(14z) = q \prod_{n=1}^{\infty} (1 - q^n)(1 - q^{2n})(1 - q^{7n})(1 - q^{14n})$, whereby $f(z)$ is some of the simplest modular forms known as **eta quotient**, and can be described in combinatorial terms. The q -expansion is $f(q) = q - q^2 - 2q^3 + q^4 + 2q^6 + q^7 - q^8 + q^9 + O(q^{10})$. Here, as alternating harmonic series, $L_E(s) = \zeta_E(s) = \frac{1}{1^s} - \frac{1}{2^s} - \frac{2}{3^s} + \frac{1}{4^s} + \frac{0}{5^s} + \dots$.

We now compare and contrast $L_E(s)$ for Elliptic curve as $\zeta_E(s)$ versus L -function for Riemann zeta function $\zeta(s)$. [1] Convergence: $\zeta_E(s)$ converges for $Re(s) > \frac{3}{2}$. $\zeta(s)$ converges for $Re(s) > 1$. [2] Trivial zeros: For $\zeta_E(s)$ occurs at $s =$ all negative integers including 0. For $\zeta(s)$, occurs at $s =$ all negative even integers but not including 0 / For $\eta_E(s)$ occurs at $s =$ all negative integers including 0. [3] Nontrivial zeros ("spectrum") obtained via Analytic continuation: For $\zeta_E(s)$, nontrivial zeros occurs ONLY at its Critical Line $\sigma = 1$. For $\zeta(s)$ there is NO nontrivial zeros / For $\eta(s)$, nontrivial zeros occurs ONLY at its Critical Line $\sigma = \frac{1}{2}$ **. Multiply $\zeta_E(s)$ by "Gamma factor" $2 \cdot (2\pi)^{-s} \Gamma(s) = \Gamma_{\mathbb{C}}(s)$ to obtain a **symmetric version of its functional equation applied to the Lambda-function given by $\Lambda_E(s) = 14^{-\frac{s}{2}} \cdot (2\pi)^{-s} \Gamma(s) \zeta_E(s)$, which satisfies $\Lambda_E(s) = \Lambda_E(1 - s)$ and having conductor of 14. This is associated with perfect Line Symmetry at vertical line $s = 1$. We can perform **Analytic normalization** for elliptic curves by shifting $\Gamma_{\mathbb{C}}(s)$ to $\Gamma_{\mathbb{C}}(s + \frac{1}{2})$ to instead obtain convergence for $Re(s) > 1$, Critical Line as $\sigma = \frac{1}{2}$ and perfect Line Symmetry as vertical line $s = \frac{1}{2}$ that is present in, for example, $\zeta_A(s)$, $\zeta_K(s)$, $\zeta(s)$, etc [as the uniformly used notation in *Generalized Riemann hypothesis*]. An important analytic ingredient for elliptic curves is a function of a complex variable, L , the Hasse-Weil zeta function [viz, $L(E, s)$ as mentioned above in relation to BSD conjecture] of E over \mathbb{Q} . This function is a **variant** of Riemann zeta function $\zeta(s)$ and Dirichlet L -function, defined as an **Euler product** with one factor for every prime number p .

Having Euler product and functional equation for $\zeta_A(s)$ [denoting Automorphic L -function $L_A(s)$ for $L_K(s)$]: $\zeta_A(s)$ converges for $Re(s) > 1$, Trivial zeros occurs at $s =$ all negative odd integers, and Non-trivial zeros ("spectrum") via analytic continuation occurs ONLY at its Critical Line $\sigma = \frac{1}{2}$. Having Euler product and functional equation for $\zeta_K(s)$ [denoting L -function for $L_K(s)$]: $\zeta_K(s)$ converges for $Re(s) > 1$,

Trivial zeros occurs at $s =$ all negative integers but not including 0, and Nontrivial zeros ("spectrum") via analytic continuation occurs ONLY at its Critical Line $\sigma = \frac{1}{2}$. Also having Euler product and functional equation for the **sum of divisors function** σ for a real or complex number z [expressed in sigma notation as $\sigma_z(n) = \sum_{d|n} d^z$, where $d | n$ is shorthand for " d divides n "], its L -function has Dirichlet L -series giving

rise to this (complex) non-alternating harmonic series $L_\sigma(s) = \zeta_\sigma(s) = \frac{1}{1^s} + \frac{3}{2^s} + \frac{4}{3^s} + \frac{7}{4^s} + \frac{6}{5^s} + \dots$ having Incompletely Predictable +ve a_n integer values 1, 3, 4, 7, 6, 12, 8, 15, 13, 18, 12, 28, 14, 24, 24, 31... that alternatingly increase and decrease in a perpetual manner [and is overall slowly increasing].

Complex multiplication for complex number $z = a + bi$ can be carried out using only three real multiplications ac, bd , and $(a + b)(c + d)$ as $\mathcal{R}[(a + ib)(c + id)] = ac - bd, \mathcal{I}[(a + ib)(c + id)] = (a + b)(c + d) - ac - bd$. The natural definition of $L(E, s)$ for elliptic curves only converges for values of s in complex plane with $Re(s) > \frac{3}{2}$ [or $Re(s) > \frac{1}{2}$ via performing **analytic normalization** by shifting $\Gamma_{\mathbb{C}}(s)$ to $\Gamma_{\mathbb{C}}(s + \frac{1}{2})$ to conform with Generalized Riemann hypothesis]. Helmut Hasse conjectured that $L(E, s)$ could be extended by analytic continuation to the whole complex plane. This conjecture was first proved by Deuring in 1941 for elliptic curves with complex multiplication. It was subsequently shown to be true for all elliptic curves over \mathbb{Q} , as a consequence of the modularity theorem in 2001. Reiterating: BSD conjecture relates the order of vanishing and the first non-zero Taylor series coefficient of the L -function associated to an elliptic curve E defined over \mathbb{Q} at the central point $s = 1$ to certain arithmetic data, the BSD invariants of E .

Elliptic curve $y^2 + xy = x^3 - x^2 - 2x - 1$ LMFDB label 49.a4 [<https://www.lmfdb.org/EllipticCurve/Q/49/a/4>]: Integral point / Torsion generator = $(2, -1)$, Conductor = 49 [with Modular form 49.2.a.a given by $q + q^2 - q^4 - 3q^8 - 3q^9 + 4q^{11} - q^{16} - 3q^{18} + O(q^{20})$ having Modular degree 1], Discriminant = -343 with all p -adic regulators being identically 1 since Analytic rank = 0 \implies finite $E(\mathbb{Q})$ solutions. At Central Point $s = 1$, Special value $L(E, 1)$ is the first non-zero value among $L(E, 1), L'(E, 1), L''(E, 1), \dots \approx 0.96665585280840577336653841951$ for elliptic curve 49.a4 computed using previous formula $\frac{1}{r!} L^{(r)}(E, 1)$; viz, $0.966655853 \approx L(E, 1) = \frac{\#\text{III}(E/\mathbb{Q}) \cdot \Omega_E \cdot \text{Reg}(E/\mathbb{Q}) \cdot \prod_p c_p}{\#E(\mathbb{Q})_{\text{tor}}^2} \approx \frac{1 \cdot 1.933312 \cdot 1.000000 \cdot 2}{2^2} \approx 0.966655853$. We now regard center of the critical strip as being $s = \frac{1}{2}$; viz, analytically normalized Central Point is $s = \frac{1}{2}$. Then computing Central value at this Central Point of elliptic curve 49.a4 that obeys Complex Multiplication is possible as per formula given previously in this paper (Remark 1.1).

Broadly, operations on L -functions using L -series include Unary operation [e.g. λ -operation], Binary operation [e.g. Addition, Multiplication, Selberg inner product], Pairing, Property, Relation, Family and Invariants [e.g. Sign, Self-dual (viz, Dirichlet coefficients a_n in L -function $L(s) = \sum_{n=1}^{\infty} \frac{a_n}{n^s}$ are real), Primitive, Degree of elliptic curves, Motivic (Arithmetic) weight w_{ar} , Algebraic weight w_{alg} (whereby **Hodge conjecture asserts that $w_{alg} = w_{ar}$ for any motivic L -function**); Conductor for various classes of elliptic curves with Analytic rank = 0 include integer values 11, 14, 15, 17, 19, 20, 21, 24, 26, 27, 30, 32, 33, 34,...; etc]. Two specific operations: Addition (Direct Sum) e.g. $L_\zeta(s) \oplus L_A(s) = L_K(s)$; and Multiplication




Figure 11. Graph of Z-function along $\Re(s) = \frac{1}{2}$ -critical line for $-\infty < t < \infty$ depicting UNIQUE nontrivial zeros spectrum for Elliptic curve with LMFDB label 389.a1 [with non-zero Analytic rank of 2] whereby we note Line Symmetry of vertical y-axis and the trajectory intersecting Origin point. Integral points are $(-2, 0)$, $(-2, -1)$, $(-1, 1)$, $(-1, -2)$, $(0, 0)$, $(0, -1)$, $(1, 0)$, $(1, -1)$, $(3, 5)$, $(3, -6)$, $(4, 8)$, $(4, -9)$, $(6, 15)$, $(6, -16)$, $(39, 246)$, $(39, -247)$, $(133, 1539)$, $(133, -1540)$, $(188, 2584)$, $(188, -2585)$.

(Tensor product or Rankin-Selberg convolution) e.g. $L_A(s) \otimes L_K(s) = L_K(s)$, $L_A(s) \otimes L_A(s) = L_\zeta(s)$, $L_\zeta(s) \otimes L_E(s) = L_E(s)$ where $L_\zeta(s)$ is identity element for \otimes .

Trivial zeros for $L_K(s) = -1, -2, -3, -4, -5, -6, -7 \dots$ (most frequent).

Trivial zeros for $L_A(s) = -1, -3, -5, -7, -9, -11, -13 \dots$ (intermediate frequency).

Trivial zeros for $L_\eta(s) = 0, -2, -4, -6, -8, -10, \dots$ (least frequent but include 0).

Trivial zeros for $L_\zeta(s) = -2, -4, -6, -8, -10, \dots$ (least frequent).

Nontrivial zeros for $L_K(s) = 6.02, 10.24, 12.99, 14.13, 16.34, 18.29, 21.02 \dots$ (most frequent).

Nontrivial zeros for $L_A(s) = 6.02, 10.24, 12.99, 16.34, 18.29 \dots$ (intermediate frequency).

Nontrivial zeros for $L_\eta(s) = 14.13, 21.02, 25.01, 30.42, 32.93, 37.58 \dots$ (least frequent).

Nontrivial zeros for $L_\zeta(s)$ DO NOT exist.

We adopt here $\zeta(\sigma + it)$ when $0 < t < \infty$ and ignore its complex conjugate $\zeta(\sigma - it)$ when $-\infty < t < 0$. As confirming the obvious relationship of trivial zeros and nontrivial zeros [located at $\sigma = \frac{1}{2}$ -Critical Line] derived from $L_K(s)$, $L_A(s)$ and $L_\zeta(s)$ [when analytically continued from Convergence for $\mathbf{R} > 1$ to the entire complex plane]; we observe these entities derived from $L_K(s)$ represent the combined entities derived from both $L_A(s)$ and $L_\zeta(s)$.

The gamma factors are $\Gamma_{\mathbb{R}}(s) = \pi^{-\frac{s}{2}} \Gamma(\frac{s}{2})$ and $\Gamma_{\mathbb{C}}(s) = 2(2\pi)^{-s} \Gamma(s)$. Required gamma factors to obtain functional equations [as completed zeta functions] will validly result in the $\Lambda(s) = \Lambda(1 - s)$ reflection relationship. Functional equations obtained for Riemann zeta function $\zeta(s)$ in $L_\zeta(s)$ requires gamma factor $\Gamma_{\mathbb{R}}(s)$; for $\zeta_A(s)$ in $L_A(s)$ requires gamma factor $\Gamma_{\mathbb{R}}(s + 1)$; for $\zeta_K(s)$ in $L_K(s)$ requires gamma factor $\Gamma_{\mathbb{C}}(s)$ [with Conductor = 4]; for $\zeta_A(s)$ in $L_A(s)$ requires gamma factor $\Gamma_{\mathbb{R}}(s + 1)$ [with Conductor = 4]; and for $\zeta_E(s)$ in $L_E(s)$ require gamma factor $\Gamma_{\mathbb{C}}(s)$ with different Conductor integer values for different elliptic curves. For Analytic normalization in elliptic curves, the required gamma factor is instead $\Gamma_{\mathbb{C}}(s + \frac{1}{2})$.

We again adopt here $\zeta(\sigma+it)$ when $0 < t < \infty$ and ignore its complex conjugate $\zeta(\sigma-it)$ when $-\infty < t < 0$. WITH applying analytic normalization, we now supply trivial zeros and nontrivial zeros [located at $\sigma = \frac{1}{2}$ -Critical Line (instead of $\sigma = 1$ -Critical Line)] derived from randomly selected Rank 2 Elliptic curve with LMFDB label 389.a1 { $y^2 + y = x^3 + x^2 - 2x$ } <https://www.lmfdb.org/EllipticCurve/Q/389/a/1> [when analytically continued from Convergence for $\mathbf{R}(s) > 1$ (instead of Convergence for $\mathbf{R}(s) > \frac{3}{2}$) to the entire complex plane; viz, $\mathbf{R}(s) > 0$]; we comparatively observe these [unrelated] **even MORE frequently occurring ** entities of trivial zeros and nontrivial zeros to be:

Trivial zeros for $L_E(s) = 0, -1, -2, -3, -4, -5, -6, \dots$

Nontrivial zeros for $L_E(s) = 2.87, 4.41, 5.79, 6.98, 7.47, 8.63, 9.63, 10.35, \dots$

Complying with simplest version of BSD conjecture: Analytic $L_E(\frac{1}{2}) = 0$ associated with this Rank 2 [viz, Rank $\neq 0$] elliptic curve \implies infinitely many $E(\mathbb{Q})$ solutions. Cf Analytic $L_E(\frac{1}{2}) \neq 0$ associated with Rank 0 elliptic curves \implies finitely many or zero $E(\mathbb{Q})$ solutions.

Euler product: $L(s) = \prod_p F_p(p^{-s})^{-1}$. Here, the reader can look up in LMFDB website the [single] "bad" prime at p 389 with $F_p(T) = 1 - T$ of degree 1, and all other [infinitely-many] "good" primes with their corresponding $F_p(T)$ e.g. at p 2, $F_p(T) = 1 + pT + pT^2$ of degree 2.

This Elliptic curve with LMFDB label 389.a1 has Dirichlet series, an (infinite) alternating harmonic series, of $L_E(s) = 1 - 2 \cdot 2^{-s} - 2 \cdot 3^{-s} + 2 \cdot 4^{-s} - 3 \cdot 5^{-s} + 4 \cdot 6^{-s} - 5 \cdot 7^{-s} + 9^{-s} + 6 \cdot 10^{-s} - 4 \cdot 11^{-s} - 4 \cdot 12^{-s} - 3 \cdot 13^{-s} + 10 \cdot 14^{-s} + 6 \cdot 15^{-s} - 4 \cdot 16^{-s} - 6 \cdot 17^{-s} - 2 \cdot 18^{-s} + 5 \cdot 19^{-s} - 6 \cdot 20^{-s} + 10 \cdot 21^{-s} + 8 \cdot 22^{-s} - 4 \cdot 23^{-s} + 4 \cdot 25^{-s} + 6 \cdot 26^{-s} + 4 \cdot 27^{-s} - 10 \cdot 28^{-s} - 6 \cdot 29^{-s} - 12 \cdot 30^{-s} + \dots$

Functional equation (analytically normalized) for this elliptic curve: $\Lambda(s) = 389^{\frac{s}{2}} \Gamma_{\mathbb{C}}(s + \frac{1}{2}) L(s) = \Lambda(1 - s)$.

(Hardy or Riemann-Siegel) **Z-function** for the Riemann zeta-function is a real-valued function defined in terms of the values of $\zeta(s)$ on the critical line via the formula $Z(t) := e^{i\theta(t)} \zeta\left(\frac{1}{2} + it\right)$, where $\theta(t)$ is the

Riemann-Siegel theta function $\theta(t) := \arg\left(\Gamma\left(\frac{2it+1}{4}\right)\right) - \frac{\log \pi}{2}t$. There is a bijection between zeros t_0

of $Z(t)$ and zeros $\frac{1}{2} + it_0$ of $\zeta(s)$. The Z-function of a general L -function is a smooth real-valued function

of a real variable t such that $|Z(t)| = |L(\frac{1}{2} + it)|$. Specifically, if we write the completed L -function as $\Lambda(s) = \gamma(s)L(s)$, where $\Lambda(s)$ satisfies the functional equation $\Lambda(s) = \varepsilon \overline{\Lambda}(1 - s)$, then $Z(t)$ is defined by $Z(t) = \frac{\varepsilon^{\frac{1}{2}} \gamma(\frac{1}{2} + it)}{|\gamma(\frac{1}{2} + it)|} L(\frac{1}{2} + it)$. The square root is chosen so that $Z(t) > 0$ for sufficiently small $t > 0$. The

multiset of zeros of $Z(t)$ matches that of $L(\frac{1}{2} + it)$ and $Z(t)$ changes sign at the zeros of $L(\frac{1}{2} + it)$ of odd multiplicity. The graph of nontrivial zeros spectrum for this NON-ZERO Analytic Rank 2 Elliptic curve with LMFDB label 389.a1 is depicted in Figure 11. **Summary by way of important observations at**

LMFDB website: We note the analogical graphs for elliptic curves with Analytic Rank 0 [viz, having zero independent basis point with infinite order which can be associated with either finitely many or zero $E(\mathbb{Q})$ solutions] DO NOT have trajectories that intersect the Origin point. This implies the simplest version of BSD conjecture is true. Elliptic curves are conventionally always of degree 2 [based on the degree




Figure 12. Graph of Z-function along $\Re(s) = \frac{1}{2}$ -critical line for $-\infty < t < \infty$ depicting UNIQUE nontrivial zeros spectrum for Dirichlet eta function, the *analytic continuation* of Riemann zeta function [with Analytic rank of 0] whereby we note Line Symmetry of vertical y-axis and the trajectory DO NOT intersect the Origin point. Integral basis is 1.

of their L-functions being the number $J + 2K$ of Gamma factors occurring in their functional equations]. For $x^2 + 1 = 0$, this "simplest" defining polynomial of degree 2 DO NOT have first trivial zero located at $s = 0$ and DO NOT have first nontrivial zero located at $t = 0$. Riemann zeta function $\zeta(s)$ is the most basic L-function associated with the simplest 1-variable polynomial equation $P(s) = x$ of degree 1. It has the analytically continued Dirichlet eta function $\eta(s)$, again of degree 1, having trivial zeros and nontrivial zeros (spectrum) on $\Re(s) = \frac{1}{2}$ -critical line whereby this spectrum [Figure 12; associated with Analytic rank 0] also DO NOT intersect the Origin point \implies its first nontrivial zero IS NOT located at $t = 0$. Finally, when based on Polar graph such as Figure 2 at $\frac{1}{2}$ -critical line for [Analytic rank 0] Riemann zeta function / Dirichlet eta function, we observe the first Gram[$y=0$] Point is located to the left of Origin point. Then, similar Polar graphs plotted for any zeta functions with non-zero Analytic rank will always show equivalent first Gram[$y=0$] Point to be located AFTER the first nontrivial zero, or Gram[$x=0, y=0$] Point, that is located on the Origin point.

(Received: July 22, 2024)

(Revised: 0, 2024)

Prof. Dr. John Yuk Ching Ting, Dental and Medical Surgery, 12 Splendid Drive, Bridgeman Downs, Queensland 4035, Australia. Affiliated with University of Tasmania, Churchill Avenue, Hobart, Tasmania 7005, Australia

E-mail : jycting@utas.edu.au

ORCID: 0000-0002-3083-5071