

# Anisotropy of Light Speed Due to Earth's Own Rotation

by

Florian Michael Schmitt

July 2024, Berlin

## Abstract

The Sagnac effect can be interpreted to demonstrate an anisotropy of light speed as a first-order effect on the scale of Earth's own rotation, but not for higher velocities such as Earth's orbital speed or its speed relative to the CMB. Following logic reason, experiments investigating the second-order effect should also exhibit this deviation due to Earth's own rotation, but not for higher velocities. Specifically, interferometer experiments with optical resonators would be capable of verifying this matter. This paper shows that none of the existing experiments provide a statement on this issue. The detection of such a second-order effect from Earth's own rotation would provide clues to new physics.

## 1. Introduction

An analysis of relevant interferometer experiments, especially those using optical resonators, yields two important results: First, the test theories are used in a way that disregard Earth's own rotation from the outset, so only the influence of orbital speed and higher velocities on anisotropy can be ruled out with increasing precision. Second, various effects are excluded or averaged out from the measurement results. These include supposed imbalances of the rotating measurement apparatus or unspecified "spurious signals". A reanalysis of the raw data of the experiments is suggested, provided they exist in unprocessed form. Moreover, it would be sensible to conduct an experiment with optical resonators that focuses solely on the anisotropy on the scale of Earth's own rotation. The following examines existing experiments regarding these aspects.

## 2. Classical Interferometer Experiments, i.e., Apparatus of the Michelson-Morley Type

At an Earth's rotational speed of about 464 m/s, a shift of the interference fringes of the experiment in the order of  $v^2/c^2$  would be expected. Assuming a light speed of  $2.998 \times 10^8$  km/s, this corresponds to a squared anisotropy of about  $2.4 \times 10^{-12}$  m/s. Considering the light wavelength and the arm length of 2 m in the Michelson-Morley apparatus, this corresponds to a shift of the interference fringes by an order of  $10^{-5}$  fringes. Even a hundredfold increase in arm length would not yield a measurable result. All classical interferometer experiments are therefore unsuitable for demonstrating the second-order effect due to Earth's own rotation.

## 3. Interferometer Experiments Using Optical Resonators

Only experimental setups that are rotatably mounted, analogous to the classical apparatuses, are considered, as the shift of an interference or the measurement of the beats of the resonators' resonance deviations requires that the direction of the resonators can be changed relative to the

velocity vector of Earth's own rotation. Experiments that do not meet this condition are therefore inapt and not considered. The only three dealing with a rotatable apparatus are:

### 1.1 Brilliet and Hall, 1978

The paper by Brilliet and Hall points to a spurious signal, quote: "*—and a spurious nearly sinusoidal frequency shift at the table rotation rate  $f$ .*" Since this is a signal of the simple rotational frequency, but we are looking for a signal of the double frequency due to squaring, this signal should not be relevant. However, the paper further states: "*we find that taking data in blocks of  $N$  table rotations ( $N \approx 8-50$ ) is helpful in minimizing the cross coupling of these noise sources into the interesting Fourier bin at 2 cycles per table revolution.*" Thus, a first averaging of the data over 8 to 50 rotations is already taking place, which should have eliminated a large part of our sought signal. It is further described: "*To discriminate between this spurious signal and any genuine ether effect, we made measurements for 12 or 24 sidereal hours*" and "*Averaging after this rotation leads, as shown in Fig. 2, to a typical 1-day result below 1 +/- 1 Hz.*" At this point, it becomes clear that a rotation-dependent signal is not being sought at all but completely averaged out, and the search is exclusively for a diurnal signal. The signal that would result from a rotation of the apparatus is therefore not further recorded. The published result data of the experiment are thus completely inconclusive for our sought signal. If raw data still exist for this experiment, it would be interesting to know the uninterpreted magnitude of the signals in connection with the double rotation frequency of the apparatus.

### 1.2 Eisele et al, 2008

Within the paper by Eisele et al, divergent data are averaged beforehand, the cause of which is attributed to an unavoidable tilt of the apparatus: "*Sensitivities of the beat to tilts were determined at the beginning of each run.*" and "*The means of the amplitudes of the tilt modulations at  $2\omega$  were less than  $0.2 \mu\text{rad}$ , leading to an effect on the means of the coefficients of  $B$ ,  $C$  of less than  $5 \times 10^{-17}$ .*" Despite averaging, there is a phase shift of  $0.2 \mu\text{rad}$  here, which is already a multiple ( $10^{-8}$ ) of our sought signal (order of  $10^{-12}$ ). The evaluation of the measurement results is then done using the Robertson-Mansouri-Sexl (RMS) test theory. The paper states: "*The RMS test theory (with the effects of Earth's rotational velocity neglected) leads to expressions for  $B(t)$ ,  $C(t)$  with similar structure.*" In further analysis, the data are then interpreted only with regard to  $B(t)$  and  $C(t)$ , always neglecting Earth's own rotation. The published result data of the experiment are thus inconclusive for our sought signal for both reasons mentioned above. Here too, the raw data would be highly interesting and should be interpreted without applying the test theory, or at least taking into account Earth's own rotation.

### 1.3 Herrmann et al, 2018

In this experiment as well, the sought signals are suppressed from the outset. An initial analysis of the paper by Herrmann et al deals with the error signals, particularly those at double the rotational frequency of the apparatus: "*For example, gravitational or centrifugal forces that act on the resonators may get modulated with the turntable rotation and therefore modulate the length of the resonators. However, most of these effects lead to a modulation at a rate of  $\omega_{tt} = 2\pi/T_{tt}$  so that they are in principle distinguishable from the anisotropy signal searched for at  $2\omega_{tt}$ . Moreover, if the data spans more than one day, systematic effects with a fixed phase in the laboratory average out in the analysis for an anisotropy of  $c$  that is fixed relative to a sidereal frame.*" Apparently, the slight errors at double the rotational frequency can only be attributed to a variety of causes, so the strategy is chosen to investigate a daily modulation: "*This daily modulation is essential to discriminate an anisotropy signal from constant or slowly varying systematic effects caused by active rotation as described in Section I. Only systematic effects subjected themselves to a 23.93 h modulation would mimic such a sidereal anisotropy signal.*" By no means can our sought signal be found within a diurnal averaging. And finally,

the paper even specifically states: *“These samples extend over 450 s each such that we may neglect a possible modulation due to Earth's rotation within each sample.”* Furthermore, the RMS test theory again is applied in a way that disregards Earth's own rotation from the outset.

#### **4. Conclusion**

All experiments that could provide information about a second-order anisotropy of light speed concerning Earth's own rotation have been interpreted in a way that a signal at double the rotational frequency of the apparatus and in the sought magnitude is either (partly together) suppressed in the error analysis, averaged out through long-term investigation, or eliminated due to the way the test theory is applied. If the raw data of even one of the experiments were evaluated such that such a signal emerged, it would indicate new physics.

Ch. Eisele, A. Y. (2008). Laboratory Test of the Isotropy of Light Propagation at the  $10^{-17}$  Level. *Physical Review Letters* PRL 103, 090401 (2009).

Hall, A. B. (1978). Improved Laser Test of the Isotropy of Space. Colorado, USA: *Physical Review Letters*, Volume 42, Number 9.

Michelson, A. A., Morley, E.W. (1887). On the Relative Motion of the Earth and the Luminiferous Ether (Vols. 1887, 34 (203): 333–345 1881). *American Journal of Science*.

S. Herrmann, A. S. (2018). Rotating optical cavity experiment testing Lorentz invariance at the  $10^{-17}$  level. Berlin: arXiv:1002.1284v1 [physics.class-ph] 5 Feb 2010.

Sagnac, Georges. (1913). L'éther lumineux démontré par l'effet du vent relatif d'éther dans un interféromètre en rotation uniforme. In: *Comptes Rendus*. 157, S. 708–710.