

Refutation of Wiltshire's "Timescape cosmology"

Warren D. Smith (warren.wds@gmail.com) January 2025.

Abstract. I use previously-published observed astronomical facts to refute Wiltshire's recently-hyped "Timescape cosmology" in 1 page.

Introduction

Recently, there has been worldwide hype trumpeting the claim, mainly by New Zealand cosmologist David L. Wiltshire, that the universe's "accelerating expansion," detected experimentally by supernova cosmology projects in (what was at the time) a big surprise that won the 2011 physics [Nobel](#) prize, actually is a mere *illusion*: cosmic expansion actually is *slowing*, just as one would have naively expected. Wiltshire proposes as an improvement over the long-standing [FLRW](#) (Friedmann-Lemaitre-Robertson-Walker) Λ CDM model, his "timescape" model.

FLRW is a famous exact solution of the Einstein general relativity equations (and Λ /matter/radiation equations-of-state) under the Friedmann approximate assumption that the universe at any time is filled with a spatially-constant-density mixture of matter (gas, dust, and "cold dark matter"), radiation, and Einstein cosmical constant Λ . Because the FLRW equations have a known exact solution in terms of elliptic functions, it is mathematically rigorous. Fits of Λ CDM parameters to a vast number of observations (yielding a nonzero repulsive-signed cosmical constant Λ) have yielded excellent agreement. Two textbooks which cover this (which may not be the best anymore, but possess the advantage that I read them) are Misner, Thorne, Wheeler 1973 and Ciufolini & Wheeler 1995. For confrontations with observation more recent than those books, start with the 2011 Nobel prize lectures by Saul [Perlmutter](#), Brian P. [Schmidt](#), and Adam G. [Riess](#) (all also viewable as online videos on the Nobel prize website: [P](#), [S](#), [R](#)) as well as later publications by the Planck satellite team.

Timescape instead proposes an *inhomogeneous* universe, with that inhomogeneity *increasing* with time. Gas condensed into galaxies, which segregated into galaxy-clusters and superclusters, distributed into sheet-like and filament-like structures rich in galaxies, with "voids" in between containing far fewer galaxies. Timescape posits that the "accelerating expansion" of the universe, is *not* caused by nonzero Λ , but rather is an illusion caused by the fact that we humans live in the *Milky Way* galaxy (part of "local group" galaxy-cluster which in turn is a member of the "Virgo supercluster"), rather than the *middle of a void*. Relativistic time-dilation effects (**RTDE**) cause time to pass at different rates in those two places, and the increase of that rate-discrepancy (as the universe evolved to become more inhomogeneous), as interpreted by humans who (since their thinking was FLRW-based) did not take it into account, is what is responsible for this illusion. Wiltshire pointed out that the era of development of void/sheet/etc structure *coincided* with the era when FLRW claimed that Λ took over as the main driver term in the Friedmann equation, thus "accelerating expansion."

The recent **worldwide hype** was triggered by the [2024 paper](#) by Seifert, Lane, Galoppo, Ridden-

Harper, Wiltshire (SLGRHW24) claiming the confrontation between (a) observations (the "Pantheon+ type Ia supernovae spectroscopic dataset") versus (b) their computer simulations of timescape, shows "very strong evidence" that Timescape is *superior* to Λ CDM.

How large RTDEs is Wiltshire claiming? Here are three time-dilation numbers to consider:

- a. **1.000008** based on this simple numerical "sanity check": the escape velocity from our galaxy is believed about 500 km/sec and from clusters and superclusters about 1000 km/sec $< 0.004c$. This should cause the time dilation factor between void interiors and galaxy interiors to be < 1.000008 .
- b. **1.001** based on the 10^{-3} in this quote from SLGRHW24: "In standard cosmology, differences from average FLRW expansion are assumed to be mostly attributed to local Lorentz boosts – i.e., peculiar velocities – of source and observer, with gravitational potentials contributing fractional variations of $\sim 10^{-5}$ of average expansion at galaxy and galaxy cluster scales. In timescape, the same fractional variation can be up to $\sim 10^{-3}$."
- c. **1.35** (which struck me, a priori, as worryingly probably insanely too-large to possibly be true) based on the 19 Dec. 2024 press release from the Royal Astronomical Society "[Dark energy 'doesn't exist' so can't be pushing 'lumpy' Universe apart – study](#)" featured maximally-prominently (as of January 2025) on D.L.Wiltshire's web [page](#) at the University of Canterbury NZ, and on which he is listed as the sole "scientific contact." Quote: "a clock in the Milky Way would be about 35 per cent slower than the same one at an average position in large cosmic voids."

When by email I asked Wiltshire to explain the (very large!) discrepancies among these three RTDE numbers, he responded as follows (quoted from his email):

- a. The notion of escape velocity that you are using is entirely based on gravitationally bound structures existing as isolated compact bodies in empty space, or embedded in the standard cosmology, with the dipole in the CMB anisotropy spectrum being purely kinematic. Likewise the notions of time dilation that you apply are based on some sort of static approximation. E.g, in talking about the local group of galaxies we know that if we factor out the motion of the earth around the galaxy, and the motion of the Milky Way with respect to the barycentre of the local group, then in order for the dipole in the CMB to be kinematic the local group [must be] moving at 630 km/sec in the direction of Hydra. That is our "peculiar velocity" in the standard FLRW cosmology, and it is where the order of magnitude numbers of "escape velocity" that you throw around come from. In the timescape model, however, there is "peculiar expansion" of the local structures in our vicinity – the anisotropy of our own filament within the local void that is the key to understanding the cosmological dipole anisotropies. In timescape there is a quasilocal uniform Hubble expansion right down to scales of 5 Mpc which means that any peculiar velocity is actually only of order 10 km/s on a background within which expansion is uniform after recalibration of rulers and clocks across the voids. The voids have positive kinetic energy of expansion, or equivalently negative spatial curvature. It is the quasilocal uniform Hubble expansion condition in timescape, which allows one to extract a simple expansion law in what is considered to be the "nonlinear regime" in standard cosmology.

– and –

Everything you said about escape velocities etc is based on a completely empty universe in which we can calibrate clocks relative to that of an observer at spatial infinity where space is

Minowski, and has a timelike Killing vector. That is essential in defining the notion of gravitational mass in a conventional setting. Every statement you make is based on this notion then superimposed on a FLRW background. However, the universe is not empty. Furthermore, it is changing with time on cosmological scales – there is no global time symmetry. The time dilation I discuss is not related to bound systems. It relates to gradients in local expansion of expanding regions, which may be viewed as gradients in spatial curvature or equivalently, in an average sense, to the quasilocal kinetic energy of expansion. The question of how to keep two clocks calibrated over billions of years when there is a very small relative volume deceleration is a fundamentally different problem to that of time dilations for static situations within bound structures.


- b. I do not know how you come up with your numbers... I do *not* claim that time dilations are of order 1.001 in my papers. This is incorrect. From my first paper on this in 2007, [Cosmic clocks, cosmic variance and cosmic averages](#), I have always claimed time dilations of an order similar to 35%.

I thank Wiltshire for those clarifications about the RTDE numbers. The insanity of the number "1.35" now makes it a simple matter to refute his entire proposal.

The Refutation

How can two people ("Amy" and "Bob") measure their relative time-dilation factor? Amy shines her laser toward Bob with pre-agreed frequency (say) 10^{15} Hertz. Bob measures the frequency (say) 0.8×10^{15} Hertz. "Aha!" says Bob. "I deduce dilation factor 1.25." Alternatively (or "also") Bob could shine his laser toward Amy.

A. The largest known void in the observed universe is the [Boötes supervoid](#) (so-called because it lies near the constellation Boötes in the sky), an approximately spherical region of space containing only 60 galaxies instead of the 2000 expected from a volume this large. Its width is about 400 million light-years. The picture mapping the situation is taken from wikipedia which in turn got it from Richard Powell's [Atlas of the Universe](#). Robert Kirshner discovered it in 1981 in a survey of galactic redshifts. Its center is located 700 million light-years from Earth.



B. On the boundary of the Boötes supervoid (on our side of it) are the [Hercules Superclusters](#) approximately 330 million light years in diameter. Hercules is regarded as an "unusually large" supercluster.

C. In front of them, in turn, (left bottom part of the picture) is the [Northern Local Supervoid](#), containing a few small galaxies (primarily spirals) and galaxy clusters, but mostly empty. This is the supervoid nearest to Earth, with center 199 Mly away and width 339 Mly.

Now enquire: what are the observed redshifts of spectral lines produced by galaxies in those three regions?

- A. Cruzes et al 2002 "investigate the physical properties of a sample of 26 galaxies in the Boötes void and classify these galaxies based on the emission lines in their spectra." For the 26 galaxies they tabulate in their table 1 the observed redshift z values range from 0.0432 to 0.0574.
- B. Kopylova & Kopylov 2013 in their table 1 find redshifts (expressed as equivalent recessional velocities) ranging between 8066 and 13563 km/sec, or re-expressed (by dividing by the speed of light $c=299792.458$ km/sec) as z -values, between 0.0269 and 0.0452.
- C. Redshifts ≤ 15000 km/sec (§2 of Lindner et al 1997's arXiv version), i.e. z -values between 0 and 0.050.

From these we may [deduce](#), very conservatively, that relative time-dilation factors between void centers versus galaxies in superclusters on the void-wall are ≤ 1.0574 , and presumably actually ≤ 1.014 and even that ought to be a large overestimate. In any case, it is clear that the answer is nowhere near as large as 1.35, even for the largest void in the known universe bounded by an unusually large supercluster.

(Also, if "TimeScape" had been correct, then it seems rather peculiar that blueshifts are seen as rarely as they are.)

Discussion

In his same email to me, Wiltshire claimed that he was a proponent of "rigour" and that papers failing to possess it should be rejected, often even without refereeing. As a mathematician who has studied the issue of rigor in physics (or its lack) I am more qualified than most to talk about that topic.

For most nonlinear PDEs (partial differential equations), in particular the Navier-Stokes equations of hydrodynamics and Einstein's general relativity (GR) field equations, present-day mathematics unfortunately is largely incapable of making rigorous statements. We **can**:

1. Sometimes find (and once found, it is easy to verify), exact solutions. Unfortunately: for the vast majority of real physical problems, especially "messy" ones, usually nobody can find an exact solution.
2. Prove certain formal identities, for example some tensor identities in GR.
3. Sometimes prove some statements about the nature of solutions when they exist, for example the Schoen-Yau, later reproven by Witten (& friends and successors) "[positive mass theorems](#)" in GR.

4. In the cases of Navier-Stokes hydrodynamics and GR, solutions have been proven to exist for an initial-data-dependent positive amount (albeit perhaps very small) of time into the future, in a large class of situations ("local existence"). The latter is due to Yvonne Choquet-Bruhat and discussed by Misner, Thorne, Wheeler 1973; for the former see, e.g, Kato 1984.
5. Sometimes, using *very* difficult and long proofs, which historically very few people have been capable of producing (and I have not been one of those few) one can prove eternal existence of solutions *provided* the initial data is "smooth enough" and "small enough" in certain precisely stated norm-senses. That was first accomplished for Navier-Stokes in work of O.A.Ladyzhenskaya, and for GR vacuums near-enough to Minkowski flat space by Christodoulou & Klainerman. My guess would be that those proofs are well beyond Wiltshire's ability level. Regardless of whether that ability-guess is valid, all such theorems so far have been almost useless for practical purposes because essentially no hydrodynamical/aeronautical engineer, and no astrophysicist, ever cares about simulating data that small.

But we presently usually **cannot**

6. Prove solutions *exist* and are *unique*, and that keep existing for arbitrarily long (user-specified) future timespans.
7. And if you cannot do that, then you basically can't prove *anything*, such as: "coarse graining" alleged approximations, models of fluid "turbulence," validity to within any error bound of some algorithm for allegedly finding approximate solutions numerically, etc. Indeed, nobody, ever, has shown that the Navier-Stokes or Einstein-GR PDEs are algorithmically simulable at all. No algorithm has ever been invented that anybody can prove successfully does either.

Let me try to make that even clearer. If somebody (for example Thomas Buchert, whose work Wiltshire's "timescape" largely rested upon) says they have coarse-grained approximate equations that understand cosmology in the presence of growing inhomogeneity then *those claims cannot presently be supported by rigorous mathematics*. If somebody claims they have a computer program that simulates GR and then tells you that, by running their computer, they deduced, with, (say) error bound of 5%, some claim about some physical situation, then *those claims cannot presently be supported by rigorous mathematics*.

And indeed, in the papers I have examined by Buchert and Wiltshire on the present topic, they proved a grand total of *zero* theorems.

Because of those lacks of rigor and rigorizability, the only options we have to attempt to assess errors from such computations, are either (a) comparison versus physical experiments, or (b) guesstimates based on comparing repeated runs of the computer program with different inputs and parameters. Until and unless we can attain rigor – which we presently cannot! – those attempts will always be unreliable.

The attempts of this kind most likely to **work well**, are, for example, GR black-hole collision simulations, involving pure vacuum (*no* matter) and *no* unknown or merely approximately known parameters, *simple* initial (and final) data – the latter sometimes with support from rigorous proofs of stability of black hole spacetimes – and (thanks to [LIGO](#)) numerous experimental *observations*.

On the other hand, the attempts of this kind that I would **trust the least** (and perhaps even a priori

presume to be total garbage!) are: simulations of the entire universe involving growing messy inhomogeneities in a crucial way!

Is there, then, any hope to attain rigor about this kind of question? Let me make two speculative suggestions (or perhaps a better descriptor would be "baseless hopes"):

1. Perhaps some *exact solution* of a cosmological model featuring growing inhomogeneities (e.g. featuring some sort of periodic lattice of them) could be found.
2. Witten's "positive mass theorem" states in precise ways claims that it is *impossible* for ordinary matter, and GR, to simulate "negative mass" matter, such as the repulsive-signed Einstein cosmical constant Λ . That suggests that Wiltshire's goal of getting rid of Λ because its alleged effects are merely illusions "simulated" by ordinary matter, is not possible. Here I do *not* want to claim belief in that "suggestion." I merely want to suggest that perhaps some version of Witten et al's ideas, can be used to produce some sort of rigorous upper and/or lower bounds, on the capabilities of such simulo-effects, and/or on the validities of "coarse graining" approximation notions.

I am not optimistic about either plan, but these seem the two avenues most worth investigating if you want "rigour."

Obviously, the FLRW cosmology is not an exactly correct depiction of reality, because the universe really is inhomogeneous, and that really must exert some cosmological effect. Wiltshire et al are right about that. The problem is that these effects have been, up until now, only a small perturbation to FLRW (1% level or less). Furthermore, if FLRW is correct and if Einstein Λ truly is the reason for the accelerating expansion, then I expect the effect of such inhomogeneities also to remain small *forever*. That is because in the future, Λ will convert the universe into a closer and closer approximation to exponentially-expanding "de Sitter spacetime" containing a vacuum only slightly polluted by ordinary and dark matter and radiation, where the "pollution level" will become exponentially small. Eventually any inhabitant of the Milky Way will only be able to see the few galaxies gravitationally bound to it, with all others unseeable since over the de Sitter "horizon."

Here is a (quite incomplete) chronologically-ordered list of papers with disagreeing conclusions:

- Siegel & Fry 2005: "We evaluate the effect of inhomogeneity energy on the expansion rate of the universe. Our method is to expand to Newtonian order in potential and velocity but to take into account fully nonlinear density inhomogeneities... [We conclude] the contributions due to inhomogeneities never mimic the effects of dark energy or induce an accelerated expansion."
- Buchert & Räsänen "Backreaction in late-time cosmology" 2012: Dismisses Siegel & Fry (and many others of their ilk) with the single sentence "Estimates in first order perturbation theory cannot resolve the issue, because the average of the first order perturbation vanishes, and the contribution of the square of the first order perturbations is *not gauge invariant* without the contribution of the intrinsic second order perturbations." Buchert and Wiltshire then proceeded never to cite Siegel & Fry ever again, apparently considering the matter settled by this single sentence.

But: *why* should we have or want gauge invariance? Why does it matter? B&R simply do not say. The Nobel prize winning "BCS model" for superconductivity was not gauge-invariant either, and that did not seem to hurt its experimental validity in the slightest. (And the FLRW uniform-cosmology model has had very good quantitative success, far exceeding that of

BCS.)

- Adamek et al 2015: Effectively dismisses Buchert & Räsänen's title and that single sentence and the Wiltshire 35% claim with: "The large-scale homogeneity and isotropy of the Universe is generally thought to imply a well-defined background cosmological model. It may not. Smoothing over structure adds in an extra contribution, transferring power from small scales up to large. Second-order perturbation theory implies that the effect is small... We investigate this using two different N-body numerical simulations: a 3D Newtonian and a 1D simulation which includes all relevant relativistic effects. We show that while perturbation theory predicts an increasing *backreaction* as more initial small-scale power is added, in fact the virialization of structure saturates the backreaction effect at the same level independently of the equality scale. This implies that backreaction is a *small* effect independently of initial conditions. Nevertheless, it may still contribute at the percent level to certain cosmological observables."
- Saulder et al 2018 computer simulates TimeScape reaching the opposite conclusion to the computer simulations of Sefiert et al 2024 (which never cited it)! "Using large-scale galaxy surveys such as SDSS and 2MRS, we test the variation of expansion expected in the Λ CDM model versus a more generic differential expansion using our own calibrations of bounds suggested by timescape cosmology. Method. Our test measures the systematic variations of the Hubble flow towards distant galaxies groups as a function of the matter distribution in the lines of sight to those galaxy groups. We compare the observed systematic variation of the Hubble flow to mock catalogues from the Millennium Simulation in the case of the Λ CDM model, and a deformed version of the same simulation that exhibits more pronounced differential expansion. Results:... statistical tests... consistently yield results preferring Λ CDM over our approximated model of timescape."
- Wiltshire published a 2019 "comment" responding to Saulder et al, which he summarized to me as "Since they did not have better simulations Saulder et al deformed Newtonian N-body simulations. However, this misses a key aspect of timescape; there is no backreaction." However that misses a key aspect of backreaction: it is too tiny to matter (say Adamek et al).

It is rather amusing to watch this war of words between these two camps, who however mostly just ignored each other for 20 years while happily publishing their own stuff. The result is as clear as mud.

You may be wondering how it can be that many different publications by many different authors over 2 decades about TimeScape and related cosmologies have managed, via competing computer simulations and competing human analyses, to reach such vastly disagreeing conclusions! I believe a big reason is: "because this entire area is entirely nonrigorous"! That is why I wrote this paper: I considered my reasoning here from observational data so simple and convincing that it settles the argument.

References

Julian Adamek, Chris Clarkson, Ruth Durrer, Martin Kunz: [Does small scale structure significantly affect cosmological dynamics?](#) Phys.Rev.Lett. 114,5 (2015) #051302.

Thomas Buchert: [Dark Energy from structure: a status report](#), Gen. Relat. Gravit. 40 (2008) 467-527.

- Thomas Buchert: [Toward physical cosmology: focus on inhomogeneous geometry and its non-perturbative effects](#), Classical & Quantum Gravity 28 (2011) #164007.
- Thomas Buchert & Syksy Räsänen: [Backreaction in late-time cosmology](#), Ann.Rev.Nucl.Part.Sci. 62 (2012) 57-79.
- Ignazio Ciufolini & John A. Wheeler: Gravitation and inertia, Princeton University Press 1995. QC173.59.G44C58.
- Demetrios Christodoulou & Sergiu Klainerman: The global nonlinear stability of Minkowski space, Princeton University Press (Princeton Mathematical Series #41) 1993. QC173.59.S65C57.
- Shawn Cruzen, Tara Wehr, Donna Weistrop, Ronald J. Angione, Charles Hoopes: [Spectroscopy of galaxies in the Boötes void](#), The Astronomical Journal 123 (Jan.2002) 142-158.
- Ethan R. Siegel & J.N.Fry: The Effects of Inhomogeneities on Cosmic Expansion, The Astrophysical Journal 628,1 (July 2005).
- Toshio Kato: Strong L^p -solutions of the Navier-Stokes equations in \mathbb{R}^m , with application to weak solutions, Mathematische Zeitschrift 187 (1984) 471-480.
- Robert P. Kirshner, Augustus Oemler Jr, Paul L. Schechter, Stephen A. Sackett: [A million cubic megaparsec void in Boötes?](#), Astrophysical Journal 248 (Sep.1981) L57-60; [A survey of the Boötes void](#), Astrophysical Journal 314 (Mar.1987) 493-506.
- F.G.Kopylova & A.I.Kopylov: Investigation of properties of galaxy clusters in the Hercules supercluster region, Astronomy Letters 39,1 (2013) 1-16.
- Olga A. Ladyzhenskaya: The Mathematical Theory of Viscous Incompressible Flows (2nd edition), Gordon & Breach, New York, 1969.
- Ulrich Lindner, Jaan Einasto, Maret Einasto, Wolfram Freudling, Klaus Fricke, Erik Tago: [The structure of supervoids. I. Void hierarchy in the Northern Local Supervoid](#), Astronomy and Astrophysics 301 (Sep.1995) 329-347. Related online paper on ArXiv: [astro-ph/9711046](#).
- Charles W. Misner, Kip S. Thorne, John A. Wheeler: Gravitation, Freeman 1973. QC178.M57.
- Christoph Saulder, Steffen Mieske, Eelco van Kampen, Werner W. Zeilinger: [Hubble flow variations as a test for inhomogeneous cosmology](#), Astronomy & Astrophysics 622 (Feb.2019) A83. [COmment](#) by D.L.Wiltshire, Astronomy & Astrophysics 624 (Apr.2019) A12.
- Antonia Seifert, Zachary G. Lane, Marco Galoppo, Ryan Ridden-Harper, David L. Wiltshire: [Supernovae evidence for foundational change to cosmological models](#), Monthly Notices Royal Astronomical Soc. Letters 537 (2025) L55.
- David L. Wiltshire: [Cosmic clocks, cosmic variance and cosmic averages](#), New Journal of Physics 9 (Oct.2007) #377 (66 pages).
- David L. Wiltshire: [Cosmic structure, averaging and dark energy](#), (2013-2016). Wiltshire's

description: "Based on lecture notes given to a Brazilian Cosmology School. The most comprehensive introduction I have, at a level for graduate students in cosmology, although all the material on observational tests there is a decade out of date."

David L. Wiltshire: [Average observational quantities in the timescape cosmology](#), Physical Review D 80 (Dec.2009) #123512. Wiltshire description: "Proposes a comprehensive series of observational tests" – but not my test here!

19 Dec. 2024 press release from Royal Astronomical Society: "[Dark energy 'doesn't exist' so can't be pushing 'lumpy' Universe apart – study](#)" featured maximally-prominently (as of January 2025) on D.L.Wiltshire web page, and citing him as the sole "scientific contact."