

# Deduction of the Electron Charge

Antonio Mejias<sup>1</sup>

<sup>1</sup>Territorial Polytechnic University of Mérida. (UPTM). antoniojmejiasm@gmail.com

## Abstract

In this work the charge of the electron will be deduced from Gauss's flow law.

**Keywords:** electron, Gauss's law, speed of light, electron mass, Bohr radius

## 1. Introduction

The charge of electron is considered a fundamental constant of nature, which means its value must be determined experimentally. The current value is known with very high precision due to a series of rigorous experiments. The most famous is Millikan's oil drop experiment. [1]

However, in this work it will be deduced from other more fundamental constants such as the electron radius, the speed of light and electron mass.

## 2. Gauss's Flux Law and Electron Charge

Gauss's flux law: [2,3]

$$\Phi = \iiint \nabla \cdot \vec{F} dV \quad (1)$$

Since  $\vec{F} = \omega^2 \vec{r} \mu_e = \frac{v^2}{r^2} \vec{r} \mu_e$  [4] is the centripetal force of the electron, where  $\mu_e$  is the reduced mass of the electron in the hydrogen atom,  $v$  is the velocity of the electron in the hydrogen atom, and  $r$  is the Bohr radius.

Then, from (1), the following is obtained:

$$\Phi = 4\pi v_e^2 r_b \mu_e \quad v_e = \frac{\hbar}{\mu_e r_b}$$

From here, we get:

$$\frac{\Phi}{4\pi} = \frac{\hbar^2}{r_b \mu_e} \quad (2)$$

## 3. Electron Charge in ues

From (2):

$$\sqrt{\frac{\Phi}{4\pi}} = e = \pm \hbar \sqrt{\frac{1}{r_b \mu_e}} \quad (3)$$

Given:

$$\hbar = 1.055 \times 10^{-27} \text{ erg} \cdot \text{s}$$

$$r_b = 5.3 \times 10^{-9} \text{ cm}$$

$$\mu_e = 9.11 \times 10^{-28} \text{ gr}$$

Substituting these values into (3):

$$e = \pm 4.8 \times 10^{-10} \text{ ues} [5]$$

## 4. Origin of the Electric Permittivity $\epsilon_0$ and Magnetic Permeability $\mu_0$ Constants

As these constants are related to the deduction of electrical charge in Coulombs, it is incorrect to use them without knowing their true origin.

Speed of light squared is  $c^2 = 9 \times 10^{16} \frac{m^2}{s^2}$ . The trick is to multiply and divide this quantity by  $4\pi$  and by the units  $\frac{N}{A^2}$ , and then distribute them in a special way as shown below: [6]

$$c^2 = \frac{4\pi}{4\pi} \times 9 \times 10^{16} \frac{1}{\frac{s^2}{m^2}} = \frac{1}{\left(\frac{4\pi}{10^7}\right) \left(\frac{1}{4\pi \times 9 \times 10^9}\right) \left(\frac{N}{A^2}\right) \left(\frac{A^2}{N}\right)}$$

Ampere (A)  $\times$  second (s) = Coulomb (C)

$$c^2 = \frac{1}{\left(\frac{4\pi}{10^7}\right) \left(\frac{N}{A^2}\right) \left(\frac{1}{4\pi \times 9 \times 10^9}\right) \frac{C^2}{N \cdot m^2}}$$

$$\mu_0 = \frac{4\pi}{10^7} \frac{N}{A^2}$$

$$\epsilon_0 = \left(\frac{1}{4\pi \times 9 \times 10^9}\right) \frac{C^2}{N \cdot m^2} = 8.84 \times 10^{-12} \frac{C^2}{N \cdot m^2}$$

Therefore:

$$c^2 = \frac{1}{\mu_0 \epsilon_0}$$

## 5. Electron Charge in Coulombs

Also from formula (2), multiplying both sides by  $\epsilon_0$  but in MKS units, we obtain the following:

$$\Phi \epsilon_0 = \frac{4\pi \hbar^2 \epsilon_0}{r_b \mu_e}$$
$$\sqrt{\Phi \epsilon_0} = e = \pm \hbar \sqrt{\frac{4\pi \epsilon_0}{r_b \mu_e}} \quad (4)$$

Using:

$$\epsilon_0 = 8.84 \times 10^{-12} \frac{C^2}{N \cdot m^2}$$

$$\mu_e = 9.11 \times 10^{-31} kg$$

$$r_b = 5.3 \times 10^{-11} m$$

$$\hbar = 1.055 \times 10^{-34} J \cdot s$$

Substituting these values into (4):

$$e = \pm 1.6 \times 10^{-19} C$$

## 6. Conclusions

The charge of electron is not a fundamental constant it is the result of the combination of other fundamental constants, such as the Bohr radius, the mass of the electron, and the reduced Planck constant.

The sign of the charge arises from the mathematical nature of the square root, naturally allowing for the existence of electrons, positrons.

Likewise, Magnetic Permeability constants  $\mu_0$  and Electric Permittivity constants  $\epsilon_0$  are not fundamental constants but rather mathematical arrangements.

## 7. References

1. MILIKAN, ROBERT ANDREWS. On the elementary electrical charge and the Avogadro constant. *Physical Review*, 1913, vol. 2, no 2, p. 109.
2. MURRAY R. Spiegel. *Vector Analysis*, McGraw Hill. ISBN 9684228805.
3. RAYMOND A. Serway. *Physics Volume II*, 4th ed. Mexico: McGraw Hill. ISBN 0030156548.
4. FRANCIS W. Sears, MARK W Zemansky, HUGH D. Young, 1986, *University Physics*, 6th ed. Mexico: Addison-Wesley Iberoamericana. ISBN 0201640139.
5. CHARLES KITTEL, WALTER D. Knight, MALVIN A. Ruderman, 1968, *Mechanics Volume 1*.
6. "The unification of the 4 forces and more." (September 26, 2021). Deduction of the magnetic permeability and electric permittivity constant [Video]. YouTube. <http://www.youtube.com/watch?v=u8WvcUnWTtQ>.

