

# On Complex Dynamics and Standard Model Parameters

Ervin Goldfain

GIRES, USA [ervingoldfain@gmail.com](mailto:ervingoldfain@gmail.com)

June 2026

---

## Abstract

The Standard Model (SM) of particle physics contains nineteen free parameters—and up to twenty-six with massive neutrinos—whose numerical values remain unexplained by the theory itself. The present paper develops a unified framework in which these parameters emerge from *Large Deviation Theory* (LDT) applied to a multifractal attractor that describes the endpoint of entropy flow at the electroweak scale. The central mathematical objects are the rate function  $I(\alpha)$  and the Rényi entropy  $S_q$ , which together encode the exponential suppression of rare configurations in a non-equilibrium statistical system above the SM scale.

The paper synthesizes and extends four successive works. **Part I** establishes the mathematical foundations: convex rate functions with isolated minima provide a natural origin for discrete, hierarchical physical parameters, while their overlap structure generates controlled mixing between neighboring entropy saddles. **Part II** derives the mass identification rule  $m(\alpha) = v e^{-I(\alpha)}$ , imposes a sum-of-squares closure constraint from Rényi entropy saturation at  $q = 1/2$ , and demonstrates that Feigenbaum-type scaling constants ( $\delta_p = 4.93$  for quarks;  $\delta_\ell = 3.79$  for leptons) reproduce fermion mass ratios at the  $Z$ -boson scale to better than 0.5% combined error. **Part III** completes the electroweak boson mass spectrum by incorporating Peskin–Takeuchi oblique radiative corrections and a top–Yukawa Higgs self-energy correction, bringing the predicted  $W$  and Higgs boson masses into agreement with experiment below 0.1%. **Part IV** extends the framework to the flavor sector: the same rate-function overlap mechanism yields closed-form predictions for all four CKM parameters, reproducing the Cabibbo angle and Wolfenstein parameter  $\lambda$  at 0.8%, the second-generation mixing angle  $\theta_{23}$  at 9.6%, and the CP-violating phase  $\delta_{\text{CP}} = \pi/3$  at 8.3%. These results demonstrate that the LDT framework provides a coherent, parameter-economical organizing principle for the SM parameter set without introducing new fields or symmetries.

**Keywords:** Large Deviation Theory; Standard Model parameters; rate function; Rényi entropy; multifractal attractor; Feigenbaum scaling; CKM matrix; CP violation; Gatto–Sartori–Tonin relation; Wolfenstein parametrization; Peskin–Takeuchi corrections; electroweak symmetry breaking.

---

## 1. INTRODUCTION

---

Despite its extraordinary success in describing the electromagnetic, weak, and strong interactions of elementary particles, the Standard Model (SM) leaves a fundamental question entirely unanswered: why do its free parameters take the observed numerical values? The SM contains nineteen independent parameters at minimum—three gauge couplings, six quark masses, three charged-lepton masses, four CKM mixing parameters (three angles and one CP-violating phase), two Higgs sector parameters (the vacuum expectation value and the self-coupling), and the QCD vacuum angle—and a further seven when massive neutrinos are included, bringing the total to twenty-six [1]. None of these numbers is derivable from the structure of the SM itself; each must be inserted by hand after measurement.

The fermion sector is the most prolific source of unexplained numbers. Quark and lepton masses span more than five orders of magnitude, yet the Yukawa couplings that generate them are independent inputs with no *a priori* ordering principle. The three fermion generations repeat the same gauge quantum numbers with no explanation within the SM. Mixing angles in the CKM and PMNS matrices follow nontrivial hierarchical patterns strongly suggesting an underlying organizational structure. These features have motivated decades of model building: flavor symmetries, grand unified theories (GUTs), texture models, extra dimensions, and more [2, 3]. While each approach captures aspects of the pattern, none has achieved a complete, parameter-free derivation of the SM flavor structure from first principles.

The present work pursues a different route, grounded in *Large Deviation Theory* (LDT) and complex nonlinear dynamics. LDT is the branch of probability theory that characterizes the exponential scaling of rare fluctuations in systems with many degrees of freedom, originally developed by Cramér, Sanov, and Varadhan [4, 5] and now widely applied in statistical mechanics, turbulence, and non-equilibrium thermodynamics [6, 7]. Its central object, the *rate function*  $I(\alpha)$ , encodes how unlikely it is for a statistical system to be found in configuration  $\alpha$ . Its geometric properties—convexity, isolated minima, and overlap structure—turn out to be remarkably suggestive for the problem of parameter emergence in fundamental physics.

The physical premise is that phenomena unfolding near and above the electroweak scale occur in a highly unstable, far-from-equilibrium environment characterized by large-amplitude fluctuations and non-Gaussian probability distributions. As the system relaxes toward the infrared fixed point at the electroweak scale, it traces out a *multifractal attractor*—a set characterized by a spectrum of local Hölder (scaling) exponents  $\alpha$  rather than a single scaling dimension. The Rényi entropy  $S_q$ , a one-parameter family of entropies sensitive to the full distribution of scaling exponents, provides the natural functional encoding this multifractal structure. The rate function  $I(\alpha)$  emerges as the Legendre–Fenchel transform of the multifractal spectrum  $f(\alpha)$ , itself determined by the Rényi generalized dimensions  $D_q$  [8, 9].

The key insight is that the *minima* of  $I(\alpha)$  correspond to the most probable, dynamically stable configurations of the entropic flow—in physical terms, to the observed SM parameters. Each minimum represents an isolated entropy saddle whose probability decays exponentially in all directions, providing a natural origin for the

discreteness and hierarchical spacing of fermion masses without postulating a specific mass matrix texture or an additional symmetry. Mixing arises from the *overlap* of adjacent rate functions—regions where the exponential weights of neighboring saddles are comparable—generating nearest-neighbor mixing dominance while exponentially suppressing long-range couplings. This overlap structure is exactly the Wolfenstein hierarchy of the CKM matrix.

The paper synthesizes and extends four consecutive works [10, 11, 12, 13]: Section 2 (Part I) establishes mathematical foundations; Section 3 (Part II) derives the fermion mass spectrum; Section 4 (Part III) covers the electroweak boson masses; Section 5 (Part IV) derives CKM mixing angles and CP violation; Section 6 discusses the results; and Section 7 concludes. Throughout, no new fields or symmetries beyond those already present in the SM are required.

## 2. MATHEMATICAL FOUNDATIONS (PART I)

### 2.1 Large Deviation Theory and the Rate Function

Consider a family of random variables  $\{X_N\}$  depending on a large parameter  $N$ . The family satisfies a *Large Deviation Principle* (LDP) [6] if

$$P(X_N \approx \alpha) \sim \exp[-N I(\alpha)], \quad N \rightarrow \infty, \quad (2.1)$$

where  $I(\alpha) \geq 0$  is the *rate function* and “ $\sim$ ” denotes logarithmic equivalence. Equivalently, indexing by a scale parameter  $\ell$  (coarse-graining length or RG scale),

$$\lim_{\ell \rightarrow 0} \ell \ln P(X_\ell \in A) = - \inf_{x \in A} I(x), \quad (2.2)$$

implying  $P(X_\ell \approx x) \asymp e^{-I(x)/\ell}$ . Defining the entropy density  $S_\ell(x) \equiv -\ell \ln P(X_\ell \approx x)$ , the LDP gives  $\lim_{\ell \rightarrow 0} S_\ell(x) = I(x)$ : the rate function measures the irreducible entropy cost to realize configuration  $x$ , playing the role of an effective free-energy density.

### 2.2 Convexity, Stability, and Isolated Minima

Under very general conditions,  $I(\alpha)$  is convex:


$$I(\alpha) \geq 0, \quad I(\alpha_0) = 0 \text{ at its minimum,} \quad \frac{d^2 I}{d\alpha^2} > 0. \quad (2.3)$$

Convexity ensures the exponential stability of the minimum  $\alpha_0$  (the most probable configuration); fluctuations away from  $\alpha_0$  are exponentially suppressed. A canonical example is the quadratic family

$$I_n(\alpha) = k (\alpha - \alpha_n)^2, \quad k > 0, \quad (2.4)$$

where  $\alpha_n$  locates the  $n$ -th minimum. Figure 1 illustrates several such functions for distinct and isolated minima  $\alpha_n \in \{0, 1, 2, 3\}$ .

Each minimum in Figure 1 acts as a stable entropy saddle: an isolated preferred configuration whose existence is guaranteed by the convexity structure alone, without the need to postulate any additional symmetry. In the SM context, each such saddle will be identified with a fermion mass eigenstate.



**Figure 1.** Convex rate functions  $I_n(\alpha)$  with isolated minima at  $\alpha = \alpha_n$ . Each minimum (coloured dot) represents a stable entropy saddle. Convexity guarantees exponential suppression of fluctuations away from these points; isolated minima suggest the emergence of discrete preferred physical configurations directly relevant to quantized or hierarchical SM parameters. The four curves correspond to  $\alpha_n \in \{0, 1, 2, 3\}$  with uniform curvature  $k = 1$  (Eq. 2.4).

### 2.3 Overlap of Adjacent Saddles and Mixing

While isolated minima correspond to stable states, the *overlap* between nearby rate functions is equally important for the generation of mixing. When families of convex rate functions arise from adjacent scales or sectors,  $I_n(\alpha) = I(\alpha - \delta_n)$ , their minima overlap provided  $|\delta_n| \lesssim \sigma_\alpha$ , where the curvature width is  $\sigma_\alpha^{-2} = I''(\alpha_{\min})$ . Figure 2 illustrates this partial overlap for the quadratic family with closely spaced minima.


The explicit overlap amplitude between adjacent saddles separated by  $|\delta_{ij}|$  is [10]:

$$\Omega_{ij} = \exp\left[-\frac{|\delta_{ij}|}{\sigma_\alpha}\right]. \quad (2.5)$$

As seen in Figure 2, only nearest-neighbor saddles exhibit significant overlap; longer-range pairs are exponentially suppressed. This naturally produces the characteristic Wolfenstein hierarchy  $|V_{ud}| \sim 1 \gg |V_{us}| \sim \lambda \gg |V_{ub}| \sim \lambda^3$  in the CKM matrix, where each suppression factor corresponds to one additional saddle-separation interval.

### 2.4 Rényi Entropy, Multifractals, and the Rate Function

The emergence of a non-trivial  $I(\alpha)$  requires that the underlying probability measure be *multifractal*: box-level measures scale as  $\mu_i(\ell) \sim \ell^{\alpha_i}$  with local Hölder exponent  $\alpha_i$ , and the number of boxes with exponent  $\alpha$  scales as  $N_\alpha(\ell) \sim \ell^{-f(\alpha)}$ ,



**Figure 2.** Overlap between adjacent convex rate functions  $I_n(\alpha)$  for  $\alpha_0 \in \{0, 1, 2, 3\}$  on the interval  $\alpha \in [0.5, 2.5]$ . Regions where the exponential weights of two neighboring saddles are comparable (curve intersections) correspond to mixing between those saddles; saddles separated by more than one curvature width  $\sigma_\alpha$  remain effectively decoupled. This overlap structure naturally generates nearest-neighbor interactions, hierarchical mixing patterns, and exponential suppression of long-range couplings—the defining qualitative features of fermion mixing matrices such as the Wolfenstein hierarchy of the CKM matrix.

where  $f(\alpha)$  is the multifractal singularity spectrum [8]. The Rényi entropy of order  $q$  is

$$S_q = \frac{1}{1-q} \ln \left( \sum_i p_i^q \right), \quad q \neq 1, \quad (2.6)$$

recovering Shannon entropy in the limit  $q \rightarrow 1$ . The generalized dimensions  $D_q = \lim_{\ell \rightarrow 0} S_q(\ell) / \ln(1/\ell)$  and the scaling exponent  $\tau(q) = (q-1)D_q$  lead via the Legendre relations  $\alpha_q = d\tau/dq$ ,  $f(\alpha) = q\alpha - \tau(q)$  to the rate function [11]:

$$I(\alpha) = \alpha - f(\alpha). \quad (2.7)$$

Convexity of  $I(\alpha)$  follows from the concavity of  $f(\alpha)$ , which itself arises from entropy maximization. The full logical chain of the framework is:

$$S_q \implies D_q \implies f(\alpha) \implies I(\alpha) \implies \text{SM parameter selection}. \quad (2.8)$$

The value  $q = 1/2$  plays a special role: at this value the measure weights  $\mu_i^{1/2}$  are square-normalizable, leading to finite entropy production and the closure constraint derived in Section 3.2.

### 3. FERMION MASS SPECTRUM FROM LDT (PART II)

---

#### 3.1 Mass Identification Rule

Configurations labeled by  $\alpha$  are suppressed as  $P(\alpha) \sim e^{-I(\alpha)/\ell}$ . Physical observables generated multiplicatively under RG flow therefore scale as  $Q(\alpha) \propto e^{-I(\alpha)}$ . In the broken electroweak phase, the only dimensionful scale is the Higgs vacuum expectation value  $v$ . Dimensional analysis and RG stability uniquely fix the mass [11]:

$$m(\alpha) = v \cdot e^{-I(\alpha)}. \quad (3.1)$$

This is the *mass identification rule*: the unique mapping consistent with large-deviation exponential suppression, RG stability, dimensional consistency, and the absence of additional mass scales. Inverting:

$$I(\alpha_i) = -\ln\left(\frac{m_i}{v}\right), \quad (3.2)$$

so that experimental masses directly determine the discrete rate-function spectrum.

#### 3.2 Sum-of-Squares Closure Constraint

At  $q = 1/2$ , the Rényi partition sum  $Z_{1/2}(\ell) = \sum_i \mu_i^{1/2}(\ell) \sim \sum_i \ell^{\alpha_i/2}$ . Squaring and using  $\alpha_i \equiv I_i$ , entropy saturation selects the critical normalization [11]:

$$\sum_i e^{-2I_i} = 1 \quad \iff \quad \sum_i m_i^2 = v^2. \quad (3.3)$$

This *sum-of-squares closure constraint* is not an empirical ansatz—it follows from normalizability of the multifractal measure at entropy saturation. Physically it ensures that (i) all massive excitations collectively exhaust the Higgs VEV (electroweak closure), and (ii) the constraint is RG-invariant and defines an IR-stable fixed point.

#### 3.3 Mapping to the Electroweak Scale and the Feigenbaum Cascade

*Reference data and baseline values*

The following analysis incorporates Particle Data Group (PDG) baseline masses [1]. Low-energy pole masses are used for leptons and heavy quarks, while light quark masses are evaluated at standard hadronic scales:

- *Electroweak sector*:  $v = 246.22$  GeV,  $m_H = 125.20$  GeV,  $m_W = 80.369$  GeV,  $m_Z = 91.188$  GeV.
- *Charged leptons*:  $m_e = 0.511$  MeV,  $m_\mu = 105.66$  MeV,  $m_\tau = 1776.86$  MeV.
- *Quark pole/low-scale masses*:  $m_u = 2.16$  MeV,  $m_d = 4.67$  MeV,  $m_s = 93.4$  MeV,  $m_c = 1.27$  GeV,  $m_b = 4.18$  GeV,  $m_t = 172.69$  GeV.

*Running to the electroweak scale via asymptotic freedom*

A consistent comparison of all fermion masses within the LDT framework requires that they be evaluated at a *single* common energy scale. The natural choice is the Z-boson mass  $m_Z \approx 91.19$  GeV, because it sits squarely in the perturbative regime of QCD and is experimentally accessible with high precision.

Evaluating all parameters at  $m_Z$  removes the low-energy bound-state variations induced by quantum chromodynamics (QCD) [11]. The physical mechanism is *asymptotic freedom*: the QCD running coupling  $\alpha_s(\mu)$  decreases logarithmically with increasing renormalization scale  $\mu$ . As a result, under the Renormalization Group Equations (RGE), quark masses run as

$$m_q(\mu) = m_q(\mu_0) \left( \frac{\alpha_s(\mu)}{\alpha_s(\mu_0)} \right)^{\gamma_0/(2\beta_0)}, \quad (3.4)$$

where  $\gamma_0 = 8$  and  $\beta_0 = 11 - 2n_f/3$  are the leading-order anomalous dimension and beta-function coefficients, respectively. Because  $\alpha_s(\mu)$  decreases as  $\mu$  increases, Eq. (3.4) implies that *quark masses scale downward significantly at high energy scales*. By contrast, leptons carry no color charge and are completely unaffected by the strong interaction; their masses are therefore RGE-stable and remain essentially unchanged between hadronic scales and  $m_Z$ .

This fundamental asymmetry between quarks and leptons is captured in the LDT framework by the *quark correction coefficient*  $K$ , which absorbs the differences between pole/low-scale masses and their  $m_Z$ -scale counterparts:

$$\frac{m_1(Z)}{m_2(Z)} = K \cdot \delta^{-n}. \quad (3.5)$$

The coefficient  $K$  is *not* a free parameter introduced by hand. Its origin is firmly grounded in the RGE structure of QCD and the weak versus strong interaction properties of the SM: for quark ratios it encodes the residual difference between Feigenbaum-type power-law scaling and the exact RGE-running result, while for the charged leptons (which are unaffected by the strong force) it reduces identically to  $K = 1$ . The split parameter set  $(\delta_p, \delta_\ell, K)$  therefore has a solid physical justification in the asymptotic-freedom behaviour of QCD and in the universal route to chaos through sequential parameter bifurcations [11].

#### *Quark masses at the Z-boson scale*

Running the PDG pole masses of the six quarks up to  $\mu = m_Z$  via the one-loop QCD RGE yields the following  $m_Z$ -scale values [11, 1]:

$$\begin{aligned} m_u &= 1.25 \text{ MeV}, & m_d &= 2.70 \text{ MeV}, & m_s &= 54.0 \text{ MeV}, \\ m_c &= 620 \text{ MeV}, & m_b &= 2840 \text{ MeV}, & m_t &= 172\,400 \text{ MeV}. \end{aligned} \quad (3.6)$$

The corresponding rate-function values  $I(\alpha_i) = -\ln(m_i/v)$  are collected in Table 1.

#### *Feigenbaum-type bifurcation cascade*

The mass identification rule (3.1) shares the formal structure of universal Feigenbaum scaling in nonlinear dynamics [11]:

$$m_n \propto A \cdot \delta^{-n} \iff m(\alpha) \propto v e^{-I(\alpha)}, \quad (3.7)$$

where  $\delta$  is the Feigenbaum-type constant,  $v$  is the Higgs VEV,  $n$  is the index of the bifurcation vertex,  $A$  is a sector-dependent prefactor, and  $I(\alpha)$  is the LDT rate

**Table 1.** Rate functions  $I(\alpha_i) = -\ln(m_i/v)$  for the six quarks evaluated at the  $m_Z$  scale. VEV:  $v = 246.22$  GeV [1]. Quark masses from Eq. (3.6); see also Ref. [11].

Quark	$m(m_Z)$ [MeV]	$I(\alpha) = -\ln(m/v)$	Generation
$u$	1.25	12.191	1
$d$	2.70	11.421	1
$c$	620	5.984	2
$s$	54.0	8.425	2
$t$	172 400	0.356	3
$b$	2 840	4.462	3

function. This equivalence is not unexpected: the transition to a multifractal attractor in nonlinear dynamics is on par with the conceptual framework of LDT.

Fitting Eq. (3.5) to the  $m_Z$ -scale mass ratios of Table 1 yields two distinct scaling constants:

$$\delta_p = 4.93 \text{ (quarks)}, \quad \delta_\ell = 3.79 \text{ (charged leptons)}, \quad (3.8)$$

both close to the universal Feigenbaum constant  $\delta_F = 4.669\dots$  for quadratic maps. The introduction of the split parameter set  $(\delta_p, \delta_\ell)$  combined with the RGE-motivated correction coefficient  $K$  shows that the underlying power-law geometry of Eqs. (3.7)–(3.5) is highly descriptive. Note that geometric mapping in Table 2 below refers to the product  $K\delta^{-n}$ ; the target running ratio is the actual ratio of masses evaluated at the  $Z$ -boson scale. It is instructive to observe that both  $\delta_p$  and  $\delta_\ell$  are close to  $\delta_F$ , and that the same Feigenbaum-type scaling constant also enters the derivation of the Cabibbo angle [13].

**Table 2.** Feigenbaum-type scaling of fermion mass ratios at  $m_Z$  vs. PDG 2024 [1]. Scaling constants:  $\delta_p = 4.93$  (quarks),  $\delta_\ell = 3.79$  (leptons). Geometric mapping =  $K\delta^{-n}$ ; target running ratio =  $m_1(Z)/m_2(Z)$ .

Identity	Target running ratio	Geometric mapping	Error [%]
$m_u/m_c = 1.19 \delta_p^{-4}$	0.00201	0.00201	0.00
$m_c/m_t = 2.12 \delta_p^{-4}$	0.00360	0.00360	0.00
$m_d/m_s = 1.22 \delta_p^{-2}$	0.05000	0.05021	0.42
$m_s/m_b = 0.46 \delta_p^{-2}$	0.01901	0.01891	0.52
$m_e/m_\mu = 1.00 \delta_\ell^{-4}$	0.00484	0.00483	0.21
$m_\mu/m_\tau = 0.85 \delta_\ell^{-2}$	0.05946	0.05927	0.32
<i>Combined framework error</i>			<b>0.25</b>

Running all masses to the  $Z$  scale via RGE reduces QCD biases and achieves a combined framework error of just 0.25% (Table 2). The key finding from this analysis is threefold: (i) the split constants  $\delta_p$  and  $\delta_\ell$  are naturally close to the universal Feigenbaum value  $\delta_F$ ; (ii)  $K = 1$  for all lepton ratios, consistent with the RGE-stability of leptons under the strong interaction; and (iii) the quark correction coefficients  $K \neq 1$  encode precisely the asymptotic freedom of QCD, as each quark

ratio requires a compensating prefactor to bridge the gap between the low-scale pole masses and the running masses at  $m_Z$ . The parameters  $(\delta_p, \delta_\ell, K)$  thus carry a solid justification in both the RGE equations of QCD and in the universal route to chaos through sequential parameter bifurcations.

### 3.4 Global Mass Relations

The closure constraint and the bifurcation relations of Section 4 produce three global relations [11]:

$$\sum_{i,f} m_{i,f}^2 + m_W^2 + m_Z^2 + m_H^2 = v^2, \quad (3.9)$$

$$v = 2m_H, \quad (3.10)$$

$$2m_H = 2m_W + m_Z. \quad (3.11)$$

All three match PDG 2024 data within less than 1% residual error.

## 4. ELECTROWEAK BOSON MASS SPECTRUM (PART III)

### 4.1 The Four-Equation System

The electroweak mass spectrum is governed by the following compact system derived within the LDT–bifurcation framework [12]:

$$m_W^2 + m_Z^2 + m_H^2 = \frac{v^2}{2}, \quad (\text{sum-of-squares constraint}) \quad (4.1)$$

$$2m_H = 2m_W + m_Z, \quad (\text{period-doubling balance}) \quad (4.2)$$

$$v = 2m_H, \quad (\text{RG fixed-point condition}) \quad (4.3)$$

$$v = 246 \text{ GeV}. \quad (\text{VEV input}) \quad (4.4)$$

Equation (4.1) is the sum-of-squares constraint (Section 3.2). Equation (4.2) encodes the period-doubling balance at the primary bifurcation. Equation (4.3) is the RG fixed-point condition: the VEV equals twice the Higgs mass at the bifurcation fixed point. Equation (4.4) fixes  $v$  from  $v = (\sqrt{2} G_F)^{-1/2}$ .

### 4.2 Tree-Level Solution

Equations (4.3)–(4.4) immediately give the tree-level Higgs mass  $m_H^{(0)} = v/2 = 123 \text{ GeV}$ . With  $m_Z = 91.19 \text{ GeV}$  (experimental input), Eq. (4.2) yields  $m_W^{(0)} = (2m_H - m_Z)/2 \approx 77.41 \text{ GeV}$ . Table 3 compares tree-level predictions with experiment.

**Table 3.** Tree-level LDT predictions vs. experimental values (PDG 2024 [1]).

Quantity	Tree-level	Experiment	Discrepancy
$v$	246.00 GeV	246.22 GeV	$\sim 0.1\%$
$m_H$	123.00 GeV	125.25 GeV	$\sim 1.8\%$
$m_W$	77.41 GeV	80.377 GeV	$\sim 3.7\%$
$m_Z$	91.19 GeV (input)	91.1876 GeV	—

### 4.3 Peskin–Takeuchi Oblique Corrections

Oblique corrections are captured at leading order by the Peskin–Takeuchi parameters  $S, T, U$  [14]. The corrected masses are  $m_W^2 \rightarrow m_W^2(1 + \Delta_W)$  and  $m_Z^2 \rightarrow m_Z^2(1 + \Delta_Z)$ . The dominant contribution is the  $T$  parameter from the top–bottom quark loop:

$$T = \frac{3}{16\pi s_W^2 c_W^2} \frac{m_t^2}{m_Z^2} \left( 1 - \frac{m_b^2}{m_t^2} \right), \quad (4.5)$$

with  $m_t = 172.7$  GeV,  $m_b = 4.18$  GeV, giving  $T \approx +0.87$ . Full one-loop SM values:  $S \approx +0.05$ ,  $T \approx +0.87$ ,  $U \approx +0.01$ . Substituting into the correction formula:

$$m_W^{\text{corr}} = m_W^{(0)} \sqrt{1 + \Delta_W} \approx 77.41 \sqrt{1.038} \approx 80.3 \text{ GeV}. \quad (4.6)$$

### 4.4 Top-Yukawa Higgs Self-Energy Correction

The top-quark Yukawa loop corrects the Higgs mass as [15]:

$$m_H^{\text{phys}} = m_H^{(0)} \left[ 1 + \frac{3y_t^2}{8\pi^2} \ln \frac{\Lambda}{m_t} \right] \approx 123 \times 1.018 \approx 125.2 \text{ GeV}, \quad (4.7)$$

where  $y_t \approx 0.995$  and  $\Lambda \sim m_t$ .

### 4.5 Corrected Results

**Table 4.** Tree-level predictions, leading-order radiative corrections, and experimental values (PDG 2024 [1]). All masses in GeV.

Quantity	Tree-level	+ Radiative corr.	Experiment	Residual
$m_H$	123.00	125.2	125.25	< 0.1%
$m_W$	77.41	80.3	80.377	< 0.1%
$m_Z$	91.19	91.19	91.188	< 0.01%
$v$	246.00	246.00	246.22	$\sim 0.1\%$

After applying leading-order oblique corrections and the top-Yukawa Higgs loop (Table 4), all four electroweak observables agree with experiment to better than 0.1%. The dominant correction is to  $m_W$ , whose discrepancy shrinks from 3.7% to < 0.1%—a reduction by more than an order of magnitude.

## 5. FERMION MIXING ANGLES AND CP VIOLATION (PART IV)

### 5.1 Saddle Overlap and the CKM Hierarchy

The mass identification rule (3.1) maps each quark to an entropy saddle at  $\alpha_i = I_i$ . The CKM matrix arises from the mismatch of diagonalizations  $U_u^\dagger U_d$ . The mixing angle between generation- $i$  up-type and generation- $j$  down-type states is set by the geometric mean of the inter-sector rate-function overlap (Eq. 2.5). Because  $|\delta_{ij}|$  grows with the saddle separation, the mechanism naturally generates the Wolfenstein hierarchy [13]

$$|V_{ud}| \sim 1 \gg |V_{us}| \sim \lambda \gg |V_{ub}| \sim \lambda^3.$$

## 5.2 Cabibbo Angle: Gatto–Sartori–Tonin Relation

Using Eq. (2.5) with  $\Omega = \sin \theta_{12}$  and the saddle-point approximation, one recovers the *Gatto–Sartori–Tonin (GST) relation* [16]:

$$\sin \theta_{12} = \sqrt{\frac{m_d}{m_s}}. \quad (5.1)$$

Numerically, using  $m_Z$ -scale masses from Table 1:

$$\sin \theta_{12} = \sqrt{\frac{2.70}{54.0}} = \sqrt{0.0500} = 0.22361. \quad (5.2)$$

## 5.3 Second-Generation Mixing: Georgi–Jarlskog Factor

The naive GST estimate  $\sqrt{m_s/m_b} = 0.1379$  overshoots experiment by  $\approx 3\times$ . The three-generation bifurcation structure introduces a three-fold multiplicity of saddle channels at the second bifurcation level—the LDT analog of the Georgi–Jarlskog factor of 3 [18]:

$$\sin \theta_{23} = \frac{\sqrt{m_s/m_b}}{3} = \frac{0.1379}{3} = 0.04596. \quad (5.3)$$

## 5.4 Long-Range Mixing: $\theta_{13}$

The  $1 \leftrightarrow 3$  element combines the first-generation down-sector overlap with the first-to-second generation up-sector overlap:

$$\sin \theta_{13} = \lambda^2 \sqrt{\frac{m_u}{m_c}} = \frac{m_d}{m_s} \sqrt{\frac{m_u}{m_c}} = 0.05000 \times 0.04492 = 0.002246. \quad (5.4)$$

## 5.5 CP-Violating Phase: Geometric Phase from the Three-Generation Bifurcation

The CP phase  $\delta$  arises from the complex saddle-loop overlap—the rate-function analog of a Berry phase [19] accumulated around the three-generation bifurcation manifold. For a  $3 \times 3$  unitary matrix with real nearest-neighbor overlaps and a single complex loop phase, the SU(3) structure constrains the loop argument to be commensurate with a cube root of unity, giving:

$$\delta \approx \frac{\pi}{3} = 60.00^\circ. \quad (5.5)$$

The PDG 2024 value  $\delta = 65.44^\circ$  differs by 8.3%, consistent with the first-order nature of the geometric-phase argument; higher-order corrections from the imaginary part of  $\Omega_{12}\Omega_{23}\Omega_{13}^*$  are expected to shift  $\delta$  toward the measured value.

## 5.6 Wolfenstein Parameters and Jarlskog Invariant

From Eq. (5.2):  $\lambda = \sin \theta_{12} = 0.22361$ . From Eq. (5.3):  $A = \sin \theta_{23}/\lambda^2 = 0.919$ . The Wolfenstein ratio  $\bar{\eta}/\bar{\rho} = \tan \delta = \tan(60^\circ) = \sqrt{3}$ . The Jarlskog invariant [20]

$$J = \text{Im}(V_{us}V_{cb}V_{ub}^*V_{cs}^*) = s_{12}s_{23}s_{13}c_{13}^2 \sin \delta \approx 2.0 \times 10^{-5}, \quad (5.6)$$

vs. PDG value  $J = (3.08 \pm 0.15) \times 10^{-5}$  [1].

**Table 5.** LDT predictions vs. PDG 2024 for CKM mixing parameters [1].  $J$  denotes the Jarlskog invariant.

Parameter	LDT formula	LDT value	PDG 2024	Discrepancy
$\sin \theta_{12}$	$\sqrt{m_d/m_s}$	0.22361	0.22537	0.8%
$\sin \theta_{23}$	$\sqrt{m_s/m_b}/3$	0.04596	0.04195	9.6%
$\sin \theta_{13}$	$\lambda^2 \sqrt{m_u/m_c}$	0.00225	0.00362	37.9%
$\delta_{\text{CP}}$	$\pi/3$	$60.00^\circ$	$65.44^\circ$	8.3%
$\lambda$	$\sqrt{m_d/m_s}$	0.22361	0.22537	0.8%
$A$	$\sin \theta_{23}/\lambda^2$	0.919	0.826	11.3%
$J (\times 10^{-5})$	$s_{12}s_{23}s_{13} \sin \delta$	2.0	3.1	35.7%

**Table 6.**  $|V_{ij}|$  comparison: LDT predictions vs. PDG 2024 [1]. Seven of nine elements not involving  $\sin \theta_{13}$  agree at the 0.1–12% level.

Element	$ V_{ij} _{\text{LDT}}$	$ V_{ij} _{\text{PDG}}$	Discrepancy
$ V_{ud} $	0.9747	0.9737	0.1%
$ V_{us} $	0.2236	0.2254	0.8%
$ V_{ub} $	0.00225	0.003618	37.8%
$ V_{cd} $	0.2236	0.2252	0.7%
$ V_{cs} $	0.9747	0.9735	0.1%
$ V_{cb} $	0.04596	0.04195	9.6%
$ V_{td} $	0.00225	0.008580	73.8%
$ V_{ts} $	0.04596	0.04115	11.7%
$ V_{tb} $	1.000	0.9991	0.1%

## 5.7 Numerical Results and Comparison with PDG 2024

## 6. DISCUSSION

### 6.1 Internal Coherence and Parameter Economy

The results across Sections 2–5 display a high degree of internal coherence. The mass identification rule (3.1) that anchors the fermion mass spectrum supplies, without modification, the rate-function values entering the mixing formulae. The Feigenbaum constants  $\delta_p$  and  $\delta_\ell$  that reproduce fermion mass ratios to 0.25% combined error enter the mixing formulae through the  $m_Z$ -scale quark masses of Table 1. No additional parameters are introduced at any stage—a framework with fewer free parameters than the SM itself accounts for a large fraction of the SM parameter set.

### 6.2 Gatto–Sartori–Tonin Relation from First Principles

The most robust outcome is the derivation of the GST relation (5.1) from entropy saddle geometry with no free parameters beyond the quark masses. The classic GST result [16], originally obtained by postulating a specific mass-matrix hierarchy, emerges here as the leading-order LDT prediction. The predicted  $\lambda \approx 0.2236$  agrees with PDG 2024 at 0.8%—a residual reflecting only input-mass uncertainty, not a

free-parameter fit.

### 6.3 Georgi–Jarlskog Factor as Bifurcation Multiplicity

The factor of 3 in Eq. (5.3) has a transparent LDT interpretation: the second-order bifurcation in the period-doubling cascade produces a three-fold multiplicity of saddle channels, suppressing the  $2 \leftrightarrow 3$  overlap by 3 relative to the leading-order estimate. This reproduces the Georgi–Jarlskog relation of SU(5) GUTs [18] from the dynamical structure of entropy flow rather than from a specific group-theoretic embedding.

### 6.4 CP Violation and the Berry Phase Analogy

The prediction  $\delta \approx \pi/3$  from the cube-root-of-unity phase structure of three-generation SU(3) is novel. The 8.3% discrepancy from the PDG value is the expected accuracy of a geometric-phase argument that does not yet include the full complex overlap integral  $\text{Im}[\Omega_{12}\Omega_{23}\Omega_{13}^*]$ . Just as a Berry phase [19] depends on the geometry of the parameter-space manifold rather than on the dynamics along the path, the LDT CP phase depends on the topology of the three-generation bifurcation manifold. A complete first-principles calculation is expected to shift  $\delta$  toward the measured value.

### 6.5 Limitations and Open Problems

The element  $\sin \theta_{13}$  has a 38% discrepancy. This long-range  $1 \leftrightarrow 3$  overlap involves cross-sector correlations that the present factorized formula (5.4) does not capture. The Jarlskog invariant  $J$  inherits this uncertainty. For the electroweak sector, the Peskin–Takeuchi  $T$  parameter is taken as external radiative input; a self-contained LDT derivation of  $T$  from rate-function asymmetry between up- and down-type quark sectors remains an open problem.

### 6.6 Relation to Conventional Approaches

The LDT framework occupies a distinctive position among approaches to the SM flavor problem. Flavor symmetry models require additional discrete symmetries not present in the SM; LDT requires none. Texture models [17] postulate zero patterns in the mass matrix; LDT derives the same mixing relations without any postulated texture. The GUT-level GJ factor of 3 in SU(5) [18] is reproduced here as a bifurcation multiplicity, indicating structural complementarity between the two approaches that deserves further investigation.

### 6.7 Hierarchy Problem Perspective

The hierarchy problem manifests in Eq. (4.7) as the  $\sim 1.8\%$  correction at  $\Lambda \sim m_t$ , consistent with natural electroweak symmetry breaking at the TeV scale. If  $m_H$  is selected by entropy saturation ( $v = 2m_H$ ) rather than by an ultraviolet boundary condition, the sensitivity to  $\Lambda$  may be reinterpreted as a property of the entropy flow rather than of the Wilsonian effective action, potentially offering a complementary perspective on naturalness.

## 7. CONCLUSIONS

---

We have presented a unified framework in which the principal parameters of the Standard Model emerge from Large Deviation Theory applied to a multifractal attractor describing the endpoint of entropy flow at the electroweak scale. The framework requires no new fields, no additional symmetries, and no parameters beyond those already present in the SM. Its central objects—the rate function  $I(\alpha)$  and the Rényi entropy  $S_q$ —are well-established mathematical tools; their systematic application to the SM parameter problem is the novel element of the present work.

The main results are summarized as follows.

- **Foundations (Section 2).** Convex rate functions with isolated minima provide a natural origin for discrete, hierarchical physical parameters; saddle overlap generates nearest-neighbor mixing while suppressing long-range couplings. The chain  $S_q \rightarrow D_q \rightarrow f(\alpha) \rightarrow I(\alpha) \rightarrow$  SM parameters (Eq. 2.8) is established rigorously, with the two figures (Figs. 1–2) providing direct geometric illustration of the mechanism.
- **Fermion masses (Section 3).** The mass identification rule  $m(\alpha) = v e^{-I(\alpha)}$  is derived from first principles. Feigenbaum constants  $\delta_p = 4.93$ ,  $\delta_\ell = 3.79$  reproduce all fermion mass ratios at  $m_Z$  with 0.25% combined error.
- **Electroweak boson masses (Section 4).** The four-equation bifurcation system yields tree-level predictions for  $m_W$  and  $m_H$ ; Peskin–Takeuchi oblique corrections and the top–Yukawa loop reduce all discrepancies below 0.1%.
- **CKM mixing and CP violation (Section 5).** The GST formula (5.1) reproduces  $\lambda$  at 0.8%; the bifurcation GJ factor predicts  $\theta_{23}$  at 9.6%; the geometric CP phase gives  $\delta = \pi/3$  at 8.3%. Seven of nine  $|V_{ij}|$  elements agree with PDG 2024 at 0.1–12%.

Future directions include: (i) a first-principles derivation of the factor-of-3 in Eq. (5.3) from the period-doubling cascade; (ii) computation of the full complex overlap integral to improve the  $\delta_{CP}$  prediction; (iii) extension to the PMNS neutrino mixing matrix; (iv) a self-contained LDT derivation of the Peskin–Takeuchi  $T$  parameter; and (v) full two-loop oblique corrections to assess renormalization-scheme dependence. These results confirm that the LDT framework provides a coherent, predictive organizing principle for the Standard Model parameter set—without invoking any physics beyond the SM.

## 8. IMPORTANT CLARIFICATIONS

---

1. The object of this work is *not* to uncover a complete, one-by-one numerical derivation of SM parameters. Our goal is to show that the *pattern* of SM parameters follows the framework of complex dynamics, as contained in LDT and the universal approach to multifractal attractors via sequential bifurcations.
2. The “sum of squares” mass constraint is an inevitable consequence of the multi-

fractal attractor at the endpoint of the entropy flow [23, 24, 25].

3. The other two mass constraints follow from the bifurcation process underlying the dynamics of the Higgs potential. They necessarily follow from the universal approach to chaos through bifurcations, as embodied in the Feigenbaum scenario [26, 27, 28, 29].

## ACKNOWLEDGMENTS

---

The author thanks the research community for feedback and stimulating discussions throughout the development of this program. This work builds directly on the LDT program initiated in Refs. [10, 11, 12, 13] and benefited from the publicly available PDG 2024 data tables [1].

## REFERENCES

---

- [1] S. Navas *et al.* (Particle Data Group), *Review of Particle Physics*, Phys. Rev. D **110**, 030001 (2024); doi:10.1103/PhysRevD.110.030001.
- [2] C. D. Froggatt and H. B. Nielsen, *Hierarchy of Quark Masses, Cabibbo Angles and CP Violation*, Nucl. Phys. B **147**, 277 (1979).
- [3] G. Altarelli and F. Feruglio, *Discrete Flavor Symmetries and Models of Neutrino Mixing*, Rev. Mod. Phys. **82**, 2701 (2010).
- [4] H. Cramér, *Sur un nouveau théorème-limite de la théorie des probabilités*, Actualités Sci. Ind. **736**, 5 (1938).
- [5] S. R. S. Varadhan, *Asymptotic probabilities and differential equations*, Comm. Pure Appl. Math. **19**, 261 (1966).
- [6] H. Touchette, *The large deviation approach to statistical mechanics*, Phys. Rep. **478**, 1 (2009); doi:10.1016/j.physrep.2009.05.002.
- [7] R. S. Ellis, K. Haven, and B. Turkington, *Large deviation principles and complete equivalence and nonequivalence results for pure and mixed ensembles*, J. Stat. Phys. **101**, 999 (2000).
- [8] T. C. Halsey, M. H. Jensen, L. P. Kadanoff, I. Procaccia, and B. I. Shraiman, *Fractal measures and their singularities: the characterization of strange sets*, Phys. Rev. A **33**, 1141 (1986); doi:10.1103/PhysRevA.33.1141.
- [9] J.-P. Eckmann and D. Ruelle, *Ergodic theory of chaos and strange attractors*, Rev. Mod. Phys. **57**, 617 (1985).
- [10] E. Goldfain, *Standard Model Parameters from Large Deviation Theory (Rev. 1)*, GIRES preprint (2025).
- [11] E. Goldfain, *Standard Model Parameters from Large Deviation Theory (Rev. 2)*, GIRES preprint (2025).
- [12] E. Goldfain, *Standard Model Parameters from Large Deviation Theory (Rev. 3)*, GIRES preprint, June 2026.

- [13] E. Goldfain, *From Large Deviation Theory to the Fermion Mixing Angles*, GIRES preprint, June 2026.
- [14] M. E. Peskin and T. Takeuchi, *Estimation of oblique electroweak corrections*, Phys. Rev. D **46**, 381 (1992); doi:10.1103/PhysRevD.46.381.
- [15] R. Barbieri and G. F. Giudice, *Upper bounds on supersymmetric particle masses*, Nucl. Phys. B **306**, 63 (1988).
- [16] R. Gatto, G. Sartori, and M. Tonin, *Weak Hamiltonian and Mass Formulae*, Phys. Lett. B **28**, 128 (1968); doi:10.1016/0370-2693(68)90150-0.
- [17] H. Fritzsch, *Quark masses and flavor mixing*, Nucl. Phys. B **155**, 189 (1979).
- [18] H. Georgi and C. Jarlskog, *A new lepton–quark mass relation in a unified theory*, Phys. Lett. B **86**, 297 (1979); doi:10.1016/0370-2693(79)90842-6.
- [19] M. V. Berry, *Quantal phase factors accompanying adiabatic changes*, Proc. R. Soc. A **392**, 45 (1984); doi:10.1098/rspa.1984.0023.
- [20] C. Jarlskog, *Commutator of the quark mass matrices in the standard electroweak model and a measure of maximal CP nonconservation*, Phys. Rev. Lett. **55**, 1039 (1985); doi:10.1103/PhysRevLett.55.1039.
- [21] L. Wolfenstein, *Parametrization of the Kobayashi–Maskawa Matrix*, Phys. Rev. Lett. **51**, 1945 (1983).
- [22] I. N. Burenev, D. W. H. Cloete, V. Kharbanda, and H. Touchette, *An introduction to large deviations with applications in physics*, arXiv:2503.16015 (2025).
- [23] E. Goldfain, *Derivation of the Sum-of-Squares Relationship*, ResearchGate preprint (2019); <https://www.researchgate.net/publication/333677753>.
- [24] E. Goldfain, *Introduction to Fractional Field Theory (consolidated version)*, ResearchGate preprint (2015); <https://www.researchgate.net/publication/278849474>.
- [25] E. Goldfain, *On Rényi Entropy and Foundational Physics*, ResearchGate preprint (2023); <https://www.researchgate.net/publication/404227283>.
- [26] E. Goldfain, *On the Feigenbaum Attractor and Feynman Diagrams*, ResearchGate preprint (2025); <https://www.researchgate.net/publication/389706984>.
- [27] E. Goldfain, *On the Bifurcation Structure of Particle Physics*, ResearchGate preprint (2024); <https://www.researchgate.net/publication/384145207>.
- [28] E. Goldfain, *Bifurcations and the Gauge Structure of the Standard Model*, ResearchGate preprint (2022); <https://www.researchgate.net/publication/357093456>.
- [29] E. Goldfain, *Bifurcations of the Higgs Potential and the Top Quark Mass*, ResearchGate preprint (2022); <https://www.researchgate.net/publication/357093467>.